PHILOSOPHICAL TRANSACTIONS,

OF THE

ROYAL SOCIETY

OF

LONDON.

VOL. LXXXI. For the Year 1791.

PART II.

LONDON,

SOLD BY PETER ELMSLY,

PRINTER TO THE ROYAL SOCIETY.

MDCCXCI.

CONTENTS

OF

VOL. LXXXI. PART II.

VIII. *ON the Rate of Travelling, as performed by Camels; and its Application, as a Scale, to the Purposes of Geography.* By James Rennell, *Esq. F. R. S.* Page 129

IX. *On Infinite Series.* By Edward Waring, *M. D. F. R. S.* Lucasian *Professor of Mathematics in the University of* Cambridge. p. 146

X. *An Account of some Appearances attending the Conversion of cast into malleable Iron. In a Letter from* Thomas Beddoes, *M. D. to Sir* Joseph Banks, *Bart. P. R. S.* p. 173

XI. *On the Decomposition of Fixed Air.* By Smithson Tennant, *Esq. F. R. S.* p. 182

XII. *A Meteorological Journal, principally relating to Atmospheric Electricity; kept at* Knightsbridge, *from the 9th of* May, 1789, *to the 8th of* May, 1790. *By Mr.* John Read; *communicated by* R. H. A. Bennet, *Esq. F. R. S.* p. 185

XIII. *Farther Experiments relating to the Decomposition of dephlogisticated and inflammable Air.* By Joseph Priestley, *LL.D. F. R. S.* p. 213

XIV. *Expe-*

XIV. *Experiments on Human Calculi. In a Letter from Mr.* Timothy Lane, *F. R. S. to* William Pitcairn, *M. D. F. R. S.* p. 223

XV. *Chermes Lacca. By* William Roxburgh, *M. D. of* Samulcotta. *Communicated by* Patrick Ruſſell, *M. D. F. R. S.* p. 229

XVI. *The Longitudes of* Dunkirk *and* Paris *from* Greenwich, *deduced from the Triangular Meaſurement in* 1787, 1788, *ſuppoſing the Earth to be an Ellipſoid. By Mr.* Iſaac Dalby; *communicated by* Charles Blagden, *M. D. Sec. R. S.* p. 236

XVII. *On the Method of determining, from the real Probabilities of Life, the Values of contingent Reverſions in which Three Lives are involved in the Survivorſhip. By Mr.* William Morgan, *F. R. S.* p. 246

XVIII. *Abſtract of a Regiſter of the Barometer, Thermometer, and Rain, at* Lyndon *in* Rutland; *by* Thomas Barker, *Eſq.*; *with the Rain in* Hampſhire *and* Surrey; *for the Year* 1790. *Communicated by* Thomas White, *Eſq. F. R. S.* p. 278

XIX. *Deſcription of a ſimple Micrometer for meaſuring ſmall Angles with the Teleſcope. By Mr.* Tiberius Cavallo, *F. R. S.* p. 283

XXI. *Experiments and Obſervations to inveſtigate the Compoſition of* James's *Powder. By* George Pearſon, *M. D. F. R. S.*; *communicated by Sir* Joſeph Banks, *Bart. P. R. S.* p. 317

XXII. *An Account of ſome chemical Experiments on* Tabaſheer. *By* James Louis Macie, *Eſq. F. R. S.* p. 368

XXIII. *A Second Paper on Hygrometry. By* J. A. De Luc, *Eſq. F. R. S.* p. 389

PHILOSOPHICAL TRANSACTIONS.

VIII. *On the Rate of Travelling, as performed by Camels; and its Application, as a Scale, to the Purposes of Geography.* By James Rennell, *Esq. F. R. S.*

Read March 17, 1791.

AMONGST the discoveries and improvements of various kinds, that may be expected from the very laudable, and, as it concerns mankind in the gross, no less humane and benevolent institution of AN ASSOCIATION FOR PROMOTING DISCOVERIES IN THE INTERIOR PARTS OF AFRICA, that of the geography of the Continent in question may be expected to make the quickest progress: since, in every kind of distant research, whether successful or otherwise, the act of enquiry alone brings materials to the geographer. But, even with every

ordinary advantage, the geography of a continent must neceſſarily be ſlow in its progreſs towards maturity; *we*, therefore, can only expect to witneſs its early infancy; and cannot, from a view of the preſent, allow ourſelves to predict what its future features may be. Let it be our care, however, to foſter it; and contribute all within our power towards its improvement and perfection.

In a caſe where there is ſo little probability, even in a long courſe of time, of obtaining many fixed points by celeſtial obſervations (though I hope that one at leaſt will be attempted in the central part of Africa, in our time), it is fortunate that the mode of travelling happens to be ſuch, as ſerves to furniſh a remarkably equal ſcale: the rate of the camel's movement appearing to be, beyond all others, the leaſt variable; whether we examine it by portions of days, or of hours. In the preſent ſtate of things, the former mode alone can be uſed; becauſe few or none of the African travellers carry watches with them: but it may be hoped, that at no very diſtant period, the time employed on the road may be obtained with ſuch a degree of exactneſs, as to furniſh the geographer with materials of a far better kind, than any of thoſe, formed on computation, that have hitherto been exhibited.

Theſe remarks occurred on the reſult of an examination, which (though for a different purpoſe) I lately made into the rate of the camel's movement on the Arabian deſert, between Aleppo, Bagdad, and Buſſorah: for it appeared to me, that if the African caravans are compoſed of the ſame kind of camels, and are governed in their motions and œconomy by the ſame circumſtances, as thoſe which croſs the Arabian deſerts; there is no ſcale, of the computed kind, that can be more applicable to the African geography, than that formed on the camel's rate

of

of travelling. I ſhall therefore detail the examples from whence I have drawn the proportions for the hours and days journey of the camel, under the two different degrees of burthen, which conſtitute what is commonly denominated the LIGHT, and the HEAVY caravan.

The routes which furniſh the above examples are determined in their horizontal, or direct diſtance, by the reſpective poſitions of ALEPPO, BAGDAD, and BUSSORAH: all of which have their latitudes and longitudes fixed by celeſtial obſervations *. Theſe routes are five in number: and although ſome of the journals that deſcribe them, contain leſs information than others, yet all of them have the time given with a ſufficient degree of preciſion, to enable me to found a general rule on. Three of theſe routes lead acroſs the GREAT DESERT, or that between Aleppo and Buſſorah; the other two are acroſs the LITTLE DESERT, or that between Aleppo and Bagdad.

The firſt of the Great Deſert routes was traced by a Mr. CARMICHAEL in 1751. The manuſcript copy of his Journal was obligingly communicated by my friend Dr. PATRICK RUSSELL; and it manifeſts a great degree of ingenuity and perſeverance in this way. The Author declares, that he was determined to keep a regiſter of the courſes by a compaſs, and to compute, comparatively, if not abſolutely, the intermediate diſtance on each courſe; by counting the ſteps or paces of the camel on which he rode, during a certain interval of time; and afterwards meaſuring a number of them on the ground. The particulars of this operation ſhall be given hereafter; and

* Aleppo, in Conn. des Temps, lat. 36° 11′, long. 37° 9′, reckoned from Greenwich.

Bagdad, by M. BEAUCHAMP, lat. 33° 22′, long. 44° 21′, Greenwich.

Buſſorah, by Capt. RITCHIE, lat. 30° 30′, long. 47° 33′, Greenwich.

although Mr. CARMICHAEL failed in the attempt to aſcertain his road diſtance by this method, yet his proceſs has furniſhed others with the means of aſcertaining the whole diſtance in the aggregate, and of proportioning the parts throughout. For, as the direct diſtance is given by the celeſtial obſervations, and a complete traverſe table by the journal, the *data* are perfect. And when the reader is informed that Mr. CARMICHAEL's whole line of bearing, *by compaſs*, between Aleppo and Buſſorah, nearly 720 Britiſh miles, coincided with the bearing line given by the celeſtial obſervations; by which it appears that the error could amount only to the mean quantity of the variation throughout, which might have been from ſix to ſeven degrees at that time (1751); he will give Mr. CARMICHAEL credit for much general accuracy. And it is not improbable, that even a conſiderable portion of the above error may have ariſen from the imperfection of his inſtrument *.

The ſecond journal was kept by Colonel CAPPER, in 1778, and was publiſhed ſeveral years ago; and the third, which contains little more than the time in detail, was communicated by my friend Mr. HUNTER, who croſſed the deſert in 1767.

The time given between Aleppo and Buſſorah, by theſe journals reſpectively, is as follows:

By Mr. CARMICHAEL	.	322 hours.
Colonel CAPPER	.	310
Mr. HUNTER	.	299½

* I find, by Mr. DRUMMOND's chart of the road between Aleppo and Antioch (1747), that the variation was then about 6 degrees weſterly. This is proved by comparing his magnetic bearing line between thoſe places, with that given by the difference of latitude In the head of the Gulf of Perſia, the variation was 7 degrees in 1785.

But

But to ſhew that this difference aroſe chiefly from the variations in the route acroſs the CHALDEAN DESERT (between Mesjid Ali and Buſſorah; ſee the annexed ſketch, Tab. III.), I ſhall proceed, firſt, to explain that part of the ſubject; and afterwards to exhibit the particulars in proof.

Mesjid Ali (or Ali's Moſque) is ſituated at about two thirds of the diſtance, and as nearly as poſſible in the line of direction, between Aleppo and Buſſorah; and is a ſort of land-mark to the caravans which paſs the common boundary of the Arabian and Chaldean deſerts. Its geographical poſition is deduced from bearings and latitudes taken by M. NIEBUHR and others: and therefore, as far as general geography is concerned, it may almoſt be regarded as a fixed point. Not that the truth of its poſition will in any ſhape affect the preſent head of enquiry; which is entirely directed towards a compariſon of the ſpaces of time, employed between certain points of the route, by each traveller reſpectively.

Now, that portion of the Deſert route between Mesjid Ali and Buſſorah, being ſubject to great variation in the track, as appears by the journals of different travellers; whilſt the much larger portion of it, between Mesjid Ali and Aleppo, is very nearly the ſame at all times; it is very clear, that this latter portion furniſhes the propereſt ground on which to form the compariſon: and the particulars are as follow:

Aleppo

	CARMICHAEL.	CAPPER.	HUNTER.	
	H. M.	H. M.	H. M.	
Aleppo to Hagla	11 5	11 24	10 0	Hagla.
Hagla to Ain il Koom	37 30	41 4	35 0	to Taiba.
Ain il Koom to Uklet Hauran	80 10	78 41	81 30	to Uklet Hauran.
Hagla to Uklet Hauran	117 40	119 45	116 30	
Uklet Hauran to Al Kadder	53 50	54 45	51 30	
Hagla to Al Kadder	171 30	174 30	168 0	
Al Kadder to Rackama, oppoſite Meſjid Ali	21 45	19 50	19 30	
Hagla to Rackama	193 15	194 20	187 30	
Aleppo to Rackama	204 20	205 44	197 30	

On the Little Deſert I have two examples of time, from Mr. IRWIN in 1781, and Mr. HOLFORD in 1780; both of whom kept regular journals.

	IRWIN.	HOLFORD.
	H. M.	H. M.
Aleppo to Ain il Koom.	52 0	46 27
Ain il Koom to Annah on the Euphrates	76 0	80 15
Aleppo to Annah	128 0	126 42

It appears by the journals, that Mr. IRWIN deviated from the uſual track in the firſt part of his route; and that Mr. HOLFORD did the like in the latter part of his; each to avoid an enemy: ſo that it may be preſumed, that the deviations nearly balanced each other (ſee the ſketch).

Between Annah and Bagdad, theſe gentlemen made part of their journey in the caravan of loaded camels, and partly with *light* camels (that is, without any other load than the rider). Mr. IRWIN employed 62½ hours: but the laſt 15 hours, on

light camels, were at an accelerated rate of half a mile *per* hour, or one fifth part, above the ordinary rate; according to his idea, which I have no doubt was a very juſt one: and this accelerated rate ſhould add 3 hours to the 15, to reduce it to caravan time; making 65½ hours inſtead of 62½. Mr. HOLFORD's journey, by the ſame *ratio*, muſt be reckoned at 68: but as this part of the two journies is obviouſly too inaccurate to draw any concluſions from, in the way of compariſon, I ſhall only make uſe of Mr. IRWIN's time (to which no ſolid objection can be urged) when I calculate the rate of the camel's travelling.

We have now ſeen, that on a journey of about 200 hours, between Aleppo and Mesjid Ali, two accounts differ only 1 hour 24 minutes; and a third differs from the mean of the other two ſeven hours and an half. And we may obſerve, that if the ſtage from Aleppo to Hagla be taken out of the queſtion, the number of Mr. HUNTER's hours would be nearer on an equality with the others by about an hour and a quarter (ſee p. 134.). The reaſon of the different reports of the diſtance between Aleppo and Hagla, appears to be, that travellers commonly join the caravans either at Hagla or on the road to it; and they, travelling by a quicker conveyance than camels afford, and then adjuſting the time to the caravan rate, make different eſtimates of the diſtance. Or there may be ſome other cauſe which has not been explained. Four different perſons give the time as follows:

	H.	M.		H.	M.
CARMICHAEL	11	5	CAPPER	11	24
HUNTER	10	0	HOLFORD	9	12

So that the proper point of outſet in making the compariſon, is

is Hagla. And, reckoning from thence, we have in the first table (p. 134.) the numbers 193¼, 194¼, and 187¼, for the time between Hagla and Mesjid Ali, in the three journies respectively: and the same table affords also the following comparisons between different places on the route:

In one instance,	80⅛, and 78⅔;
In a second,	117¾, 119¼, and 116½;
In a third,	53¼, 54¼, and 51½;
And in a fourth,	171½, 174½, and 168.

Again, between Aleppo and Annah on the Euphrates, the numbers in the second table stand thus:

128, and 126½.

I think I need not produce any more examples to prove the equal rate of motion of a camel that is in any degree loaded; or rather of a number of camels together, where the rate will be determined by the *slow-going ones*: and whatsoever rate, in actual distance, may be deduced from these examples, must be applied to loaded camels travelling in a body together, and not to light camels, or those chosen for speed, whose rate appears to be at least ⅓th greater. By a light camel is meant one that has only a man, or a very small quantity of baggage, on it; whereas a camel's load is 500 to 600 pounds; and camels so loaded, form what is termed the HEAVY caravan. LIGHT caravan, on the contrary, is applied to camels under a *moderate* load, or perhaps little more than half loaded. And with respect to camels, either *moderately* or *fully loaded*, I can perceive no difference in their hourly rate of motion: the difference alone appears in the length of their day's journey; as we shall perceive hereafter. A camel, it is said, will not permit himself to be over-laden; and this may be the reason why the load does not affect his rate of motion.

It

It appears, that the direct distance between Aleppo and Bussorah, is 621 geographic miles, or 720 British, nearly. And Mr. CARMICHAEL's route, traced by a compass through all its principal bendings, and calculated trigonometrically, gives 688 geographic miles, or of British 797. It follows then, of course, that as the same gentleman was 322 hours on the road, the mean hourly rate of the camel's motion, was 2,475 British miles. Colonel CAPPER's route, though easily traced on the map, is not correct enough in its particulars, to serve as an authority equal to Mr. CARMICHAEL's; and the like may be said of Mr. HUNTER's: but they must both be allowed to corroborate Mr. CARMICHAEL's in a general way; for as nearly as Colonel CAPPER's route can be traced, over the CHALDEAN Desert (and, as we have before observed, the track is nearly the same at all times, in all other parts of the Desert) the hourly rate of his camels was 2,51 *per* hour; and that of Mr. HUNTER's 2,585.

We come now to the LITTLE DESERT route. It has been noticed, that Mr. IRWIN employed 128 hours on his journey from Aleppo to Annah; and 65¼ more (allowing for his accelerated rate 3 hours, see p. 135.) between Annah and Bagdad; altogether 193¼ hours between Aleppo and Bagdad. The direct distance between those places is 393 geographic miles; and by the route traced by Mr. IRWIN, the *road distance* comes out about 414½, or British miles 480 *. And this number, divided by 193¼, gives 2,48 *per* hour for the camel's rate; or

* Not that the distance between those places is so much as 480 miles by the direct road: it is probably less than 470. But Mr. IRWIN's party took a circuitous course to the southward, between Aleppo and Ain jl Koom, to avoid an enemy that lay in the way. He estimated his distance at 493 miles.

within a very ſmall fraction of Mr. CARMICHAEL's rate; his being, as we have juſt ſeen, 2,475.

I think I may venture to reſt the calculation of the loaded camel's mean hourly rate of travelling, on the experiments of Mr. CARMICHAEL and of Mr. IRWIN; both of whom appear to have taken much pains with the detail of their journies*; and then it will ſtand as under:

Mr. CARMICHAEL on 322 hours 2,475 } Mean 2,478 Britiſh miles.
Mr. IRWIN on 193½ . . 2,48 }

We have mentioned above, the reſult of Colonel CAPPER's and of Mr. HUNTER's time, which gave a rate ſo very near to CARMICHAEL's and IRWIN's: and it may not be amiſs to add to theſe, the reſult of Mr. HOLFORD's; as well as the eſtimates of the camel's rate, formed by ſeven different perſons. All theſe I have placed in one point of view, in the following table.

	CARMICHAEL.	IRWIN.	CAPPER.	HUNTER.	HOLFORD.	PLAISTED.	Anonymous.
	Brit. mi.						
Eſtimated rates	2,29	2,55	2,25	2,33	2,24	2,3	2,5
Experiments	2,475	2,48	2,51	2,585	2,5	——	——

Mean of the ſeven eſtimates, 2,35.
Mean of the five experiments, 2,51.
Mean of CARMICHAEL's and IRWIN's, 2,478.

* Mr. IRWIN alſo took the bearings of his courſe by a compaſs, though not ſo much in detail as Mr. CARMICHAEL; but Mr. IRWIN not only remarked the time, but the particular rate of travelling, on each day; which appeared to vary from 2 to 3 *per* hour, but was commonly 2½; and the mean of all 2,55 Britiſh miles.

Before

Before I quit the ſubject of the hourly rate, I ſhall obſerve, that the road diſtances in Mr. CARMICHAEL's traverſe table are often thrown together in lines of very conſiderable length, ſuch as 20 miles and upwards; and very commonly in lines of 5, 6, and 7, and yet are all conſidered as ſtraight lines. By this mode, it is probable, that many ſmall inflexions paſſed unnoticed: and therefore the rate taken from the reſults of CARMICHAEL's and IRWIN's obſervations, will be rather under the mark than otherwiſe; but it is obvious, that no kind of rule can be found to correct it by. It is certain, that fewer inflexions are likely to occur in the Deſert routes, than in almoſt any other; not to mention that the long lines of diſtance happen chiefly in the open, level part of the route. The road diſtance exceeds the direct diſtance, by *one-fourteenth part* of the latter only, between Aleppo and Mesjid Ali; amounting to a fraction of ,168 Britiſh mile on each hour; or in the proportion of 7¼ miles on each hundred of direct diſtance. This, I confeſs, is much leſs than I could poſſibly have ſuppoſed; and which nothing ſhort of actual experiment could have induced me to believe. On the whole road between Aleppo and Buſſorah, the proportion is nearly 11 on each hundred, on CARMICHAEL's route: but his route over the Chaldean Deſert was unuſually circuitous; and cannot be admitted to have any weight, in the determination of this queſtion.

One would expect that the inflexions of which no account is taken by Mr. CARMICHAEL, would amount to at leaſt half as much as thoſe which are taken notice of; and ſuch addition would make the whole hourly rate 2,56 inſtead of 2,475. But this is merely a ſuppoſition; and it is poſſible, that the rate may not be higher than 2,52, the mean of the four other experi-

ments. At all events, the error can be but ſmall: and poſſibly, all circumſtances taken into the caſe (and particularly this remarkable one, that of three perſons who attempted to aſcertain the rate, by counting and meaſuring the camel's footſteps, none reckoned it higher than 2¼, and one went ſo low as 2¼), I think the rate of two miles and an half *per* hour may be uſed, as differing but a ſhade from the general reſult; and as having the moſt manageable fraction.

Thus it appears to me, that the hourly rate of the camel may be applied as a very uſeful ſcale to the African geography; whenſoever the uſe of watches ſhall be adopted by the native travellers employed by the AFRICAN ASSOCIATION *; and with ſtill greater advantage, of courſe, if Europeans are employed. And if Mr. CARMICHAEL could deſcribe the general bearing, on a line of more than 700 Britiſh miles, ſo nearly as within 6 or 7 degrees of the truth; and that with a pocket compaſs; nothing more need be ſaid concerning the advantages that may be derived from the uſe of that valuable inſtrument, aided by ſuch a ſcale as I have been deſcribing.

The mean length of the day's journey of the camel, varies according to the degree in which it is loaded: and in this particular it is that the ſtate of the camel, as to its burthen, operates on its progreſs. It is neceſſary to obſerve, that whatſoever remarks I may offer, on the ſubject of the camel's day's journey, are meant to be applied only to the *mean* rate on journies of conſiderable length; ſince any other kind of experiment would be of no uſe in geography: I ſhall therefore conſider only the progreſs of the LIGHT and HEAVY CARAVANS, in which

* In the Memoirs of ABDUL KURRIM, a Caſhmerian of diſtinction, he informs us, that he kept an account of the time, on the road between *Bagdad* and *Mecca*, by means of an European watch, in the year 1740.

the

the camels are left to purſue their journey quietly and at leiſure; and with the regularity of a machine: and not that of the LIGHT CAMELS, which are not only freed from incumbrance, but are alſo urged on.

I have two examples of the heavy kind, and three of the light kind, where the time has been regularly kept: beſides a third example of the heavy kind, where the neceſſary regularity is wanting, but yet containing within itſelf, evidence ſufficiently ſtrong to corroborate the other two.

The HEAVY caravans were thoſe of Mr. CARMICHAEL and M. HOLFORD; the firſt of 1000 camels, of which 600 were loaded, went, on a journey of 45 days, at a mean, each day, ... H. M. 7 10

The ſecond, with 50 loaded camels, on a journey of 15 days 7 40

Mean of the two, 7 25

The third, TEIXEIRA, with 130 loaded camels, on a journey of 21 days, *about* 7 30

Mean of all, *per* day, 7 27

The LIGHT caravans were,

			H. M.
Meſſ. IRWIN, CAPPER, HUNTER,	from 80 to 100 camels,	21 days,	9 12
		33 —	8 38
		34 —	8 45
		Mean of the three	8 52

Here then the mean of the heavy caravan day is under ſeven hours and an half; and that of the light caravan between eight and three quarters, and nine hours.

Some

Some of the ordinary watering places on the Deſert, being from 3 to 5 days journey aſunder, it may be ſuppoſed, that the length of the intermediate journies, would be determined by the known and approved ſtandard of a proper day's walk for a camel: for as they often arrive at the watering-place early in the day, it appears, that the caravan chiefs, do not, by any means, divide the ground between ſuch watering places, into equal portions for marches. This being the caſe, one may expect to arrive at the knowledge of what is at leaſt *intended* for a proper day's journey for a camel, under the circumſtances of the caſe: and indeed the reſult is ſuch as to prove what I have before advanced, reſpecting the length of the mean journies of both kinds. For having ſelected from the five above-mentioned journals, the length of the *apparently optional* journies in each; it appears, that the heavy caravans went 7 h. 51 m. on a mean of 24 ſuch days: and the light caravans 9 h. 8 m. on a mean of 38 days. In both of which caſes, as might have been expected, the length of the mean *optional* day, is ſomewhat longer than that of the whole journey; in which there is a wider field for delays and accidents. The compariſon is as follows:

	Heavy caravan.			Light caravan.		
		Diſtance in Britiſh miles.			Diſtance in Britiſh miles.	
	H. M.	at 2½ *per* hour.	at 2,56	H. M.	at 2½ *per* hour.	at 2,56.
Mean daily rate of the whole journey . .	7 27	18,64	19,06	8 52	22,17	22,7
Of optional days . .	7 51	19,62	20,1	9 8	22,8	23,38

Thus

Thus the mean daily rate of the heavy caravan, appears to be 18,64 Britiſh miles, reckoning two miles and an half for each hour; and 19,06 if taken at 2,56: and the mean rate of the light caravan 22,17 miles, at 2½; 22,7 at 2,56.

In order to apply this ſcale with effect, to the African geography, it is neceſſary to ſtate the number of days that the caravans uſually halt on the road; for as yet I have only conſidered their rate of motion: but it is evident, that if the length of the journey in the groſs, is given, the requiſite information will not be obtained, without a previous knowledge of the time loſt by neceſſary, or unavoidable halts on the road. My enquiries have furniſhed me with an account of 13 halts, to 149 days of travelling; or, which is the ſame thing, 13 halts out of 162 days, reckoned from the time of departure, to the time of the arrival of the caravans at the place of deſtination: that is, 1 halt to 12½ travelling days. This, of courſe, muſt be deducted from the aggregate of the diſtance: or, ſhould it be averaged on each day, the heavy caravan day muſt be reckoned at 17,14 miles inſtead of 18,64; and that of the light caravan 20,4, inſtead of 22,17; when the hourly rate is taken at two miles and an half.

It alſo remains to be ſtated, from the proportion that the road diſtance bore to the direct diſtance, by the trace of Mr. CARMICHAEL's route; what length in direct diſtance, and in geographic miles, may be allowed for each day, for the heavy caravan, on ſimilar lengths of journey, and over ſimilar tracts of country. It appears then, that on the 28 days between Aleppo and Rackama (oppoſite Mesjid Ali) the mean length of the day's journey, in direct diſtance, is about 15¼ geographic miles: and on the whole 45 days between Aleppo and Buſſorah, 13,8 ſuch miles. But this is without any allow-

ance for halts; which, as has been obſerved before, require a deduction of 8 parts in 100, to be made from the groſs amount of the whole journey, when applied to the purpoſes of geography.

I have already taken notice, that Mr. CARMICHAEL counted the camel's ſteps, in order to aſcertain a ſcale of diſtance; and I ſhall now give the reſult of his obſervations, as well as of Mr. HOLFORD's, who alſo counted the ſteps, and meaſured the length of a number of them on the ground. Mr. HUNTER's experiment was on too ſmall a ſcale to ground a calculation on. It is certain, that theſe reports of the number of ſteps during certain portions of time, and the meaſurement of a certain number of thoſe ſteps on the ground, furniſh a reſult that does not agree with the experiments on the great ſcale; ſuch as we have juſt related, and which appear to be ſuſceptible of greater accuracy than thoſe made in detail. But it will, nevertheleſs, be proper to give the reſults, and to make ſome remarks on them; if it be only to prevent any perſon in future from founding a calculation on them.

Mr. CARMICHAEL counted the double ſteps, or rather the return of the ſame foot, of a camel on which he rode, for an hour together, on 20 different days; at times when, from the nature of the ground, he thought the greateſt variation took place, in the rate of motion. He found the greateſt number of ſteps to be 2420, the leaſt 2086; and the mean of the whole 20 hours, was 2200. Mr. HOLFORD reckoned the greateſt 2240, leaſt 2060; mean 2150. They both report the double ſtep to be 5 feet and an half. The reſult of the former account is 2,29 Britiſh miles *per* hour; of the latter 2,24; and each allowed his diſtance accordingly, in his journal: though nothing is more certain than that their computed diſtances fall very

ſhort

Scanderoon
Aleppo
Salt Lake
Euphrates River
Beer
Racca
Hagla
Irwin, 1781.
Carmichael's
Desert
Kirkeria ruins
Ain-il-Koom
Taiba ruin
Syrian
Route, 1751.
Holford.
Hems
Palmyra
Jub il Gannam
1780.
Damascus
Arabian
Mesopo
LITTLE
GREAT
Uklet Ha
Geographic Miles.
Carmichael's Route.
Irwin's.
Capper's where it deviates from Carmichael's.
Holford's.
Teixeira's across the Chaldean Desert.

Sketch of the ROUTES across the DESERTS, Between Aleppo and Buſsorah. 1791.



ſhort of the truth. Mr. CARMICHAEL, for inſtance, reckons the diſtance from Aleppo to Buſſorah, by the road, at 720 Britiſh miles, although the direct diſtance itſelf ſcarcely falls ſhort of it. And Mr. HOLFORD's road diſtance alſo falls very ſhort. Even Mr. CARMICHAEL's higheſt number of paces, would exceed the mean rate given by the experiment at large, by a *ſixtieth* part only.

As theſe gentlemen's experiments differ only 50 ſteps, in the mean number, during the hour (one being 2200, the other 2150), that is, a 44th part, the error muſt be looked for elſewhere; and it probably originated in their meaſuring too ſmall a number of ſteps on the ground to found their calculation on.

The reaſon of this great variation in the number of paces, in a given time, is the plenty or ſcarcity of the Deſert ſhrubs, on which the camels feed, as they go on; and thus ſuch expements become almoſt uſeleſs, unleſs the quality of the Deſert was deſcribed in every part. As the hourly rate of Mr. CARMICHAEL, coincides with that of Mr. IRWIN, within a very ſmall fraction, although the one travelled in November and December, the other in March and April; it appears, that the ſeaſons have little or no effect in this particular: and it is therefore highly probable, that the ſhrubs may flouriſh in ſome parts of the Deſert, and be dried up in others, at one and the ſame ſeaſon.

IX. *On Infinite Series.* *By* Edward Waring, *M. D. F. R. S.* Lucasian *Profeſſor of Mathematics in the Univerſity of* Cambridge.

Read March 24, 1791.

1. MERCATOR firſt publiſhed the continuation of the common method of diviſion to an infinite ſeries of terms proceding according to the dimenſions of a variable quantity; NEWTON did the ſame for the common method of extraction of roots. Dr. BARROW before applied the ſame principles in ſome eaſy examples to find the aſymptotes of curves.

2. The methods of diviſion and extraction of roots were long before taught; but the continuation of them *in infinitum* would have been uſeleſs, as the areæ of curves, whoſe ordinates are $ax^{\frac{n}{m}}$ (where x denotes the abſciſs, and a, n, and m invariable quantities) had not been diſcovered many years before the time of MERCATOR's Publication, and conſequently it would have been of little uſe to transform an ordinate or fluxion, whoſe area or fluent is unknown, into another form, of which the area, &c. is equally unknown.

3. Sir ISAAC NEWTON extended the rule for raiſing a binomial (to any affirmative power) to negative powers, the extraction of roots and fractional indexes, by applying the law of the ſeries for affirmative powers to them, and continuing it *in infinitum*. M. DE MOIVRE extracted the root, &c. of a multinomial by a ſeries of a ſimilar nature; but theſe methods will

will only apply in the moſt ſimple caſes, when not more than one root is to be extracted. In every complicate caſe (*viz.* the extraction of roots of quantities which involve the roots of compound quantities) of irrational quantities, recourſe muſt be had to the old methods of multiplication, diviſion, and extraction of roots.

4. If a root of a complicate irrational quantity be required by a ſeries proceeding according to the dimenſions of x; firſt reduce all the irrational quantities contained under the root by multiplication, diviſion, and extraction of roots into ſerieſes proceeding according to the dimenſions of x, ſo that the terms of the leaſt dimenſions be conſtituted firſt, if an aſcending ſeries be required, and ſo on; and the contrary, if a deſcending; then add the ſeveral ſums together, and extract the root of the reſulting ſum by a ſeries which proceeds according to the dimenſions of x, and it will be the root required.

Ex. Let the value of the quantity $\sqrt[3]{(x^{-1}+a+\overline{b+cx}\times\sqrt[m]{(d+ex+\sqrt{(f+gx^2)})+\sqrt{(b+kx^2)}})}$ be required by an aſcending ſeries; firſt, extract the innermoſt root $\sqrt{(f+gx^2)}=f^{\frac{1}{2}}+\frac{gx^2}{2f^{\frac{1}{2}}}-$ &c. which add to the quantity $d+ex$ contained under the ſame root (m), and place the terms of the ſum according to the dimenſions of x, and it becomes $(d+f^{\frac{1}{2}})+ex+\frac{gx^2}{2f^{\frac{1}{2}}}-$ &c. of which extract the (m) root, and there reſults

$$\left.\begin{aligned}(d+f^{\frac{1}{2}})^{\frac{1}{m}}+\tfrac{1}{m}\times\overline{d+f^{\frac{1}{2}}}^{\frac{1-m}{m}}\times ex+\tfrac{1}{m}\times\tfrac{1-m}{2m}\times\overline{d+f^{\frac{1}{2}}}^{\frac{1-2m}{m}}\times e^2\\ +\tfrac{1}{m}\times\overline{d+f^{\frac{1}{2}}}^{\frac{1-m}{m}}\times\frac{g}{2f^{\frac{1}{2}}}\end{aligned}\right\}x^2+$$

&c. multiply this ſeries into $b+cx$, and thence is derived

$b \times \overline{d+f^{\frac{1}{n}}}^{\frac{1}{m}} + \frac{1}{m} b \times \overline{d+f^{\frac{1}{n}}}^{\frac{1-m}{m}} \times e \Big\} x + \&c. = P$; extract the root

$+ \quad c \times \overline{d+f^{\frac{1}{n}}}^{\frac{1}{n}}$

$\sqrt{b+kx^2} = b^{\frac{1}{2}} + \frac{kx^2}{2b^{\frac{1}{2}}} + \&c. = Q$; add the three quantities P, Q, and $x^{-1} + a$ contained under the ſame root $\sqrt[3]{}$ together, and the ſeries reſulting, whoſe terms are conſtituted according to the dimenſions of x, will be $x^{-1} + (a + b \times \overline{d+f^{\frac{1}{n}}}^{\frac{1}{m}} + b^{\frac{1}{2}} = A) +$ $\left(\frac{1}{m} b \times \overline{d+f^{\frac{1}{n}}}^{\frac{1-m}{m}} \times e + c \times \overline{d+f^{\frac{1}{n}}}^{\frac{1}{n}} = B\right) x + \&c.$; of which extract the cube root $\sqrt[3]{}$, and it will be $x^{-\frac{1}{3}} + \frac{1}{3}Ax^{\frac{2}{3}} + \frac{1}{3}Bx^{\frac{5}{3}} + \&c.$ the root required.

5. The principal uſe of reducing quantities into ſeries proceeding according to the dimenſions of the variable quantity is as before mentioned for finding the area of a curve from its ordinate; or, which correſponds, the integral from its naſcent or evaneſcent increment; but the ſerieſes deduced ſhould converge, otherwiſe from them cannot be found the area or integral. In the Meditationes Analyticæ a method was firſt publiſhed of finding when theſe ſeries will converge and when not, *e. g.* the ſeries $a + bx + cx^2 + dx^3 + \&c. = \int (A + Bx + Cx^2 + Dx^3 + \&c.)^n \times \dot{x} = P$ will converge when (x) either affirmative or negative is leſs than the leaſt root (α) of the equation $A + Bx + Cx^2 + Dx^3 + \&c. = o$, if the roots are poſſible. A ſimilar rule is given when ſome of the roots are impoſſible. The ſeries will diverge when x is greater than α, and the caſes are given in which it will converge when $x = \alpha$. The ſeries deſcending according to the dimenſions of x will converge when x is greater than the greateſt root of the equation, &c. Theſe principles are fur-

ther applied in the ſame Book to complicate irrational algebraical functions of x, &c.

Hence moſt commonly the ſeries for the area contained between two ordinates, or integral between two different increments deduced by the common method will diverge; on which account, in the ſame Book, is given a method by interpolation of finding the area or integral contained between any two different values of x by converging ſeries, if the area, &c. is finite.

6. To find whether a given value $(+a)$ is leſs than the leaſt affirmative or negative root (x) of a given algebraical equation $A+Bx+Cx^2+Dx^3+\&c.=0$, if all its roots are poſſible; transform the equation into another, whoſe root z is the reciprocal of the root $x=\frac{1}{z}$ of the given equation, and for z in the reſulting equation write reſpectively $v+a$ and $v-a$; and if from the former ſubſtitution all the terms become negative or affirmative, and from the latter they become alternately negative and affirmative, then will a be leſs than the leaſt root of the given equation. If in the ſame manner, in the given equation for x be ſubſtituted $v+a$ and $v-a$, and the terms reſult as before, then will (a) be greater than the greateſt root affirmative or negative of the given equation.

7. When the integral of an algebraical quantity, whoſe increments are finite, is required; firſt, by the method given in Medit. Analyt. inveſtigate the integral in finite terms, if it can be expreſſed by them; but if not, reduce it into infinite ſerieſes of which the integral of each of the terms can be found, and alſo the ſerieſes for finding the integral contained between the two different given values of the variable quantity may converge.

Serieſes of this kind have been given in the Medit. Analyt. and innumerable of a like kind may be added for finding integrals

integrals by converging ſerieſes either aſcending or deſcending, of which the given increments are either finite or evaneſcent.

7. 2. It may be obſerved, that generally the particular caſe of which the increments are naſcent or evaneſcent may be deduced from the general, in which the increments are finite; and conſequently in many caſes the general will, *mutatis mutandis*, correſpond to the particular; *e. g.* 1. the integral cannot be expreſſed in finite algebraical terms, when any factor in the denominator of the increment has not a ſucceſſive correſpondent one; which is analogous to the caſe of the ſimple diviſor in the denominator of a fluxion publiſhed in the Quadrature of Curves. 2. Nor can it be expreſſed by the above-mentioned terms, when the dimenſions of the variable quantity in the denominator exceed its dimenſions in the numerator by unity, which correſponds to a ſimilar caſe in fluxions firſt given in Medit. Analyt. To theſe may ſeveral others be added.

8. Let the fluent of the fluxion $(A+Bx^{n}+Cx^{2n}+\&c.)^{m}\times x^{\theta-1}\dot{x}=ax^{\theta}+bx^{\theta+n}+cx^{\theta+2n}+\&c.=(A+Bx^{n}+Cx^{2n}+\&c.)^{m+1}\times \overline{\alpha x^{\theta}+\beta x^{\theta+n}+\gamma x^{\theta+2n}+\&c.}=a'x^{\lambda}+b'x^{\lambda-n}+c'x^{\lambda-2n}+\&c.=(A+Bx^{n}+Cx^{2n}+\&c.)^{m+1}\times \overline{\alpha' x^{\lambda'}+\beta' x^{\lambda'-n}+\gamma' x^{\lambda'-2n}+\&c.}$.

1. If the infinite aſcending ſeries $\alpha x^{\theta}+\beta x^{\theta+n}+\gamma x^{\theta+2n}+\&c.$ converges, then will the ſeries $ax^{\theta}+bx^{\theta+n}+cx^{\theta+2n}+\&c.$ converge, and nearly in the ſame ratio; and *vice verſâ*, if the former diverges, the latter will alſo diverge. In the ſame manner if the infinite deſcending ſeries $\alpha' x^{\lambda'}+\beta' x^{\lambda'-n}+\gamma' x^{\lambda'-2n}+\&c.$ converges, then will the ſeries $a'x^{\lambda}+b'x^{\lambda-n}+c'x^{\lambda-2n}+\&c.$ converge; and if the former diverges, then will the latter alſo diverge; and in all the caſes nearly in the ſame ratio, except only when their convergency is the leaſt.

2. If the series $ax^{\theta} + bx^{\theta+n} + cx^{\theta+2n} +$ &c. converges, then will the series $a'x^{\lambda} + b'x^{\lambda-n} + c'x^{\lambda-2n} +$ &c. diverge, unless in cases of the slowest convergency, where $x = \pm v\sqrt[r]{\pm 1}$, and all the roots of the given equation are of the formula $\pm v\sqrt[r]{\pm 1}$; when $x = \pm v\sqrt[r]{\pm 1}$, the successive terms of the infinite series deduced from algebraical quantities by the preceding method will ultimately, that is, at an infinite distance come more near to a ratio of equality with each other than any assignable difference.

3. If the fluxion be $(A + Bx^{n} + Cx^{2n} \ldots\ldots x^{rn})^{m} \times x^{\theta-1}\dot{x}$, then will $\lambda = mrn + \theta$, and $\lambda' = \theta - rn$.

Many more propositions concerning infinite series and their convergency are given in the Medit. Analyt.

4. Let the fluxion be $\overline{a + bx^{n}}^{m} \times x^{n}\dot{x}$, of which find the fluent in a series ascending according to the dimensions of x^{n}, and it will be $a^{m} \times x^{b+1} \left(\frac{1}{b+1} + \frac{m}{b+n+1} \times \frac{b}{a}x^{n} + m \cdot \frac{m-1}{2(b+2n+1)} \times \frac{b^{2}}{a^{2}}x^{2n}\right.$ $\left. + m \cdot \frac{m-1}{2} \cdot \frac{m-2}{3(b+3n+1)} \times \frac{b^{3}}{a^{3}}x^{3n} + \&c.\right) = \frac{b^{m}x^{mn+b+1}}{mn+b+1} +$ $\frac{mb^{m-1}ax^{mn+b+1-n}}{mn+b+1-n} + m \cdot \frac{m-1}{2} \times \frac{b^{m-2}a^{2}x^{mn+b+1-2n}}{mn+b+1-2n} +$ &c. $= \overline{a + bx^{n}}^{m+1} \times \left(\frac{x^{b+1}}{b+1a} - \frac{\overline{m+1}(n)+b+1}{\overline{b+1} \cdot \overline{b+1+n}} \times \frac{b}{a^{2}}x^{b+n+1} + \frac{(\overline{m+1}(n)+b+1).(\overline{m+2}(n)+b+1)}{\overline{b+1} \cdot \overline{b+1+n} \cdot \overline{b+1+2n}} \right.$ $\left. \times \frac{b^{3}}{a^{3}} \times x^{b+2n+1} - \frac{\overline{(m+1} \cdot n+b+1) \cdot \overline{(m+2n+b+1)} \cdot \overline{(m+3n+b+1)}}{\overline{b+1} \cdot \overline{b+1+n} \cdot \overline{b+1+2n} \cdot \overline{b+1+3n}} \times \right.$ $\left. \frac{b^{3}}{a^{3}}x^{b+3n+1} + \&c.\right) = (bx^{n} + a)^{m+1} \times \left(\frac{1}{(mn+b+1)b} \times x^{b+1-n} - \right.$ $\left. \frac{\overline{b+1-n}a}{mn+b+1 \cdot \overline{mn+b+1-n}b^{2}} \times x^{b+1-2n} + \right.$ $\left. \frac{\overline{b+1-n} \times \overline{b+1-2n} \times a^{2}}{mn+b+1 \times mn+b+1-n \times mn+b+1-2n \times b^{3}} \times x^{b+1-3n} - \&c.\right)$ (L). The

first

firſt ſeries, which aſcends according to the dimenſions of x^n, terminates, when $\frac{mn+b+1}{n}$ is a whole negative number; the ſecond, which deſcends according to the dimenſions of the ſame quantity x^n, terminates when $\frac{b+1}{n}$ is a whole affirmative number. The fluent will terminate both ways when $\frac{b+1}{n}$ is a whole affirmative and m a negative number greater than $\frac{b+1}{n}$.

5. When m is a whole number or $=0$, the fluent $\int \overline{a+bx^n}^m x^b \dot{x}$ can always be found in finite terms of x, or in the above-mentioned finite terms, together with the log. of x; which appears from reducing $\overline{a+bx^n}^m$ into ſimple terms $a^m + ma^{m-1}bx^n + m \cdot \frac{m-1}{2} a^{m-2} b^2 x^{2n} +$ &c., and multiplying them into $x^b \dot{x}$, and finding the fluents of the reſulting fluxions: but the ſeries found by the preceding method will not always terminate when m is a whole number, and the ſeries findable as above mentioned; when properly corrected, it may be rendered findable, or, which means the ſame thing, the ſeries may be made to terminate.

6. When the ſeries which expreſſes the fluent of a fluxion terminates, we may begin either from one end of the ſeries, or the other; for example, in finding the fluent of the fluxion $\overline{a+bx^n}^m \times x^b \dot{x}$, either aſſume the ſeries $\overline{a+bx^n}^{m+1} \times (\alpha x^{b+1} + \beta x^{b+n+1} + \gamma x^{b+2n+1} +$ &c.) or the ſeries $\overline{a+bx^n}^{m+1} \times (A + Bx^n + Cx^{2n} +$ &c.); the former, as is before mentioned, terminates when

when $\frac{mn+b+1}{n}$ is a whole negative number; the latter when $\frac{b+1}{n}$ is a whole affirmative number.

In like manner assume for a descending series $(bx^n+a)^{m+1} \times (\alpha' x^{b+1-n} + \beta' x^{b+1-2n} + \gamma' x^{b+1-3n} + \&c.)$, or $(bx^n+a)^{m+1} \times (Ax^{-m+1n} + Bx^{-m+2n} + Cx^{-m+3n} + \&c.)$; the former will terminate when $\frac{b+1}{n}$ is a whole affirmative; the latter when $\frac{mn+b+1}{n}$ is a whole negative number.

It appears, therefore, that a series will terminate equally by an ascending or descending series; and the end of the one ascending series is the beginning of its correspondent descending one. All these serieses, which do not terminate, proceed on *in infinitum*; one term in the former series becomes infinite, when $b+1+\varkappa n=0$; and in the latter (L) when $mn+b+1-\varkappa n=0$, $\varkappa$ being a whole affirmative number.

10. It has been observed in the Medit. Analyt. that if some quantities contained in the given irrational ones are much less or greater than the rest, it may be preferable in the former case to reduce them into serieses not proceeding according to the dimensions of x, but according to the dimensions of those quantities; and in the latter case according to the reciprocal dimensions of them; and particularly so if the fluent or integral of the terms of the resulting serieses can be found in finite terms, or by tables already calculated.

From similar principles to those before given may be found when the resulting series will converge, and when not.

This method will in many problems be useful, when the value of a near approximate is known.

Of this caſe I ſhall ſubjoin a few examples, of which ſome have been already publiſhed in the Medit. Analyt.

Ex. 1. Let the ſine and coſine of a given arc A be reſpectively s and c; then will the ſine and coſine of an arc $A+e$, where e bears a very ſmall ratio to A, be reſpectively $s+\frac{ce}{r}-\frac{s}{1\cdot 2r^2}e^2-\frac{ce^3}{1\cdot 2\cdot 3r^3}+\frac{s}{1\cdot 2\,3\cdot 4r^4}e^4+$ &c. (the ſigns proceed by pairs alternately negative and affirmative) $=s\left(1-\frac{1}{1\cdot 2r^2}\times e^2+\frac{1}{1\cdot 2\cdot 3\cdot 4r^4}\times e^4-\text{&c.}\right)+c\left(\frac{e}{r}-\frac{e^3}{1\cdot 2\cdot 3r^3}+\frac{e^5}{1\cdot 2\cdot 3\cdot 4\cdot 5r^5}-\text{&c.}\right)$; and $c-\frac{s}{r}e-\frac{1}{1\cdot 2r^2}ce^2+\frac{1}{1\cdot 2\cdot 3r^3}\times se^3+$ &c. (the ſigns proceed as before by pairs alternately negative and affirmative) $=c\left(1-\frac{1}{1\cdot 2r^2}e^2+\frac{1}{1\cdot 2\cdot 3\cdot 4r^2}e^4-\text{&c.}\right)-\frac{s}{r}\left(e-\frac{1}{2\cdot 3r^2}e^3+\frac{1}{2\cdot 3\cdot 4\cdot 5r^4}e^5-\text{&c.}\right)$

This ſeries can alſo eaſily be derived from NEWTON's ſeries by plane trigonometry, and will converge much ſwifter than the ſeries $A-\frac{A^3}{1\cdot 2\cdot 3r^2}+\frac{A^5}{2\cdot 3\cdot 4\cdot 5r^4}-$ &c. &c. if e bears a ſmall proportion to A.

Ex. 2. Let e be a ſmall quantity in proportion to t, and the given fluxion $\frac{\dot{t}}{r^2+(t+e)^2}=\frac{\dot{t}}{r^2+t^2}-\frac{2te+e^2}{(r^2+t^2)^2}\dot{t}+\frac{(2te+e^2)^2}{(r^2+t^2)^3}\times\dot{t}-\frac{(2te+e^2)^3}{(r^2+t^2)^4}\times\dot{t}+$ &c. $=\frac{\dot{t}}{r^2+t^2}-\frac{2t\dot{t}}{(r^2+t^2)^2}\times e-\frac{(r^2+t^2-2^2\times t^2)\dot{t}}{(r^2+t^2)^3}\times e^2+2t\times\frac{(2r^2+t^2-2^2t^2)\dot{t}}{(r^2+t^2)^4}\times e^3+$ &c. (P); the co-efficient of a term $\dot{t}\times e^m$ of this ſeries will be a fraction whoſe numerator is $\overline{r^2+t^2}^m-\overline{m+1}\cdot\frac{m}{2}\times 2^2\times t^2\times\overline{r^2+t^2}^{m-1}+\overline{m+2}\cdot\frac{m+1}{2}\cdot\frac{m}{3}\cdot\frac{m-1}{4}\times 2^4\times t^4\times\overline{r^2+t^2}^{m-2}-\overline{m+3}\cdot\frac{m+2}{2}\cdot\frac{m+1}{3}\cdot\frac{m}{4}\cdot\frac{m-1}{5}\cdot\frac{m-2}{6}\times$

$\times 2^6 \times t^6 \times \overline{r^2+t^2}^{m-3}$ + &c., and denominator $(r^2+t^2)^{2m+1}$; and the co-efficient of a term $\dot{t} \times e^{2m+1}$ will be a fraction whose numerator is $2t \times \left(\overline{m+1} \cdot \overline{r^2+t^2}^{m} - \overline{m+2} \cdot \frac{m+1}{2} \cdot \frac{m}{3} \times 2^2 \times t^2 \times \right.$ $\left. \overline{r^2+t^2}^{m-1} + \overline{m+3} \cdot \frac{m+2}{2} \cdot \frac{m+1}{3} \cdot \frac{m}{4} \cdot \frac{m-1}{5} \times 2^4 t^4 \times \overline{r^2+t^2}^{m-2} - \right.$ &c.) and denominator $(r^2+t^2)^{2m+2}$; m being a whole number. The continuation of these series is too evident to need enunciation, as must generally be the case when the number of factors contained in the successive terms continues the same or increases in arithmetical progression; and the factors themselves increase or diminish in an arithmetical progression: the terms proceed alternately + and − by pairs.

2. 2. Let the given fluxion be $\frac{\dot{e}}{r^2+(t+e)^2} = \frac{\dot{e}}{r^2+t^2} - \frac{\overline{2te+e^2} \times \dot{e}}{(r^2+t^2)^2} +$ $\frac{(2te+e^2)^2 \times \dot{e}}{(r^2+t^2)^3}$ − &c. $= \frac{\dot{e}}{r^2+t^2} - \frac{2t\dot{e}}{(r^2+t^2)^2} \times e - \frac{(r^2+t^2-2^2t^2)\dot{e}}{(r^2+t^2)^3} \times e^2$ + &c. (Q); this series becomes the same as the preceding by substituting in it $\dot{e}$ for $\dot{t}$; in this series e is considered as variable, in the preceding t.

The fluent of the former series (P) is $\int \frac{\dot{t}}{r^2+t^2} + \frac{e}{r^2+t^2} -$ $\frac{t}{(r^2+t^2)^2} \times e^2 - \frac{(r^2+t^2-2^2t^2)}{3(r^2+t^2)^3} \times e^3$ − &c.; the fluent of the latter series (Q) is the same as of the former (P) except the first term $\int \frac{\dot{t}}{r^2+t^2}$: the co-efficient of the term e^b in the fluents will be the same as the co-efficient of the term e^{b-1} in the series (P) and (Q) divided by b.

This series, when the tangent of a given arc is known, finds the arc whose tangent differs very little from the given tangent.

11. Let t be increased or diminished by a quantity e in (P), where e bears a very small ratio to any root or value of t in

the equations $P=0$ or $\frac{1}{P}=0$; and the resulting quantity expanded into a series $a+be+ce^2+de^3+$&c. proceeding according to the dimensions of e, and this series be multiplied respectively into $\dot{t}$ and $\dot{e}$; and the fluents of the resulting fluxions found; then will the former differ from the latter by $\int a\dot{t}$; for the former will be $\int a\dot{t}+ae+\frac{\dot{a}}{2\dot{t}}\times e^2+\frac{\ddot{a}}{2\cdot 3\dot{t}^2}\times e^3+\frac{\dddot{a}e^4}{2\cdot 3\cdot 4\dot{t}^3}+$&c., and the latter $ae+\frac{\dot{a}}{2\dot{t}}e^2+\frac{\ddot{a}e^3}{2\cdot 3\dot{t}^2}+$&c.

2. In the same manner, if more than one variable quantities x, y, z, &c. are contained in (P), which are increased by small increments or decrements α, β, γ, &c., may the increments or decrements of the quantity (P) be deduced from the incremential theorem.

Ex. Let the quantity (P) be $(a+bx^n)^m$, and a, b, x, n, and m be increased by very small increments α, β, γ, δ, and ε; then will $\overline{a+\alpha+\overline{b+\beta}\times\overline{x+\gamma}^{n+1}}^{m+1}=(a+bx^n)^m+(a+bx^n)^m\times$ log. $\overline{a+bx^n}\times\varepsilon+m\times(a+bx^n)^{m-1}\times(\alpha+\beta x^n+b(x^n\delta\times\text{log. }x+nx^{n-1}\gamma))+$&c.

The terms are to be so placed, according to the dimensions of the increments or decrements, that the greatest may first occur.

12. Let some compound quantities be increased by any small quantities variable or invariable, but not the variable quantities contained in them; then reduce the given quantity into a series proceeding according to the dimensions of the small quantities, and find the fluent, integral, &c. as required.

Ex. Let the given quantity be $\frac{x^m\dot{x}}{(x^n+a^n+\beta)^r}=\frac{x^m\dot{x}}{(x^n+a^n)^r}-$

$r\beta$

$\frac{r\beta\dot{x}x^{m}}{(x^{n}+a^{n})^{r+1}}+r\cdot\frac{r+1}{2}\times\frac{\beta^{2}\dot{x}x^{m}}{(x^{n}+a^{n})^{r+2}}-r\cdot\frac{r+1}{2}\cdot\frac{r+2}{3}\times\frac{\beta^{3}\dot{x}x^{m}}{(x^{n}+a^{n})^{r+3}}+$ &c., where the quantity $x^{n}+a^{n}$ is increaſed by a ſmall quantity β; then will $\int\frac{x^{m}\dot{x}}{(x^{n}+a^{n}+\beta)^{r}}=\int\frac{x^{m}\dot{x}}{(x^{n}+a^{n})^{r}}-\frac{\beta}{n\times a^{n}}\left(\frac{x^{m+1}}{(x^{n}+a^{n})^{r}}-\overline{m+1}\right.$

$\left.\overline{-rn}\times P\right)+\frac{r\beta^{2}}{2\cdot n\times a^{n}}\left(\frac{x^{m+1}}{(x^{n}+a^{n})^{r+1}}-(m+1-\overline{r+1}n)\times Q\right)$

$\frac{r\cdot\overline{r+1}\beta^{3}}{2\cdot3\cdot n\times a^{n}}\left(\frac{x^{m+1}}{(x^{n}+a^{n})^{r+2}}-(m+1-\overline{r+2n})\times R\right)+\frac{r\cdot\overline{r+1}\cdot\overline{r+2}\cdot\beta^{4}}{2\cdot3\cdot4\cdot n\times a^{n}}$

$\left(\frac{x^{m+1}}{(x^{n}+a^{n})^{r+3}}-(m+1-\overline{r+3}n)\times S\right)$ &c. The continuation of the ſeries is evident; the letters P, Q, R, S, &c. denote the fluents $\int\frac{x^{m}\dot{x}}{(x^{n}+a^{n})^{r}}$, $\int\frac{x^{m}\dot{x}}{(x^{n}+a^{n})^{r+1}}$, $\int\frac{x^{m}\dot{x}}{(x^{n}+a^{n})^{r+2}}$, &c.

From the length of the arc of an hyperbola or ellipſe given may be deduced by this ſeries the length of a correſpondent arc of an hyperbola or ellipſe, of which the equations expreſſing the relation between its abſciſſæ and ordinates differ only by very ſmall quantities from the equations expreſſing the relation between the abſciſſæ and ordinates of the former.

13. The ſame principles may be applied to the reſolution of algebraical, fluxional, incremential, &c. equations.

Ex. Let L*m*M, &c. (Tab. IV. fig. 1.) be a given circle, whoſe center is C and radius (r), and it be required to find an arc LM, ſo that the area LSM deſcribed round a given point S contained between the lines LS, SM, and the arc of the circle LM be equal to a given area (a).

Find an arc Lm = A nearly equal to the arc required LM, of which to radius 1 ſubſtitute s for the ſine, and c for the coſine, and write e for mM = LM − Lm and b for SC; then will

$\mathrm{LM}\times\frac{r}{2}\left(\overline{\mathrm{A}+e}\times\frac{r}{2}\right)=\mathrm{SC}\times\frac{r}{2}\times\text{fin. }s\text{ arc}:\mathrm{LM}\left(\frac{b}{2}\times(\mathrm{A}+e-\right.$

$\mathrm{A}+e$

$\frac{\overline{A+e^3}}{2\cdot 3r^2}+\frac{\overline{A+e^5}}{2\cdot 3\cdot 4\cdot 5r^4}$ &c.)) $=\frac{A}{2}\times r\pm\frac{b}{2}\times rs+\frac{r}{2}\pm\frac{b}{2}\times c\times e\mp$ $\frac{b}{2r}\times\left(\frac{1}{2}\times se^2+\frac{1}{2\cdot 3r}\times ce^3-\frac{1}{2\cdot 3\cdot 4r^2}\times se^4-\frac{1}{2\cdot 3\cdot 4\cdot 5r^3}\times ce^5+\right.$ &c.) $=a$; in this equation for $\frac{2a-Ar\mp b\times rs}{r\pm bc}$ substitute π, and for $r\pm bc$ write t, and the equation resulting will be $e\mp$ $\frac{b}{2tr}se^2\mp\frac{b}{1\cdot 2\cdot 3t^2}\times ce^3\pm\frac{b}{1\cdot 2\cdot 3\cdot 4t^3}\times se^4\pm\frac{b}{1\cdot 2\cdot 3\cdot 4\cdot 5t^4}\times ce^5$ $\mp$ &c. $=\pi$. From it find e in terms proceeding according to the dimensions of π, and there results $e=\pi\pm$ $\frac{bs}{1\cdot 2tr}\times\pi^2\pm\frac{3b^2s^2\pm bct}{2\cdot 3t^3r^2}\times\pi^3\pm$&c.

This method of resolving KEPLER's and other problems of cutting a given area described round a point, whether focus or not, in a circle, when an approximate sufficiently near to the area to be found is given, will converge as swift as any known method.

The resolution of this problem may be deduced somewhat different by the following methods. Let the letters b, r, a, denote the same quantities as before, and s be the sine of the arc Lm to radius 1, and $s+o$ the sine of the arc LM nearly $=$ Lm; then will $\frac{r^2}{2}\int\frac{\dot{s}}{\sqrt{(1-s+o^2)}}\pm\frac{br}{2}\times\overline{s+o}=a$; reduce the fluxion $\frac{\dot{s}}{\sqrt{(1-s+o^2)}}$ into a series $P\dot{s}+Qo\dot{s}+Ro^2\dot{s}+So^3\dot{s}$ + &c. and find the fluents of the fluxions $Q\dot{s}$, $R\dot{s}$, $S\dot{s}$, &c., which let be B, C, D, &c., there will result the equation $o+\frac{C}{t}o^2+\frac{D}{t}o^3+$&c. $=$ $\frac{2a\mp brs-r\times A}{t}=\pi$, where $t=Br^2\pm br$ and $A=$arc Lm; find o in terms of a series proceeding to the dimensions of π, and consequently $s+o$ the sine of the arc LM required.

From

From ſimilar principles may be found the tangent of the arc required from the equation $\frac{r^2}{2}\int\frac{i}{1+(t+i)^2} \mp \frac{\overline{t+i}\times r\times b}{2\sqrt{(1+\overline{t+i}^2)}} = \alpha$, where the tangent t to rad. 1 of the arc $Lm = A$ is given. Let the fluent of the fluxion $\left(\frac{i}{1+(t+i)^2}\right) = A + Bi + Ci^2 + Di^3 + \&c.$; and $\frac{\overline{t+i}\times b}{\sqrt{(1+\overline{t+i}^2)}} = A' + B'i + C'i^2 + D'i^3 + \&c.$, the equation will become $i + \frac{rC \pm C'}{rB \pm B'} i^2 + \frac{rD \pm D'}{rB \pm B'} \times i^3 + \&c. = \frac{2\alpha - r^2 A \mp rA'}{(r^2 B \pm rB')} = \pi$; from this equation inveſtigate $i = \pi + P\pi^2 + Q\pi^3 + \&c.$, and thence $t + i$ the tangent required.

In like manner may be found the ſecant, coſine, &c. of the arc required.

14. The ſame principles may be applied to cut an area deſcribed round any given point in a given curve equal to an area α.

Let x be the abſciſs and y the ordinate of the given curve, and b the diſtance of the beginning of the abſciſs from the given points, and let (A) be the area of the curve deſcribed round the point S, when the abſciſs is x, which differs very little from the given area (α); to find the value $x + e$ of the abſciſs, when the area $= \alpha$.

Let $y = X$ a function of x, and in X for x write $x + e$, and reduce the reſulting quantity (X′) into a ſeries $X + Be + Ce^2 + De^3 + \&c.$ proceeding according to the dimenſions of e; then will the area $\int y\dot{x} = \int X\dot{x} + e\int B\dot{x} + e^2\int C\dot{x} + e^3\int D\dot{x} + \&c. = A + be + ke^2 + le^3 + \&c.$, and conſequently $A + be + ke^2 + le^3 + \&c. \pm (b \pm \overline{x+e}) \times \frac{y}{2} = A + be + ke^2 + le^3 + \&c. + \frac{1}{2}(b \pm \overline{x+e}) \times (X + Be + Ce^2 + De^3 + \&c.) = A + \frac{1}{2}(b \pm x) \times X + (b + \frac{1}{2}b \pm x \times$

$B+\frac{1}{2}X)e+(k+\frac{1}{2}(b\pm x)C+\frac{1}{2}B)e^2+(l+\frac{1}{2}\overline{b\pm x}\times D+\frac{1}{2}C)e^3+$ &c. $=\alpha$; find the value of e in a ſeries proceeding according to the dimenſions of $\frac{\alpha-A-\frac{1}{2}(b\pm x)\times X}{b+\frac{1}{2}\overline{b\pm x}\times B+\frac{1}{2}X}$, and $x+e$ will be the abſciſs required; X denotes the value of y, when the abſciſs $=x$.

From the ſame principles may the ordinate y, &c. be found.

This problem may be reſolved in the ſame manner, when X denotes an infinite ſeries deduced from an equation expreſſing the relation between the abſciſs and ordinate of the given curve.

If the given area α be the difference between two areas SPM (α') and SPQ $=\beta$ (fig. 2.); for α ſubſtitute $\alpha'-\beta$, and the operation will be the ſame as the preceding.

15. 1. Given any equations, of which the increments of the quantities contained in them can be found from each other, and given approximate values of each of the unknown quantities, which nearly correſpond to each other; to find approximations, which differ leſs from the quantities themſelves than the given ones.

Suppoſe each of the given approximate values to be increaſed or diminiſhed by ſmall increments or decrements, as e, o, i, &c. which are the approximations to be found; and from the given find the equations reſulting from this hypotheſis; and from theſe may be deduced, by ſimple equations, the approximations ſought e, i, o, &c. by neglecting in them all the powers of e, o, i, &c. except the ſimple ones, and all the products of them multiplied into each other; and conſequently the equations deduced will contain only given quantities and ſimple powers of the unknown e, i, o, &c. to be found.

2. When two or more (n) values of one (x) of the unknown quantities are nearly equal to its given approximate; then the

equation which finds the approximate to x will be a quadratic or an equation of n dimenſions.

3. The approximations found by this method will converge more or leſs, according as the approximations given are more or leſs near to one value of each of the unknown quantities than to the remaining ones, &c.

Theſe principles were printed in the Medit. Algebr. in the years 1768 and 1769.

LEMMA.

15. Let *kab* (fig. 3.) be a circle, whoſe center is o; and Po perpendicular to the plane of the circle; and the force of any corpuſcle in the circle on the particle P vary as the corpuſcle divided by the (n) power of its diſtance from the particle; to find the attraction of the circle *kab* on the particle P.

From the ſuppoſita the force of any ring contained between the neareſt concentric circles *lcd* and *hef* of which the center is o, on the particle P, will be as the area of the ring divided by the nth power of the diſtance Pb; and conſequently if Po = A, $ob = v$, $bl = \dot{v}$, and p = periphery of a circle of which the radius is 1; the attraction of the ring on P will be as $\frac{p \times v\dot{v}}{(A^2+v^2)^{\frac{n}{2}}}$, and the force of it in the direction Po as $\frac{Po}{Pb} \times \frac{p v \dot{v}}{(A^2+v^2)^{\frac{n}{2}}} = \frac{pAv\dot{v}}{(A^2+v^2)^{\frac{n+1}{2}}}$; of which fluxion the fluent is $\frac{pA}{1-n \times (A^2+v^2)^{\frac{n-1}{2}}}$; and thence the force of the circle *kab* on the particle P will be as $\frac{p \times A}{1-nPb^{n-1}} - \frac{p \times A}{1-nPO^{n-1}} \left(\frac{p}{1-nA^{n-1}} \right)$.

Cor. 1. The force of the area contained between the two circles *kab* and *lcd* on the particle P will be as $\frac{pA}{1-nPk^{n-1}} - \frac{pA}{1-nPl^{n-1}}$.

15. Let a ſolid be generated by the rotation of a given curve round a line in the axis P*o* or P*o* produced, and A the diſtance of P from *o*, and x=diſtance of any circle generated by the rotation of a point in the curve round the axis P*o* from *o*; and y the ordinate to the given curve, of which the abſciſs is x; and the ſame things be ſuppoſed as before; then will the attraction of the ſolid on the point P vary as $\int \frac{p \times \overline{A-x} \times \dot{x}}{\overline{n-1}(A-x)^{n-1}} \left(\frac{p\dot{x}}{\overline{n-1}(A-x)^{n-2}} \right) - \int \frac{p \times \overline{A-x} \times \dot{x}}{\overline{n-1} \times (\overline{A-x}^2 + y^2)^{\frac{n-1}{2}}}$.

Let the given curve be a circle, of which the center is *o*, and radius t, and conſequently $y^2 = t^2 - x^2$, and $\int \frac{p\dot{x}}{\overline{n-1}(A-x)^{n-2}}$

$$- \int \frac{p \times \overline{A-x} \times \dot{x}}{\overline{n-1} \cdot (\overline{A-x}^2 + y^2)^{\frac{n-1}{2}}} = \frac{p}{\overline{n-1} \cdot \overline{n-3} \times (A-x)^{n-3}} -$$

$$\int \frac{p\dot{x} \times \overline{A-x}}{\overline{n-1} \times (A^2 + t^2 - 2Ax)^{\frac{n-1}{2}}} = \frac{p}{\overline{n-1} \cdot \overline{n-3}(A-x)^{n-3}} - \frac{p}{n-1} \left(\frac{1}{n-3} + \right.$$

$$\left. \frac{A^2+t^2}{\overline{n-5} \cdot \overline{n-3}A^2} - \frac{x}{n-5A} = \frac{\overline{n-4}A^2 + t^2 - \overline{n-3}Ax}{\overline{n-5} \cdot \overline{n-3} \times A^2} \right) \times (A^2 + t^2 - 2Ax)^{\frac{3-n}{2}}.$$

By ſubſtituting $-t$ and t for x in the preceding expreſſion the attraction of the whole globe on the particle P will be found $\frac{p}{n-1} \left(\frac{A^2 + \overline{n-3}At + t^2}{\overline{n-5} \cdot \overline{n-3} \times A^2} \times \overline{A+t}^{3-n} \text{ (L)} - \frac{A^2 - \overline{n-3}At + t^2}{\overline{n-5} \cdot \overline{n-3} \times A^2} \times \right.$ $\left. \overline{A-t}^{3-n} \text{ (H)} \right)$.

16. If the particle be ſituated within the globe, and conſequently A leſs than t or $-t$, and n be an odd number negative

or

or affirmative; or fraction, of which the numerator is odd; then the preceding reſolution will be phyſically juſt; for in this caſe, if the attraction on one ſide of the particle be affirmative, the attraction on the other will mathematically be negative, that is, phyſically oppoſite: but if n be an even number, or fraction, of which the numerator is an even and denominator an odd number, the mathematical ſolution will not agree to the phyſical; for in the former the force on both ſides will be affirmative, in the latter the forces will be oppoſite, and therefore phyſically the force in this caſe will vary as $L+H$, and not as $L-H$, which is the force in both the caſes when A is greater than t and $-t$. The ſame may be applied to the more general reſolution.

17. Let ABCD (fig. 4.) be a globe, of which the diameters AB and CD are ſituated at right angles to each other, and AHBL be a ſpheroid generated by the revolution of an ellipſe on its axis AB, to which let HL nearly equal to CD = AB be the conjugate, and P a point in the axis BA produced; to find the attraction of the ring contained between the globe and the ſpheroid on the point P, on the ſuppoſition that the force of any corpuſcle in the ring on the particle P varies as the magnitude of the corpuſcle directly, and the nth power of its diſtance from the corpuſcle inverſely.

Let $AB=CD=2t$, $CD=HL\ (2c)+2CH=2c+2e$, and conſequently $OH=c=t-e$, where e has a very ſmall ratio to t, $oP=A$, $po=x$, and pM parallel to $CD=y$; then, by the preceding lemma, the attraction of the circle whoſe radius is pM on the point P will be $\frac{p\times\overline{A-x}}{n-1\,Pp^{n-1}}-\frac{p\times\overline{A-x}}{n-1\,PM^{n-1}}$; and in like manner the attraction of the circle, whoſe radius is pm,

will be $\frac{p \times \overline{A-x}}{n-1 \cdot Pp^{n-1}} - \frac{p \times \overline{A-x}}{n-1 \cdot Pm^{n-1}}$, and consequently the attraction of the ring contained between the two circles is $\frac{p \times \overline{A-x}}{n-1 \cdot Pm^{n-1}}$

$$- \frac{p \times \overline{A-x}}{n-1 \cdot PM^{n-1}} = \frac{p}{n-1} \times \left(\frac{A-x}{\left(\overline{A-x}^2 + \frac{\overline{t-x}^2}{t^2} \times t^2 - x^2\right)^{\frac{n-1}{2}}} - \right.$$

$$\left. \frac{A-x}{\left(\overline{A-x}^2 + t^2 - x^2\right)^{\frac{n-1}{2}}} \right) = \frac{p}{n-1} \times \frac{\overline{n-1} \times \overline{A-x} \times \left(t - \frac{x^2}{t}\right)}{\left(A^2 + t^2 - 2Ax\right)^{\frac{n+1}{2}}} \times e + \&c. \times e^2 +$$

$\&c. = p \times \left(t - \frac{x^2}{t}\right) \times \frac{A-x}{\left(A^2 + t^2 - 2Ax\right)^{\frac{n+1}{2}}} \times e$ nearly, which multiplied into $\dot{x}$, and the fluent of the resulting fluxion found, it will be $p \times \left(A^2 + t^2 - 2Ax\right)^{\frac{1-n}{2}} \times \left(\left(\frac{1}{n-7 \times tA} = d \right) x^3 - \left(\frac{1}{n-5t} + \right.\right.$

$\left.\left. \frac{3d \times \overline{A^2 + t^2}}{n-5A} = c \right) x^2 - \left(\frac{t}{n-3A} + 2c \times \frac{A^2 + t^2}{n-3 \times A} = b \right) x + \frac{t}{n-1} - \frac{b \times \overline{A^2 + t^2}}{n-1A} \right)$ e(M).

If the attraction of the whole ring contained between the sphere and spheroid be required, substitute in the fluent (M) for $\dot{x}$, t and $-t$, and proceed as in the preceding cases; in like manner may be found the attraction of the ring contained between any two values of x.

18. The same principles may be applied to find the attraction of the above-mentioned ring, when the line Pp is not perpendicular to the circles pM, pm, CD, &c., and does not cut the diameters CD, M′M, &c. into two equal parts. They may be further applied for finding the attraction of rings contained between any other given solids, of which the equations differ by very small quantities from each other; for example,

between two ſpheroids, of which the axes in the one do not differ much from correſpondent axes in the other, in which caſe the fluents found of the following fluxions may be uſeful.

$$1.\ \int \frac{\dot{x}}{(t^2-x^2)^{\frac{n-1}{2}}} = \frac{1}{n-3t^2} \times \frac{x}{(t^2-x^2)^{\frac{n-3}{2}}} + \frac{n-4}{n-3 \cdot n-5t^4} \times \frac{x}{(t^2-x^2)^{\frac{n-5}{2}}}$$

$$+ \frac{\overline{n-4} \cdot \overline{n-6}}{\overline{n-3} \cdot \overline{n-5} \cdot \overline{n-7}t^6} \times \frac{x}{(t^2-x^2)^{\frac{n-7}{2}}} + \frac{\overline{n-4} \cdot \overline{n-6} \cdot \overline{n-8}}{\overline{n-3} \cdot \overline{n-5} \cdot \overline{n-7} \cdot \overline{n-9}t^8} \times$$

$$\frac{x}{(t^2-x^2)^{\frac{n-9}{2}}} \cdot \cdot \cdot \cdot \cdot \frac{\overline{n-4} \cdot \overline{n-6} \cdot \overline{n-8} \cdot \cdot \cdot 3}{\overline{n-3} \cdot \overline{n-5} \cdot \overline{n-7} \cdot \overline{n-9} \cdot \cdot \cdot 2t^{n-3}} \times \frac{x}{t-x} +$$

$$\frac{\overline{n-4} \cdot \overline{n-6} \cdot \overline{n-8} \cdot \cdot \cdot 3 \cdot 1}{\overline{n-3} \cdot \overline{n-5} \cdot \overline{n-7} \cdot \overline{n-9} \cdot \cdot 2t^{n-3}} \times \int \frac{\dot{x}}{t^2-x^2} \text{ if } n \text{ be an odd number; but}$$

$$2.\ \int \frac{\dot{x}}{(t^2-x^2)^{\frac{n-1}{2}}} = \frac{1}{n-3t^2} \times \frac{x}{(t^2-x^2)^{\frac{n-3}{2}}} + \frac{n-4}{n-3 \cdot n-5t^4} \times \frac{x}{(t^2-x^2)^{\frac{n-5}{2}}}$$

$$+ \frac{\overline{n-4} \cdot \overline{n-6}}{\overline{n-3} \cdot \overline{n-5} \cdot \overline{n-7}t^6} \times \frac{x}{(t^2-x^2)^{\frac{n-7}{2}}} \cdot \cdot \cdot \cdot \cdot +$$

$$\frac{\overline{n-4} \cdot \overline{n-6} \cdot \overline{n-8} \cdot \overline{n-10} \cdot \cdot \cdot \cdot 4 \cdot 2}{\overline{n-3} \cdot \overline{n-5} \cdot \overline{n-7} \cdot \overline{n-9} \cdot \overline{n-11} \cdot \cdot \cdot \cdot 3 \cdot 1t^{n-2}} \times \frac{x}{(t^2-x^2)^{\frac{n-2}{2}}}, \text{ if } n \text{ be even.}$$

Theſe principles may be applied to the finding approximations in very many philoſophical problems.

Cor. 1. From hence may be deduced the ſubſequent arithmetical theorems.

$$1.\ \frac{1}{2m-1} + \frac{2m-2}{2m-1 \cdot 2m-3} + \frac{\overline{2m-2} \cdot \overline{2m-4}}{\overline{2m-1} \cdot \overline{2m-3} \cdot \overline{2m-5}} +$$

$$\frac{\overline{2m-2} \cdot \overline{2m-4} \cdot \overline{2m-6}}{\overline{2m-1} \cdot \overline{2m-3} \cdot \overline{2m-5} \cdot \overline{2m-7}} \cdot \cdot \cdot \cdot \cdot \frac{\overline{2m-2} \cdot \overline{2m-4} \cdot \overline{2m-6} \cdot \cdot \cdot \cdot 2}{\overline{2m-1} \cdot \overline{2m-3} \cdot \overline{2m-5} \cdot \overline{2m-7} \cdot \cdot \cdot 3 \cdot 1} =$$

1; or, which is the ſame, $\frac{1}{2m-3} + \frac{2m-4}{2m-3 \cdot 2m-5} +$

$2m-4$

$$\frac{2m-4\times 2m-6}{2m-3\times 2m-5\times 2m-7}\ldots+\frac{2m-4\cdot 2m-6\cdot 2m-8\ldots 4\times 2}{2m-3\cdot 2m-5\cdot 2m-7\cdot 2m-9\ldots 3\times 1}$$

$=1$, or $\frac{1}{2m-5}+\frac{2m-6}{2m-5\cdot 2m-7}\ldots+\frac{2m-6\cdot 2m-8\ldots 4\times 2}{2m-5\cdot 2m-7\cdot 2m-9\ldots 3\times 1}$

$=1$, &c.

2. $$\frac{1}{2m-2}+\frac{2m-3}{2m-2\cdot 2m-4}+\frac{2m-3\cdot 2m-5}{2m-2\cdot 2m-4\cdot 2m-6}+$$

$$\frac{2m-3\cdot 2m-5\cdot 2m-7}{2m-2\cdot 2m-4\cdot 2m-6\cdot 2m-8}\ldots+$$

$$\frac{2m-3\cdot 2m-5\cdot 2m-7\cdot 2m-9\ldots 7\cdot 5}{2m-2\cdot 2m-4\cdot 2m-6\cdot 2m-8\cdot 2m-10\ldots 6\cdot 4}+2\times$$

$$\frac{2m-3\cdot 2m-5\cdot 2m-7\cdot 2m-9\ldots 3}{2m-2\cdot 2m-4\cdot 2m-6\cdot 2m-8\cdot 2m-10\ldots 2}=1, \text{ or}$$

$$\frac{1}{2m-4}+\frac{2m-5}{2m-4\cdot 2m-6}+\frac{2m-5\cdot 2m-7}{2m-4\cdot 2m-6\cdot 2m-8}\ldots+$$

$$\frac{2m-5\cdot 2m-7\cdot 2m-9\ldots 5}{2m-4\cdot 2m-6\cdot 2m-8\ldots 4}+2\times$$

$$\frac{2m-5\cdot 2m-7\cdot 2m-9\cdot 2m-11\ldots 5\cdot 3}{2m-4\cdot 2m-6\cdot 2m-8\cdot 2m-10\cdot 2m-12\ldots 4\cdot 2}=1, \text{ &c.}$$

3. By expanding the terms of the preceding fluents may be deduced the arithmetical theorems, *viz.*

1. $$\frac{1}{n-3}\times n-3\times n-1\cdot n+1\cdot n+3\ldots n+m+\frac{n-4}{n-3\cdot n-5}$$

$$\times n-5\cdot n-3\cdot n-1\cdot n+1\ldots n+m-2+\frac{n-4\cdot n-6}{n-3\cdot n-5\cdot n-7}$$

$$\times n-7\cdot n-5\cdot n-3\cdot n-1\ldots n+m-4+$$

$$\frac{n-4\cdot n-6\cdot n-8}{n-3\cdot n-5\cdot n-7\cdot n-9}\times n-9\cdot n-7\cdot n-5\cdot n-3\ldots n+m-6$$

$$+\ldots+\frac{n-4\cdot n-6\cdot n-8\cdot n-10\ldots 3}{n-3\cdot n-5\cdot n-7\cdot n-9\cdot n-11\ldots 2}\times 2\cdot 4\cdot 6\cdot 8\ldots$$

$$m+5+\frac{n-4\cdot n-6\cdot n-8\cdot n-10\ldots 3}{n-3\cdot n-5\cdot n-7\cdot n-9\cdot n-11\ldots 2}\times 2\times 4\times 6\times 8\ldots$$

$m+5$

$\overline{m+5} \times \frac{1}{m+6} = \overline{n-1} \cdot \overline{n+1} \cdot \overline{n+3} \cdot \overline{n+5} \ldots \overline{n+m+2} \times \frac{1}{m+6}$ if n be an odd number.

2. $\frac{1}{n-3} \times \overline{n-3} \cdot \overline{n-1} \cdot \overline{n+1} \cdot \overline{n+3} \ldots \overline{n+2m+1} +$
$\frac{n-4}{n-3 \cdot n-5} \times \overline{n-5} \cdot \overline{n-3} \cdot \overline{n-1} \cdot \overline{n+1} \ldots \overline{n+2m-1} +$
$\frac{\overline{n-4} \cdot \overline{n-6}}{n-3 \cdot n-5 \cdot n-7} \times \overline{n-7} \cdot \overline{n-5} \cdot \overline{n-3} \cdot \overline{n-1} \ldots \overline{n+2m-3} +$
$\frac{\overline{n-4} \cdot \overline{n-6} \cdot \overline{n-8}}{n-3 \cdot n-5 \cdot n-7 \cdot n-9} \times \overline{n-9} \cdot \overline{n-7} \cdot \overline{n-5} \cdot \overline{n-3} \ldots \overline{n+2m-5}$
$\ldots + \frac{\overline{n-4} \cdot \overline{n-6} \cdot \overline{n-8} \ldots 2}{n-3 \cdot n-5 \cdot n-7 \cdot n-9 \ldots 3 \times 1} \times 1 \cdot 3 \cdot 5 \cdot 7 \ldots \overline{2m+5} =$
$\overline{n-1} \cdot \overline{n+1} \cdot \overline{n+3} \cdot \overline{n+5} \ldots \overline{n+2m+3} \times \frac{1}{2m+7}$, if n be an even number.

4. Arithmetical theorems, somewhat different, may be deduced from taking the fluxions of the preceding fluents, and reducing the fractions resulting to a common denominator by multiplying them into $t^2 - x^2$, $(t^2 - x^2)^2$, $(t^2 - x^2)^3$, &c.

1. $\frac{1}{n-3} \times \overline{n-4} + \frac{n-4}{n-3 \cdot n-5} \times \overline{n-7} + \frac{\overline{n-4} \cdot \overline{n-6}}{n-3 \cdot n-5 \cdot n-7} \times \overline{n-10}$
$+ \frac{\overline{n-4} \cdot \overline{n-6} \cdot \overline{n-8}}{n-3 \cdot n-5 \cdot n-7 \cdot n-9} \times \overline{n-13} \ldots + \frac{\overline{n-4} \cdot \overline{n-6} \cdot \overline{n-8} \ldots 3}{n-3 \cdot n-5 \cdot n-7 \cdot n-9 \ldots 2}$
$\times \frac{7-n}{2} - \frac{\overline{n-4} \cdot \overline{n-6} \cdot \overline{n-8} \cdot \overline{n-10} \ldots 3}{n-3 \cdot n-5 \cdot n-7 \cdot n-9 \cdot n-11 \ldots 2} \times \frac{n-3}{2} = 0$, if n be odd.

2. $\frac{1}{n-3} \times \overline{n-4} + \frac{n-4}{n-3 \cdot n-5} \times \overline{n-7} + \frac{\overline{n-4} \cdot \overline{n-6}}{n-3 \cdot n-5 \cdot n-7} \times \overline{n-10}$
$+ \frac{\overline{n-4} \cdot \overline{n-6} \cdot \overline{n-8}}{n-3 \cdot n-5 \cdot n-7 \cdot n-9} \times \overline{n-13} \ldots + \frac{\overline{n-4} \cdot \overline{n-6} \cdot \overline{n-8} \ldots 2}{n-3 \cdot n-5 \cdot n-7 \cdot n-9 \ldots 3 \times 1}$
$\times \frac{4-n}{2} = 0$, if n be even; or more general, if n be even,

3. $\frac{n-4 \cdot n-6 \cdot n-8 \ldots n-2m-2}{n-3 \cdot n-5 \cdot n-7 \cdot n-9 \ldots n-2m-3} \times n-2m-4 +$

$\frac{n-4 \cdot n-6 \cdot n-8 \cdot n-10 \ldots n-2m-4}{n-3 \cdot n-5 \cdot n-7 \cdot n-9 \cdot n-11 \ldots n-2m-5} \times (m+1 \times n-2m-6-1)$

$+ \frac{n-4 \cdot n-6 \cdot n-8 \cdot n-10 \cdot n-12 \ldots n-2m-6}{n-3 \cdot n-5 \cdot n-7 \cdot n-9 \cdot n-11 \cdot n-13 \ldots n-2m-7} \times$

$(m+1 \cdot \frac{m+2}{2} \times n-2m-8-m+2) +$

$\frac{n-4 \cdot n-6 \cdot n-8 \cdot n-10 \cdot n-12 \cdot n-14 \ldots n-2m-8}{n-3 \cdot n-5 \cdot n-7 \cdot n-9 \cdot n-11 \cdot n-13 \cdot n-15 \ldots n-2m-9}$

$\times (m+1 \cdot \frac{m+2}{2} \cdot \frac{m+3}{3} \times n-2m-10-m+2 \cdot \frac{m+3}{2}) +$

$\frac{n-4 \cdot n-6 \cdot n-8 \cdot n-10 \ldots n-2m-10}{n-3 \cdot n-5 \cdot n-7 \cdot n-9 \cdot n-11 \ldots n-2m-11} \times (m+1 \cdot \frac{m+2}{2} \cdot \frac{m+3}{3} \cdot$

$\frac{m+4}{4} \times n-2m-12-m+2 \cdot \frac{m+3}{2} \cdot \frac{m+4}{3}) \ldots\ldots +$

$\frac{n-4 \cdot n-6 \cdot n-8 \ldots 2}{n-3 \cdot n-5 \cdot n-7 \ldots 3 \times 1} \times -m+2 \cdot \frac{m+3}{2} \cdot \frac{m+4}{3} \ldots\ldots \frac{n-4}{n-2m-6}$

$= 0.$

4. $\frac{n-4 \cdot n-6 \cdot n-8 \cdot n-2m-2}{n-3 \cdot n-5 \cdot n-7 \cdot n-9 \ldots n-2m-3} \times n-2m-4 +$

$\frac{n-4 \cdot n-6 \cdot n-8 \ldots n-2m-4}{n-3 \cdot n-5 \cdot n-7 \cdot n-9 \ldots n-2m-5} \times (m+1 \times n-2m-6-1) +$

$\frac{n-4 \cdot n-6 \cdot n-8 \ldots n-2m-6}{n-3 \cdot n-5 \cdot n-7 \cdot n-9 \ldots n-2m-7} \times (m+1 \cdot \frac{m+2}{2} \times n-2m-8 -$

$m+2) + \frac{n-4 \cdot n-6 \cdot n-8 \ldots n-2m-8}{n-3 \cdot n-5 \cdot n-7 \cdot n-9 \ldots n-2m-9} \times (m+1 \cdot \frac{m+2}{2} \times$

$\frac{m+3}{3} \times n-2m-10-m+2 \cdot \frac{m+3}{2}) + \ldots\ldots\ldots +$

$\frac{n-4 \cdot n-6 \cdot n-8 \ldots 5 \times 3}{n-3 \cdot n-5 \cdot n-7 \cdot n-9 \ldots 4 \times 2} \times (m+1 \cdot \frac{m+2}{2} \cdot \frac{m+3}{3} \ldots \frac{n-5}{n-2m-5} -$

$m+2 \cdot \frac{m+3}{2} \cdot \frac{m+4}{3} \ldots \frac{n-5}{n-2m-7}) -$

$n-4$

$$\frac{\overline{n-4}\cdot\overline{n-6}\cdot\overline{n-8}\ldots 5\times 3}{n-3\cdot n-5\cdot n-7\cdot n-9\ldots 4\times 2}\times\overline{m+2}\cdot\frac{m+3}{2}\times\frac{m+4}{3}\ldots\ldots$$

$\frac{n-3}{n-2m-5}=0$, if n be an odd number.

Many more arithmetical theorems may be deduced from the fluents of these and other fluxions by similar methods, which cannot, without some difficulty, be found from the common methods of finding the sums of series.

This method of finding approximations to the areas and lengths of curves, fluents of fluxions, and sums of series, &c. of which the equations, to their increments, fluxions, &c. are given from the areas of curves, fluents, &c. of which the equations to their increments, &c. differ by very small quantities from the given equations was published in the Meditat. Analyt. near twenty years ago.

I shall conclude this Paper with two theorems of some little use in the doctrine of chances.

THEOREM I.

$$H=\overline{a+b}\times\overline{a+b-1}\cdot\overline{a+b-2}\cdot\overline{a+b-3}\ldots\overline{a+b-n+1}$$

$$=a\cdot\overline{a-1}\cdot\overline{a-2}\ldots\overline{a-n+1}+n\cdot a\cdot\overline{a-1}\cdot\overline{a-2}\ldots\overline{a\ n+2}$$

$$\times b+n\cdot\frac{n-1}{2}\times a\cdot\overline{a-1}\cdot\overline{a-2}\ldots\overline{a-n+3}\times b\cdot\overline{b-1}+n\cdot$$

$$\frac{n-1}{2}\cdot\frac{n-2}{3}a\cdot\overline{a-1}\cdot\overline{a-2}\ldots\overline{a-n+4}\times b\times\overline{b-1}\cdot\overline{b-2}+$$

$$\ldots\ldots+n\cdot\frac{n-1}{2}\cdot\frac{n-2}{3}\cdot\frac{n-3}{4}\ldots\frac{n-l+1}{l}\text{(L)}\,a\cdot\overline{a-1}\cdot\overline{a-2}$$

$$\cdot\overline{a-3}\ldots\overline{a-n+l+1}\times b\cdot\overline{b-1}\cdot\overline{b-2}\ldots\overline{b-l+1}+\ldots+$$

$n \cdot \frac{n-1}{2} a \cdot \overline{a-1} \cdot b \cdot \overline{b-1} \cdot \overline{b-2} \ldots \overline{b-n+3} + na \cdot b \cdot \overline{b-1} \cdot \overline{b-2} \ldots \overline{b-n+2} + \frac{n-1}{2} b \cdot \overline{b-1} \cdot \overline{b-2} \ldots \overline{b-n+1}$.

If for $a+b-1$, $a+b-2$, $a+b-3$, &c., $a-1$, $a-2$, &c., $b-1$, $b-2$, &c. be substituted respectively $a+b-x$, $a+b-2x$, $a+b-3x$, &c., $a-x$, $a-2x$, $a-3x$, &c., $b-x$, $b-2x$, $b-3x$, &c., the resulting equation will equally be just; and, lastly, if for x be substituted 0, it will become the binomial theorem.

Cor. If there are two different events A and B, of which the numbers are respectively a and b, and their chances of happening also as a and b; and if A's happen, let the whole number $(a+b)$ and also the number of A's be diminished by x, and in the same manner of B's happening, and so on; then will the chance of A's happening $n-l$ times, and B's happening l times in n trials be $L \times a \cdot \overline{a-x} \cdot \overline{a-2x} \ldots \overline{a-(n-l-1)x} \times b \cdot \overline{b-x} \cdot \overline{b-2x} \ldots \overline{b-(l-1)x}$ divided by H.

In a similar manner may be found, 1. the chance of A's happening between h and k times; and, 2. the chance of A's happening (h) to B's happening (k) times; 3. of A's and B's happening respectively h and k times more than the other; 4. the chance of A's happening an even to its happening an odd number of times, &c. in (n) trials, &c. &c. &c.

THEOREM II.

$H = \overline{a+b+c+d+\&c.} \times \overline{a+b+c+d+\&c.-x} \times \overline{a+b+c+d+\&c.-2x} \ldots \overline{a+b+c+d+\&c.-(n-1)x} = a \cdot \overline{a-x} \cdot \overline{a-2x} \ldots \overline{a-n-1x+n} \cdot a \cdot \overline{a-x} \cdot \overline{a-2x} \ldots \overline{a-n-2x} \times b+c+d+\&c.$

+

$+n \cdot \frac{n-1}{2} \cdot a \cdot \overline{a-x} \cdot \overline{a-2x} \cdot \overline{a-3x} \times (b \cdot \overline{b-x} + c \times \overline{c-x}$
$+d \cdot \overline{d-x} + \&c. + 2bc + 2bd + 2cd + \&c.) + \ldots + L \times (a \cdot$
$\overline{a-x} \cdot \overline{a-2x} \ldots \overline{a-l-1x} \times b \cdot \overline{b-x} \cdot \overline{b-2x} \ldots \overline{b-m-1x}$
$\times c \cdot \overline{c-x} \cdot \overline{c-2x} \ldots \overline{c-p-1x} \times d \cdot \overline{d-x} \cdot \overline{d-2x} \ldots$
$\overline{d-q-1x} \times \&c. = K) + \&c.$, where $L = n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} \ldots$
$\frac{n-l+1}{l} \times \overline{n-l} \cdot \frac{n-l-1}{2} \cdot \frac{n-l-2}{3} \ldots \frac{n-l-m+1}{m} \times \overline{n-l-m} \cdot$
$\frac{n-l-m-1}{2} \cdot \frac{n-l-m-2}{3} \ldots \frac{n-l-m-p+1}{p} \times \overline{n-l-m-p} \cdot$
$\frac{n-l-m-p-1}{2} \cdot \frac{n-l-m-p-2}{3} \ldots \frac{n-l-m-p-q+1}{q} \times \&c.$ which is the same as the co-efficient of the term $a^l \times b^m \times c^p \times d^q \times \&c.$ in the multinomial $a + b + c + d + \&c.$ raised to the power n.

The chance of any number of events A, B, C, D, &c. of which the numbers are a, b, c, d, &c. happening l, m, p, q, &c. times respectively in a similar manner to A's and B's happening in the preceding case will be $\frac{L \times K}{H}$.

All the propositions mentioned as immediately deducible from the preceding theorem may, *mutatis mutandis*, with the same ease be applied to more events A, B, C, D, &c.

If for a, b, c, d, &c. be substituted the same letters, increased or diminished by any given quantities, the resulting equation will be equally true.

ERRATA.

VOL. LXXIX.

Page 171. *after* $+n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} \log. \overline{r-3p}$ + &c. *is omitted* but $\log. r-n \times \log. \overline{r-p} + n \cdot \frac{n-1}{2} \log. \overline{r-2p} - n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} \times \log. \overline{r-3p}$ + &c. =, which probably happened, in my abſence from the preſs, on account of the ſimilarity of the preceding quantity.

Page 172. l. 1. *for* (±M + *read* + (M±

Fig 1.



Fig 2.





Fig 3.

Fig 4.

Figure 2.

X. *An Account of ſome Appearances attending the Converſion of caſt into malleable Iron. In a Letter from* Thomas Beddoes, *M. D. to Sir* Joſeph Banks, *Bart. P. R. S.*

Read March 24, 1791.

SIR,

YOU are undoubtedly well apprized of an alteration lately introduced into our manufactories of iron, in conſequence of which the reverberatory has been ſubſtituted in the place of the finery furnace. The new proceſs is capable of being indefinitely varied. I have lately been favoured with an opportunity of obſerving one of theſe variations with every advantage I could deſire. As in this method the changes undergone by the metal during the firſt ſeries of operations lie perfectly open to inſpection, a ſhort deſcription of them may not perhaps be unworthy the notice of philoſophical chemiſts. Allow me to premiſe further, that I did not content myſelf with a ſingle examination; and, for the ſake of greater accuracy, I took minutes of the phænomena, and of the time when they occurred. A very intelligent workman was at the ſame time directed to anſwer all my queſtions, ſo that I enjoyed the benefit of his experience alſo.

In ſomewhat more than half an hour after it was put in, the charge conſiſting of 2¼ cwt. of grey pig iron was nearly melted. The workman now began to ſtir the liquid maſs: for

for this purpoſe he uſed ſometimes an iron lever, and ſometimes a kind of hoe; but he firſt turned the flame from off the metal, which is done by letting down a damper upon the chimney correſponding to that with which ordinary reverberatory furnaces are provided, and by raiſing the damper of a ſecond chimney, which proceeds immediately from the fire-place, and carries off the flame, current of air, &c. without allowing it paſs into the body of the furnace.

In 50 minutes from the commencement of the operation, the metal had become in conſequence of the conſtant ſtirring looſe and incoherent; it appeared about as ſmall as gravel; it was now alſo ſtiff, and much cooled.

55 m. from the ſame period, flame turned on again. Workman keeps ſtirring and turning over the metal; in 3 m. it becomes ſoft and ſemi-fluid; flame turned off; the hotteſt part of the maſs begins to heave and ſwell, emitting a deep blue lambent flame. The workman calls this appearance fermentation.

1 hour 1 m. blue flame breaking out over the whole maſs; heaving motion alſo general.

1 h. 13 m. metal full as hot, or, as the workman and myſelf both judged, rather hotter than at the inſtant the flame was turned off, though it is now a quarter of an hour ſince.

1 h. 18 m. where there is no heaving and no blue flame the maſs is ſenſibly cooler, and only of a dull red heat.

1 h. 20 m. workman obſerves, that the metal ſticks leſs to his tools. Pig-iron, he ſays, faſtens upon it immediately, and muſt be ſhaken off by ſtriking the other end with an hammer; as it approaches more and more towards *nature* (malleable iron) it adheres leſs; and when the tools come clear up out of the maſs, he judges it to be fermented enough.

1 h. 23 m. little heaving or blue flame; metal ſtiffer, and of a dull red; flame turned on and ſoon off again.

1 h. 26 m. by conſtant ſtirring the metal is become as fine as ſand. Workman remarks, that the flame, which reappears over the whole maſs, looks more kindly. It is evidently of a lighter blue colour.

1¼ h. flame turned on and ſoon off again. Maſs ferments ſtrongly. Hiſſing noiſe heard: this noiſe was diſtinguiſhable in ſome degree ever ſince the blue flame and heaving motion became viſible, but always faint till now.

1 h. 40 m. leſs blue flame.

1 h. 48 m. flame twice turned on and off in this interval. Metal now clots, ſtands wherever it is placed, without any tendency to flow, and no liquid pig iron now remains in the baſon of the furnace; the maſs has been conſtantly ſtirred and turned over.

1 h. 50 m. a little finery cinder appears boiling up amid the maſs. Workman attributes the increaſe of the hiſſing to this.

1 h. 53 m. ſcarce any perceptible blue flame or heaving. All the metal is now gathered into lumps, which the workman beats and preſſes with an heavy-headed tool. He brings them ſucceſſively into the hotteſt part of the furnace, into which the flame has been admitted. He now ſtops the port hole in the door at which he had introduced his tools, and applies a fierce flame for 6 or 8 minutes; the metal is then rolled.

Theſe appearances, at leaſt the moſt intereſting of them, ſeem to admit of an eaſy explanation; and I offer the following obſervations as ſupplemental to thoſe for which we are already indebted to the Swediſh and French chemiſts on this important branch of metallurgy. I aſſume the following propoſitions as already proved by theſe philoſophers. 1. That caſt iron

iron is iron imperfectly reduced, or, in other words, that it contains a portion of the basis of vital air, the oxygène of M. LAVOISIER. 2. That it contains a portion of plumbago, with which grey cast iron most abounds. 3. That plumbago consists of iron united to charcoal. 4. That fixed air, which I would rather call carbonic acid air, consists of oxygène and the constituent parts of charcoal.

The heaving or swelling motion, so conspicuous in the process, is doubtless owing to the discharge of an elastic fluid; and the lambent deep blue flame, breaking out in spots over the whole surface, shews, that this elastic fluid is an inflammable gas of the heavy kind. That no doubt might be left upon the former of these circumstances, I directed the workman to take out, at two different periods, a quantity of the metal where it was working most strongly. Both proved, on examination, to be spungy, cellular, and full of bladder holes.

The heavy inflammable air, I imagine, is produced in this manner. The oxygène of the imperfectly reduced metal combines with the charcoal to form fixed air; at the same time another portion of charcoal is thrown into an elastic state, that is, into inflammable air, and burns on the surface with a very deep blue flame, on account of the admixture of fixed air. The heat which is so obviously generated in the mass at the beginning of the fermentation, I attribute to the combination of the oxygène and charcoal; a fact which, with several others as I have already remarked on another occasion *, shews, if not the falsehood, at least the imperfection of the modern doctrine on the subject of heat. The acidifying principle, it would appear, has some power of generating heat independant of its condensation. Here abundance of elastic matter is

* Chemical opinions of a philosopher of the last century.

dis-

diſcharged; yet, notwithſtanding the heat abſorbed by its formation, and that which flows out of the metal in all directions, the whole maſs becomes hotter. The oxygène cannot be ſuppoſed to have much ſpecific or latent heat, becauſe it undoubtedly exiſts in the iron in a very condenſed ſtate. Neither does the appearance of the maſs allow me to aſcribe this generation of heat to the burning of the inflammable air at the ſurface, as will alſo be immediately evident for another reaſon. The leſs deep blue colour of the flame at a ſubſequent period in the operation is probably owing to the abſence of fixed air, or at leaſt to its being produced more ſparingly, the oxygène being now nearly conſumed. It will not appear ſurprizing, that the oxygène in this caſe ſhould be conſumed before the charcoal, if it be conſidered, 1. that grey iron contains a large portion of plumbago; and, 2. that fixed air contains a much larger quantity of oxygène than of charcoal; near three times as much, according to our beſt experiments on its formation: ſo that I aſcribe the ſubſequent fermentation accompanied with the lighter coloured flame almoſt entirely to the converſion of the charcoal into an elaſtic fluid. A very experienced philoſopher, I am well aware, has aſſerted, that water is neceſſary to this converſion; an opinion concerning the juſtneſs of which I have long entertained great doubts. Whenever I have diſtilled charcoal *per ſe*, I have found the firſt portions of gas to contain fixed air; an appearance owing, as I believe, to the decompoſition of water abſorbed from the atmoſphere; but, after continuing the proceſs for ſome time, there has ſtill been a production of inflammable air; but from this neither lime-water nor milk of lime would abſorb any portion, though when fired with vital or common air, it would produce fixed air; and if moiſture was added to the charcoal, inflammable

and fixed air would be generated anew. Moreover it appears, from the experiments of Dr. AUSTIN and ſome others, that charcoal conſiſts of the hydrogène and azote of the French chemiſts. How far it may be difficult or impoſſible entirely to convert charcoal in its ordinary ſtate into gas, is a point I wiſh to ſee more fully illuſtrated by future experiments. At preſent it ſeems obvious, that the circumſtances of the operation I have deſcribed are particularly favourable to this converſion: for, 1. not to mention the violence of the heat, we have this ſubſtance in a very attenuated ſtate, ſo that, very probably, the expanſive power of fire is very little, if at all, counteracted by the attraction of coheſion, which cannot be ſaid in the caſe of the moſt minute mechanical diviſion we can effect. 2. The attraction of the particles of the iron for one another will produce an effort to extrude the intermixed particles of charcoal, and thus enable it more readily to aſſume the elaſtic form.

Now, during the continuance of the lighter coloured blue flame, the maſs, as I obſerved, ſhews no power of generating heat within itſelf; a circumſtance which indicates that the heat produced in the former part of the operation does not depend on the burning of the gas at the ſurface; and I think inſpection will ſatisfy any one that it is produced in the heart of the maſs. It may indeed be objected, that the metal, now brought nearer to the ſtate of malleable iron, may require a greater ſupply of heat to keep it at the ſame temperature. It is leſs fuſible, as we are well aſſured. By referring back to the minutes you will obſerve, how very often it was neceſſary to turn the flame upon the maſs during this ſecond fermentation in order to keep it in a ſtate in which it could be worked.

The very copious production of elaſtic fluids during an hour, and often during a much longer ſpace, for in this inſtance the

proceſs was remarkably ſucceſsful and ſhort, does not ſeem favourable to a late ingenious hypotheſis, according to which water is the embodying principle of all elaſtic fluids. I have never indeed conſidered this as very probable, and, after the obſervations I have related, I ſee no means of defending it. Will it be ſaid, that the pig iron, as being in ſome ſort a calx of iron, contains water?

In annealing crude iron, with or without charcoal, it is well known to increaſe in all its dimenſions. I have ſeen bars originally ſtraight bent like an S, when long expoſed to heat in circumſtances where they could not extend themſelves endways. I ſuppoſe this phænomenon may be owing to a very ſmall beginning of this fermentative motion, which acts as an internal principle of expanſion. Caſt iron bars, not in contact with charcoal, would, according to this ſuppoſition, by long annealing loſe of their weight; or if the heat was too low for the elaſtic fluid to be diſcharged from their ſubſtance, they would probably bliſter like ſteel: an appearance undoubtedly owing to the generation of air. Mr. HORNE, in his Eſſay on Iron, ſomewhere remarks, that on opening theſe bliſters he has heard a whiſtling noiſe as of air ruſhing out.

During the whole of this proceſs, frequent jets of white ſparks, of a dazzling brightneſs, played from the ſurface of the metal. They would have afforded an extremely beautiful ſpectacle but for the inconvenience of looking on ſo hot a maſs. They aroſe, no doubt, from the burning of ſmall portions of iron.

The effect of ſo much ſtirring as I have noted down does not require to be explained.

The workman was clearly of opinion, that the fermentation of hard or white crude iron is leſs than of grey in this

proceſs; a fact which perfectly coincides with the preceding obſervations, ſince that ſpecies contains leſs plumbago, or in other words leſs matter fit to produce elaſtic fluids.

In order to prove the extrication of fixed air during the fermentation of the metal, I once thought of introducing lime-water in an iron veſſel within the body of the furnace; but when I conſidered that the fire-place was not divided by any partition from the body of the furnace, and that the whole building was full of burned air, I omitted the experiment from a perſuaſion that, even if the lime-water ſhould become turbid, the fixed air might come from another ſource.

I was not unmindful of the ſulphur which exiſts, as I have reaſon to believe, in every form of iron manufactured with coaks. I cannot, however, aſcribe any of the effects I obſerved to its preſence. There can be little doubt, that ſome portion was perpetually extricated with the inflammable air during the whole proceſs; for on diſſolving pieces of the ſtamped, or rather the rolled iron in weak muriatic acid, ſilver held in the extricated air was tarniſhed as much and as ſoon as by air from ſpecimens taken out of the furnace at different times during the proceſs. I could not but conclude, that the tarniſhing matter came from the iron, when I found the air from a ſolution of zinc in the ſame acid, incapable of producing the colour upon ſilver. The appearance, the want of a martial aſtringent taſte, and the diſſolving action of cauſtic alkali, led me to conclude, that the colour in each experiment with iron was derived from ſulphur.

I leave it to the adherents of phlogiſton to accommodate theſe phænomena to their doctrine; conſidering it, for my own part, as ſuperfluous to beſtow any further attention upon a

system which, after a long discussion, has been fully refuted in all its modifications, and which indeed seems on the eve of being universally abandoned.

I have the honour to be, &c.

THOMAS BEDDOES.

XI. *On the Decompoſition of Fixed Air.*

By Smithſon Tennant, *Eſq. F. R. S.*

Read March 31, 1791.

AS fixed air is produced by the combuſtion of charcoal, it has long been thought highly probable that vital air and charcoal are its conſtituent ingredients. This opinion is confirmed by the experiments of M. Lavoisier, from which he diſcovered that the weight of the fixed air which is formed during the combuſtion is nearly equal to that of the vital air and charcoal conſumed in the proceſs; and that the ſmall difference of weight may, with great reaſon, be attributed to the production of water ariſing from inflammable air contained in the charcoal. The compoſition of fixed air therefore ſeems to be determined, by uniting its conſtituent parts, with as much certainty as by that mode of proof alone it is poſſible to obtain. But as vital air has a ſtronger attraction for charcoal than for any other known ſubſtance, the decompoſition of fixed air has not hitherto been attempted. By means, however, of the united force of *two* attractions I have been able to decompoſe fixed air, and thus to determine its conſtituent parts in conſequence of their ſeparation.

It has long been known, that when phoſphoric acid is combined with calcareous earth, it cannot be decompoſed by diſtillation with charcoal: for though vital air is more ſtrongly

attracted

attracted by charcoal than by phoſphorus, yet in this compound it is retained by two attractions, by that which it has for phoſphorus, and by that which the phoſphoric acid has for lime, ſince the vital air cannot be diſengaged unleſs both theſe attractions are overcome. As theſe attractions are more powerful than that which charcoal has for vital air, if phoſphorus is applied to fixed air and calcareous earth, the vital air will unite with the phoſphorus, and the charcoal will be obtained pure. Theſe ſubſtances, in order to act upon each other, muſt be brought into contact when red-hot; and this may be eaſily effected in the following manner. Into a glaſs tube, cloſed at one end, and coated with ſand and clay to prevent the ſudden action of the heat, a little phoſphorus ſhould be firſt introduced, and afterwards ſome powdered marble. The experiment ſucceeds more readily if the marble is ſlightly calcined, probably becauſe that part which is reduced to lime, by immediately uniting with the phoſphorus, detains it to act upon the fixed air in the other part. After the ingredients are introduced, the tube ſhould be nearly, but not entirely, cloſed up; by which means ſo free a circulation of air as might inflame the phoſphorus is prevented, whilſt the heated air within the tube is ſuffered to eſcape. When the tube has remained red-hot for ſome minutes, it may be taken from the fire, and muſt be ſuffered to grow cold before it is broken. It will be found to contain a black powder, conſiſting of charcoal intermixed with a compound of lime and phoſphoric acid, and of lime united with phoſphorus. The lime and phoſphoric acid may be ſeparated by ſolution in an acid and by filtration, and the phoſphorus by ſublimation.

Charcoal, thus obtained from fixed air, appears in no reſpect to differ from the charcoal of vegetable matters. On deflagrating

ting a little of it in a ſmall retort with nitre, fixed air was immediately reproduced.—Since, therefore, charcoal, by its ſeparation from fixed air, is proved to be one of its conſtituent principles, it can hardly be doubted, that this ſubſtance is preſent whenever fixed air is produced; and that thoſe experiments, from which it is ſuppoſed that this acid may be formed without the aid of charcoal, have not been conducted with the requiſite caution.

As vital air is attracted by a compound of phoſphorus and calcareous earth more powerfully than by charcoal, I was deſirous of trying their efficacy upon theſe acids, which may from analogy be ſuppoſed to contain vital air, but which are not affected by the application of charcoal. With this intention I made phoſphorus paſs through a compound of marine acid and calcareous earth, and alſo of fluor acid and calcareous earth, but without producing in either of them any alteration. Since the ſtrong attraction which theſe acids have for calcareous earth tends to prevent their decompoſition, it might be thought that in this manner they were not more diſpoſed to part with vital air than by the attraction of charcoal. But this, however, does not appear to be the fact. I have found, that phoſphorus cannot be obtained by paſſing marine acid through a compound of bones and charcoal, when red-hot. The attraction, therefore, of phoſphorus and lime for vital air exceeds the attraction of charcoal by a greater force than that ariſing from the attraction of marine acid for lime.

XII. *A Meteorological Journal, principally relating to Atmoſpheric Electricity; kept at* Knightſbridge, *from the 9th of* May, 1789, *to the 8th of* May, 1790. *By Mr.* John Read; *communicated by* R. H. A. Bennet, *Eſq. F. R. S.*

Read April 14, 1791.

A DESCRIPTION of the inſtrument for collecting atmoſpheric electricity, uſed in the following journal.

Tab. V. repreſents the apparatus. AA is a round deal rod, 20 feet long, 2 inches diameter at the lower, and one inch at the upper end. Into the lower end of it is cemented a ſolid glaſs pillar B, 22 inches long; the lower end of the glaſs ſtands in a hole made for it in a pedeſtal of wood C, which ſlips on the fore-part of an iron bracket D, which is driven into the wall, and ſupports the whole. About 13 feet above the bracket D, is fixed to the wall a ſtrong arm of wood E, which holds perpendicularly a ſtrong glaſs tube F, through which the rod is ſlided gently upwards, till the glaſs pillar B may be lowered into the hole made for it in C. It is thus fixed, and ſtands 12 inches from the wall. The tube F is of ſufficient width to admit a caſe of cork, which is faſtened in the inſide of it, at the part where the tube is ſuſtained by the arm of wood E, ſo that the rod, when bent by the wind, cannot touch the tube or break it. The upper extremity of the rod is terminated by ſeveral ſharp-pointed wires G. Two of them are

of copper, each one-eighth of an inch thick; and, in order to ſtiffen the rod, as well as conduct more readily the electric fluid, one of thoſe wires is twiſted round the rod to the right hand, and the other to the left, as low down as the braſs collar at the vertex of the lower funnel H, to which they are ſoldered, in order to render their contact perfect. The tin funnels HH ſerve to defend the glaſſes B and F from the weather, which glaſſes are alſo covered with ſealing-wax to render their inſulation more perfect. At a convenient height from the floor, a hole is bored through the wall at I. This hole receives a glaſs tube covered with ſealing-wax, through which a ſtrong braſs wire proceeding from the rod is conveyed into the room, where juſt at the end of the glaſs tube it paſſes through a two-inch braſs ball L, and proceeding a little farther, keeps ſuſpended at its extremity a pith ball electrometer K, ſo that the electrometer may be about twelve inches diſtant from the wall. On the outſide of the wall there is a wooden box M, to keep that end of the glaſs tube dry.

At two inches diſtance of the above-mentioned braſs ball L, a bell N is ſupported by a ſtrong wire, which paſſing through another hole made in the wall, is made to communicate, by means of a good metallic continuation R, with the moiſt ground adjoining to the houſe. A braſs ball, three-tenths of an inch in diameter, is ſuſpended between the bell N and ball L, by a ſilk thread faſtened to a nail O. This ball ſerves for a clapper, by ſtriking between the ball and bell, when the electrical charge of the rod is ſufficiently ſtrong.

P is a ſmall table fixed to the wall under the bell and ball, at a convenient height above the floor, upon which Leyden bottles and other apparatus are occaſionally placed. Any perſon verſed in the ſcience of electricity, will eaſily underſtand that this

apparatus is calculated to ſhew the various degrees of atmoſpherical electricity, and at the ſame time to avoid the pernicious effects which may be occaſioned by thunder-ſtorms, or in ſhort by any great quantity of electricity in the atmoſphere.

The whole perpendicular height of both parts taken together, from the moiſt earth to the uppermoſt point at the top of the rod, is 52 feet.

Finding, however, that, notwithſtanding all the precaution I had taken to procure a good inſulation, the moiſt vapour of the atmoſphere, fixing upon the inſulating parts of the apparatus, rendered it imperfect in moiſt weather; I have lately (15th of Sept. 1790) altered the ſituation of the ſame rod, ſo that all the inſulating parts are now within the roof of the houſe. This I have effected by a hole through the roof of my houſe; by which means I now obtain a conſiderably more conſtant electricity; which, however, muſt not be ſolely attributed to the ſuperiority of my preſent mode of inſulating, but to the rod's being alſo elevated to the additional height of nine feet; ſo that I conſider its pointed part to be at preſent 61 feet above the moiſt earth.

This improvement of the apparatus, having been made after the concluſion of this journal, will be particularly deſcribed in the next, which I am now carefully continuing.

It will be neceſſary juſt to mention the method I have purſued in forming the journal of atmoſpheric electricity. This has been principally by means of the ſigns exhibited by the pith balls K, connected with the rod. When I find theſe cloſed, and not attracted by my finger, I then write no ſigns of electricity. When attracted on the approach of my finger, yet not ſufficiently charged to repel each other, I write weak ſigns of the fluid. When I find the balls open, and, on the approach of excited glaſs, the balls cloſe, I write they are electrified poſi-

tively; but, if the balls open wider, I write they are electrified negatively; and the reverſe when I uſe ſealing-wax. When the balls diverge one inch and upwards, viſible ſparks may be drawn at the braſs ball L. When ſparks are ſaid to have been perceived in any obſervation, I have generally on that account omitted to note the variable quantities of divergency in the pith balls. Their utmoſt limit of regular divergency ſeems to be about five or near ſix inches; above that they are unſteady and diſorderly. The pith balls are near two-tenths of an inch in diameter, ſuſpended by very fine flaxen threads (in the ſtate it is in from the heckle) five inches long. When I mention the diſtance of the balls in tenths of an inch, it is to be underſtood as nearly ſo as my eye can determine.

This apparatus requires a conſtant attention, eſpecially during a diſturbed ſtate of the atmoſphere. From the room in which the apparatus is placed I am ſeldom abſent one hour, excepting the time of ſleep; but, when I leave it, the laſt thing I do at night is to examine the ſtate of the electricity, and, if I find the rod unelectrified, I then place the Leyden bottle on the table P, with its knob nearly in contact with the ball L. The next morning, if I find this bottle charged, I write the kind of electricity it is charged with againſt the day in the journal, and add, *by the night bottle.*

It is preſumed, that the table is ſufficiently obvious. The two columns for poſitive and negative electricity are uſed only for the firſt obſervation of each day. I uſe FAHRENHEIT's thermometer, ſuſpended on the north outſide of a bow window. The time of making the obſervation with it, and the barometer, and alſo of the direction of the wind, has uſually been about nine o'clock in the morning.

Laſtly, it may be uſeful to obſerve, that I have always found the lower though uninſulated part of the apparatus (*viz.*

the metallic connection of the bell N with the moist earth) to be in a contrary state of electricity to the upper and insulated part, where the pith balls K are suspended. See the 22d of Aug.

Having made a memorandum of the several thunder-storms which have happened in divers parts of this island, according to the information by letters, and from news-papers, I thought it useful to insert them in this journal, in order to shew whether some contemporaneous appearances in my apparatus might not be attributed to them. This seems evidently to have been the case on the 3d of September.

Days.	Wind.	Barom.	Ther.	Sparks.	Pos.	Neg.	May 9, 1789.
		Inches.	°				
May 9	NW	30.	63	0	—	neg.	Balls open about three-tenths of an inch.
10	SW	30.15	60	0	—	neg.	But very weak.
11	E	30.24	61	0	—	neg.	Barely sufficient to separate the balls.
12	E	29.91	60	small	—	neg.	The first day I have had visible sparks; weather cloudy, but fair.
13	E	29.91	61	strong	pos.	—	The weather in the forenoon a little hazy; in the afternoon a thick fog; four o'clock a little rain fell; the rod now became highly electrified positively; the bell rang briskly. I now filled several bottles with the fluid. This strong charge in the rod did not last longer than one hour, but it remained charged positively in a less degree the rest of the day. There was this day some lightning and thunder at Salisbury, and to the west of it.
14	SE	30.	56	0	pos.	—	Balls open four-tenths of an inch.
15	E	29.66	58	small	pos.	—	A.M. and P.M. negative.
16	SE	29.83	61	0	—	neg.	A.M. and P.M.
17	S	29.91	58	0	—	neg.	Balls open six-tenths of an inch.
18	SW	29.91	57	0	—	neg.	Balls open five-tenths of an inch.
19	N	30.20	54	0	—	neg.	P.M. dark heavy weather.
20	SE	30.15	56	small	—	neg.	The rod was electrified nearly the whole day.
21	E	30. 2	54	0	—	neg.	Nearly the same as the preceding day.

Days.	Wind.	Barom.	Ther.	Sparks.	Pol.	Neg.	
		Inches.	°				
May 22	S	29.89	58	0	pof.	—	} Juſt ſufficient to indicate the kind. Serene weather.
23	S	29.90	59	0	pof.	—	
24	SE	29.75	60	0	pof.	—	
25	E	29.68	62	0	—	neg.	Six o'clock A.M. Soon after a fog with drizzling rain, by which the rod became charged poſitively.
26	NE	29.64	61	0	—	neg.	A.M.
27	SW	29.65	60	0	—	neg.	A.M. Balls open three-tenths of an inch.
28	SW	29.80	60	0	pof.	—	But very weak.
29	W	29.72	61	ſtrong	pof.	—	A very cloudy morning, though at too great a height for my rod; but in the afternoon the clouds approached much nearer, and the rod became charged pretty ſtrongly poſitive, which continued about one hour and a quarter. I charged ſome Leyden bottles with the fluid, ſome poſitive, others negative, for there were four gradual changes of the electricity.
30	S	29.70	56	0	pof.	—	Balls open half an inch.
31	SW	29.57	58	ſtrong	pof.	—	Nine o'clock A.M. a heavy ſhower of rain fell, the rod became ſtrongly charged with negative electricity, all the time the ſhower laſted, which was ſhort and ſudden. One hour afterwards, the electricity changed to a ſtrong poſitive, the bell ſuddenly began to ring, and continued to do ſo five minutes; the pith balls then cloſe ſlowly, and open negative, and continued weakly ſo the reſt of the day. At Edinburgh, ſome lightning and thunder this day.
June 1	SW	29.71	53	ſtrong	pof.	—	A cloudy ſhowery morning. The electric charge in rod was moſt beautiful this day. In about ſix hours time I obſerved ſeven changes of the electricity; five of thoſe changes were gradual. The balls opened from 3 to 4 inches, and remained ſo from 15 to 20 minutes each opening, then gradually cloſed; the other two charged ſlowly, but cloſed very quick.

June

Days.	Wind.	Barom.	Ther.	Sparks.	Pos.	Neg.	
		Inches.	°				
June 2	SW	29.85	63	0	pos.	—	But weak.
3	SW	29.92	62	0	pos.	—	A.M.
4	W	29.42	52	ſtrong	pos.	—	A cool cloudy morning. The rod was pretty ſtrongly electrified full ſeven hours to-day, the reſt of the day but weakly ſo. There were 12 changes of the electricity, all gradual, except one, which was inſtantaneous; the balls ſometimes exhibited a waving motion, and ſometimes ſudden jerks. There were ſeveral ſhowers of rain, and two of hail; during the fall of hail, the rod was moſt powerfully electrified, the bell rang very ſtrongly; the effects and appearances were ſo awful, that I kept at a good diſtance from the rod. A great deal of lightning on the eaſt ſide of Kent and Eſſex this day.
5	NW	29.52	53	0	—	neg.	Balls open half an inch.
6	W	29.78	53	ſmall	—	neg.	A.M. and P.M. poſitive; there were ſmall ſparks emitted from the braſs ball I, and the pith balls continued cloſing and opening many times, without any change of kind, for full two hours.
7	NW	29.92	52	0	—	neg.	Balls open from one to ſeven-tenths of an inch.
8	NW	30.18	50	0	—	neg.	
9	NW	30.12	57	0	—	neg.	
10	NW	30.25	56	0	—	—	Weak ſigns, balls not open.
11	NW	30.12	55	0	—	—	
12	E	30.25	56	0	pos.	—	Balls open three-tenths of an inch.
13	E	30.28	54	0	pos.	—	Balls open five-tenths of an inch.
14	NE	30.19	51	0	—	neg.	Very cloudy weather, but at too great a height to affect the rod.
15	E	30. 4	55	0	—	neg.	Balls open full half an inch.
16	E	29.90	60	0	—	neg.	Balls open near one inch.
17	S	29.81	62	ſtrong	pos.	—	Ten o'clock A.M. a ſudden ſhower of rain fell, by which the rod became highly electrified poſitively, and continued to emit ſmall ſparks at the ball L, long after the ſhower was over, without any change of the electricity.

Days.	Winds.	Barom.	Ther.	Sparks.	Pos.	Neg.	
		Inches.	°				
June 18	SE	29.89	60	ſmall	poſ.	—	The weather ſhowery; the rod pretty well electrified.
19	SW	29.88	56	0	—	neg.	Balls open ſix-tenths of an inch.
20	SW	29.65	63	0	—	—	Weak ſigns; balls not open.
21	SW	29.71	59	ſmall	—	neg.	This morning ſeveral heavy clouds paſſed over, coming from the SW, by which the rod became moderately electrified negatively. Some rain fell, which increaſed the electrical charge. But, in the afternoon, the wind and clouds put on a more ominous appearance; near five o'clock, began a ſtorm of wind, rain, lightning, and thunder; but the main weight of the ſtorm did not come near my rod. Its dire effects muſt be eaſt of London: nevertheleſs, I had thoſe uſual beautiful appearances which attend a ſtrong charge in the rod. The ſtorm laſted one full hour, during which time there were five ſucceſſive changes of the electricity, *viz.* four gradual ones, attended with ſudden jerks or ſtarts, which often diminiſhed the divergency of the pith balls from 4 inches to 2 inches or leſs. After thoſe jerks the pith balls recovered their former degree of divergency, ſometimes ſuddenly, and at other times ſlowly. The other change happened inſtantaneouſly, the pith balls collapſing and opening ſo quickly, that the eye could barely ſee their motion. There were two other inſtantaneous cloſings and openings of the balls, without a change of the electricity. Much lightning at Graveſend to-day.
22	SW	29.49	56	ſmall	poſ.	—	The rod was in charge all day.
23	SW	29.54	57	ſmall	poſ.	—	Moſt of the forenoon. Afternoon ſome rain fell, and the rod became electrified negatively, and ended poſitively.
24	SW	29.60	61	0	—	neg.	A.M.

June

Days.	Wind.	Barom.	Ther.	Sparks.	Pos.	Neg.	
		Inches.	°				
June 25	NW	29.64	55	strong	—	neg.	A very thick cloudy morning. The electric charge in the rod has been moderately strong and very fine to-day. There were five gradual changes of the electricity.
26	W	29.80	53	0	—	neg.	A.M.
27	W	29.96	52	strong	pos.	—	Cold showery weather. Except some small intervals, the rod has been strongly electrified all this day. The bell once rang briskly for a few minutes, then suddenly stopped its ringing; the pith balls closed and opened negatively, and continued so till a little after eight o'clock P.M. when they changed to positive. At Liverpool, this day, a tremendous storm of lightning, thunder, hail, and rain.
28	W	29.70	50	strong	—	neg.	A very cold morning and showery, the drops of rain were very large. The rod has been very powerfully electrified near twelve hours this day, during which time there happened eleven changes of the electricity, all gradual but one. The balls often exhibited a waving pendulous motion, without any diminution in their divergency. There were also a few of those jerks before noticed on the 21st instant.
29	W	29.90	51	strong	—	neg.	The bell rang briskly. There was no change of the electricity.
30	NW	30.10	53	0	pos.	—	Nearly all day. Weather serene and clear.
July 1	N	29.19	76	0	—	—	Weak signs, balls not open. An even dark sky, but fair.
2	N	30.3	76	0	—	—	
3	N	30.2	79	0	pos.	—	Weather serene, very hot, and a clear sky.
4	N	30.2	80	small	pos.	—	A.M. and P.M. negative. At Glasgow, some lightning and thunder.
5	NE	29.19	74	0	—	—	Weak signs. Balls not open.
6	NE	29.17	72	small	—	neg.	A heavy dark atmosphere, but fair. At Monkshiver, near Bath, much lightning and thunder.

Days.	Wind.	Barom.	Ther.	Sparks.	Pof.	eg.	
		Inches.	°				
July 7	NE	29.16	69	0	—	—	Weak ſigns of electricity; balls not open.
8	NE	30.	60	0	—	—	
9	NE	29.40	72	0	—	—	
10	NE	29.20	72	0	—	neg.	A.M.
11	E	29.15	74	0	—	neg.	A.M.
12	E	29.13	78	0	—	—	Balls not open.
13	W	29. 5	72	ſmall	—	neg.	A.M. and P.M. poſitive with ſparks. At Hereford, this day, a ſtorm of rain, lightning, and thunder.
14	SW	29. 6	69	0	—	—	Weak ſigns; balls not open.
15	NW	28. 8	52	ſmall	—	neg.	A.M. and P.M. poſitive, with bright ſparks. At Strichen, in Scotland, a heavy thunder ſtorm.
16	W	30.	60	0	—	—	Balls not open. At Edinburgh, this day, ſome lightning and thunder.
17	SW	29.10	65	ſmall	—	neg.	P.M. fine ſparks; no change of kind. At Glaſgow and Hamilton, this day, lightning, thunder, hail and rain; and alſo at Newcaſtle, in Northumberland, the ſtorm was ſevere.
18	SW	29.15	59	0	—	—	Weak ſigns; balls not open.
19	W	29.10	50	0	—	neg.	P.M.
20	NW	29.10	54	ſtrong	—	neg.	Three o'clock P.M. I ſaw a thunder ſtorm approaching. While the ſtorm remained at a conſiderable diſtance, the rod was very highly charged with negative electricity, and continued ſo three quarters of an hour; during which time diſtant thunder was heard. Sometimes the balls were affected with a jerking, at others a waving, motion. The wind now ſhifted to the SE, and a heavy rain ſoon came on; the electricity now changed to poſitive, and the bell now rings briſkly; every appearance (both within and without the room) was tremendouſly awful. I therefore ſeated myſelf upon a large inſulated ſtool, where I could with ſafety obſerve the apparatus. A Leyden bottle was undeſignedly placed with its braſs knob near to the braſs

July

Days.	Wind.	Barom.	Ther.	Sparks.	Poſ.	Neg.	
		Inches.	°				ball L. This bottle charged and ſpontaneouſly diſcharged almoſt as quick as I could notice them; and at the ſame time there was a continual flaſhing of denſe ſparks between the bell and braſs ball L. Theſe very grand appearances only laſted ten minutes, and the ſcene was terminated by a clap of thunder; but the ſtorm, and its effects on the rod, laſted near two hours. I ſaw no lightning but what was in the apparatus. There were nine gradual changes of the electricity, from negative to poſitive, &c.
July 21	SW	29.15	52	ſtrong	—	neg	I had purpoſely placed a large glaſs bowl, upon an inſulated table, in the open air, to catch the falling electrified rain. As ſoon as the bottom of the bowl was well covered with the rain water of the above ſtorm, to my great ſatisfaction, a pair of linen threads I had placed for the purpoſe diverged near two inches; the water remained electrified near ten minutes after it was taken into the houſe. P.M. a little rain fell, and the electricity changed to poſitive. At Edinburgh and Bamff, this day, there was much lightning and thunder.
22	SW	29.10	61	ſtrong	—	neg.	A very ſtrong electricity in the rod all the forenoon. 12 o'clock ſome rain fell, on which the rod became highly charged poſitively, the bell rang weakly for a long time, it then ſtopped, and the balls cloſed, and opened negative, and continued ſo full three hours. This day, near Shrewſbury, a ſtorm of rain, lightning, and thunder.
23	W	29.10	51	ſtrong	poſ.	—	Both A.M. and P.M. and ended negative as the preceding day.
24	W	29.15	71	ſmall	—	neg.	A.M. and P.M. poſitive.
25	W	29.20	71	o	—	—	Weak ſigns; balls not open. An
26	W	29.10	69	o	—	—	univerſal cloudy dark ſky.

Days.	Wind.	Barom.	Ther.	Sparks.	Pos.	Neg.	
		Inches.	°				
July 27	N	29.15	64	ſtrong	—	neg.	Four o'clock P.M. happened a ſudden and ſhort ſtorm of wind and rain. The rod became powerfully electrified, the bell rang for 20 minutes, then ſtopped, and the electricity became poſitive, and ſtrong. At Cambridge, this day, a tremendous ſtorm of lightning and thunder.
28	N	29.20	74	0	pos.	—	A.M. balls open five-tenths of an inch.
29	W	30.	56	ſmall	—	neg.	A.M.
30	NW	29.30	73	0	—	neg.	Balls open ſix-tenths of an inch.
31	W	29.20	75	0	—	neg.	P.M. a regular dark ſky, with ſmall rain, which laſted four hours; ſuch rain is never electrified ſtrongly. Air is very moiſt.
Aug. 1	W	29.79	60	0	—	—	A heavy dark atmoſphere, and a warm ſoft air. The electricity theſe ſix days amounts only to weak ſigns, balls not open.
2	W	30.12	56	0	—	—	
3	SE	30.	62	0	—	—	
4	E	30. 5	61	0	—	—	
5	E	29.79	66	0	—	—	
6	SW	30. 9	60	0	—	—	
7	W	30 27	56	0	—	neg.	P.M.
8	W	30.24	58	0	—	neg.	A.M.
9	E	30.20	66	0	—	neg.	A.M.
10	E	30.17	60	0	—	neg.	A.M.
11	NE	30.20	66	0	—	—	Balls not open.
12	E	30.16	58	ſtrong	pos.	—	P.M. a fine ſhower of rain, on which the rod became pretty well electrified; but ſoon changed to a much more ſtrong negative electricity, which afforded fine ſparks at the braſs ball I. The pith balls many times cloſed and opened, without any more change of kind.
13	E	30.14	60	ſmall	pos.	—	A.M.
14	NE	30.13	62	0	pos.	—	A.M. balls open ſeven-tenths of an inch.
15	NE	30. 9	59	ſtrong	pos.	—	A.M. P.M. a very black cloud paſſed over the rod, by which it became very ſtrongly electrified for a few minutes only; the bell rang briſkly.
16	N	30.15	62	ſtrong	pos.	—	There has been much rain to-day, and the drops very large. The rod has been

Days.	Winds.	Barom.	Ther.	Sparks.	Pos.	Neg.	
		Inches.	°				in high charge great part of the day. The ſparks at the braſs ball L were very pungent; even the air in the room, and the uninſulated bell, and other things, ſhewed that they were electrified. The bell rang briſkly at three very different times. This ſtrong charge of the rod continued full four hours; and the electricity, during that time, changed in kind ten times. Eight of them were gradual, two of them were quick, and attended with jerks. I heard ſome rumbling of thunder at a great diſtance. At Dunwich, in Suffolk, this day, much lightning and thunder.
Aug. 17	NE	30.26	61	ſmall	—	neg.	A.M.
18	NE	30.32	61	o	—	neg.	Balls open nine-tenths of an inch.
19	NE	30.25	60	o	—	—	} Weak ſigns; balls not open.
20	E	29.95	63	o	—	—	
21	SE	29.75	63	ſtrong	—	neg.	P.M. a ſtrong charge in the rod, but of ſhort duration.
22	W	29.75	56	ſtrong	—	neg.	The rod was in high charge ten hours this day, except a few ſmall intervals; there were nine gradual changes of the electricity, from negative to poſitive, and the contrary. The bell rang very briſkly at three different times during that period. There were ſeveral ſhowers of rain, and one of hail; during the latter, the electric charge in the rod was moſt intenſe; the ſparks darted between the ball and bell extremely ſharp and quick. I found the moiſture in the air of the room was now electrified, alſo the bell, and its metallic connexion with the earth, and even the bricks in the wall to which the metal is faſtened were all electrified with an electricity contrary to that in the inſulated part of the apparatus. An electrician (who had often in vain called at my houſe to ſee

Aug.

Days.	Winds.	Barom.	Ther.	Sparks.	Pof.	Neg.	
		Inches.	°				the apparatus in high charge) placed his left hand on the bell, and with his finger of the right hand approached the ball L; a denſe ſpark iſſued to it, and he received a ſmart ſhock in his arms and breaſt, like that of the Leyden bottle. We then joined hands, and made the circuit in the uſual way, between the bell and ball, and we both received a ſevere ſhock. Much lightning and thunder this day at Stirling and Dumfries, in Scotland.
Aug. 23	W	29.94	52	ſtrong	pof.	—	A.M. and P.M. negative on a fall of rain.
24	NW	30.15	58	ſmall	pof.	—	Great part of the day.
25	SW	30.19	60	o	pof.	—	A.M. but very weak in the afternoon.
26	SW	30.10	58	o	—	—	Weak ſigns; balls not open.
27	N	30. 8	59	o	—	—	
28	W	30. 3	54	o	pof.	—	A.M. a ſerene clear ſky.
29	SE	29.90	58	o	—	—	Only weak ſigns.
30	SW	29.80	60	o	—	—	
31	NW	29.75	62	o	—	neg.	P.M. from a ſhower of rain, the drops of which were very large. The electricity ſoon changed to poſitive. Though every circumſtance was favourable for a ſtrong electrification, yet the charge in the rod was but weak. My ſuſpicion led me to try the ſtate of the uppermoſt end of the rod, and I found it to be in a contrary ſtate of electricity to that at the lower end of it; the middle part of the rod was in its natural ſtate, that is, ſhewed no ſigns of being electrified; therefore the rod was only (at this time) inſenſibly electrified.
Sept. 1	SE	29.72	58	o	—	neg.	A.M.
2	S	29.60	66	ſmall	pof.	—	A.M. P.M. a ſhower of rain; the electricity became negative.
3	S	29.50	69	ſmall	pof.	—	In the forenoon. P.M. a ſtrong gale of wind. I now ſaw thunder clouds forming at a great height. Half after

Sept.

Days.	Wind.	Barom.	Ther.	Sparks.	Poſ.	Neg.	
		Inches.	°				five o'clock appearances were dreadful; in five minutes time the ſtrong gale of wind became a ſtorm from SE. This ſtorm of wind (for there was but little rain) carried the huge black clouds to the NW, for there I ſaw abundance of red lightning a little above the horizon, and I once heard a rumbling of thunder. The rod before the ſtorm was poſitive, but by it was changed to negative, and continued ſo during the whole time. The electricity of the rod often varied in ſtrength, ſometimes weak, ſometimes ſtrong, without change of kind. At Amerſham, in Berkſhire, and at the ſame time at the Earl of Aylesford's park, near Packington, in Warwickſhire, was a moſt tremendous ſtorm of hail, rain, lightning, and thunder, from 4 to 6 o'clock P.M.
Sept. 4	SE	29.50	68	ſmall	—	neg.	P.M. There were four gradual changes of the electricity in two hours.
5	SW	30.	69	ſmall	—	neg.	P.M. There has been a little rain to-day.
6	SW	29.95	59	ſmall	poſ.	—	P.M. and near ſun-ſetting negative; there was ſome ſmall rain.
7	SW	30. 3	62	0	poſ.	—	A.M. by the night bottle; that in the rod very weak.
8	SW	30.15	67	0	—	—	Only weak ſigns of electricity; balls not open.
9	SW	30.15	68	0	—	—	
10	SW	30.12	68	0	—	—	
11	N	29.95	61	0	—	—	
12	N	30.10	54	0	poſ.	—	By the night bottle.
13	W	30.10	54	0	—	—	Weak ſigns; balls not open. A dark atmoſphere.
14	SW	29.85	60	0	—	—	
15	W	29.82	48	0	—	—	
16	SW	29.74	48	ſtrong	—	neg.	A.M. There were ſeveral heavy clouds paſſed from the SW, by which the rod was highly charged with poſitive electricity. The bell rang briſkly. This charge laſted from a little before eleven o'clock

Days.	Wind.	Barom.	Ther.	Sparks.	Pof.	Neg.	
		Inches.	°				A.M. to full ſix o'clock P.M. during which time there were four gradual changes of the electricity.
Sept. 17	NW	29.92	44	0	—	neg.	By the night bottle.
18	NW	29.61	52	0	—	neg.	A.M. a more tranſient viſit of the electric fluid I never before ſaw, juſt while a ſmall black cloud paſſed over the rod, and let fall a few drops of rain; the whole time of the charge was about 4 minutes.
19	W	29.40	48	0	—	neg.	A.M. by the night bottle. The electrification of the rod very weak.
20	W	29.40	60	0	—	—	Weak ſigns; balls not open.
21	W	29.40	60	ſmall	—	neg.	A.M. by the night bottle. P.M. the rod was electrified negatively.
22	NW	30. 9	51	0	—	neg.	By the night bottle, and the rod alſo.
23	NW	30. 2	51	0	—	neg.	The electrical charge has been very weak theſe eight days, only juſt ſufficient to indicate the kind.
24	NW	30.23	61	0	—	neg.	
25	NW	30.26	49	0	—	neg.	
26	S	30.26	49	0	—	neg.	
27	SW	30.	61	0	—	neg.	
28	W	30.	51	0	—	neg.	
29	S	29.71	58	0	—	neg.	
30	S	29. 1	48	0	—	neg.	
Oct. 1	SW	29.22	53	ſmall	pof.	—	A.M. by the night bottle, which was ſufficiently charged to give a bright ſpark on making the circuit. At noon ſome rain fell, by which the rod became electrified negatively, and very ſtrong, which laſted full four hours; during that time, the electricity changed four times, which were all gradual.
2	SW	29.28	47	ſtrong	—	neg.	A.M. P.M. the rod was electrified poſitively, and afterwards negatively, with ſtrong ſparks at the braſs ball L.
3	SW	29.11	48	ſtrong	pof.	—	A.M. but much more ſtrongly poſitive in the afternoon; the bell for a ſhort time rang briſkly, after that faintly, then ſtopped, and the electricity changed to negative, which continued ſome hours, then declined gradually to weak ſigns only.

Oct.

Days.	Wind.	Barom.	Ther.	Sparks.	Poſ.	Neg.	
		Inches.	°				
Oct. 4	W	29.40	51	0	poſ.	—	A.M. by the night bottle.
5	S	29.53	45	ſmall	poſ.	—	A.M. And P.M. negative.
6	S	29.50	56	ſmall	poſ.	—	A.M. P.M. the ſame, but much ſtronger. There hath been to-day a cold ſmall rain. The electricity changed four times in two hours. This day, at Whitehaven and Lancaſter, was much lightning and thunder, rain, hail, &c.
7	W	29.27	45	ſmall	poſ.	—	A.M.; but in the afternoon a much ſtronger poſitive charge.
8	E	29.20	48	0	poſ.	—	A.M.
9	SW	29.29	46	0	poſ.	—	A.M.
10	SE	29.50	45	0	poſ.	—	A.M.
11	SE	29.80	49	0	poſ.	—	A.M.
12	S	29.46	53	0	—	—	Weak ſigns; balls not open. Dark, hazy weather.
13	N	29.50	53	0	—	—	
14	W	29.50	53	0	—	—	
15	W	29.52	56	0	—	—	
16	W	29.67	42	ſmall	poſ.	—	A.M. juſt ſufficient to emit viſible ſparks.
17	W	29.90	40	0	poſ.	—	P.M. by means of a fog.
18	S	29.67	52	0	—	neg.	P.M. balls open ſix-tenths of an inch.
19	S	29.79	50	0	—	—	Weak ſigns; balls not open.
20	SW	29.75	58	0	—	—	
21	S	29.89	53	0	poſ.	—	A.M.
22	W	30. 4	50	0	—	—	Weak ſigns of electricity; balls not open. Dark, hazy weather, and moiſt air.
23	NE	30. 7	53	0	—	—	
24	E	30.24	48	0	—	—	
25	NE	30.28	55	0	—	—	
26	E	30.27	47	0	—	—	
27	E	30.27	45	0	—	—	
28	E	30.30	40	0	—	—	
29	N	30.13	46	0	—	—	
30	E	30. 3	40	0	poſ.	—	P.M. by means of a little cold rain.
31	N	29.92	40	0	poſ.	—	P.M. a very high north wind.
Nov. 1	N	30.32	38	0	—	—	Only weak ſigns of the electric fluid; balls not open. A dark, clouded atmoſphere.
2	N	29.59	46	0	—	—	
3	S	29.19	39	0	—	—	
4	SE	29. 3	48	0	—	—	
5	S	29.29	50	0	—	—	
6	W	28.82	48	0	—	—	

Days.	Wind.	Barom.	Ther.	Sparks.	Pof.	Neg.	
		Inches.	°				
Nov. 7	W	28.90	43	ſmall	—	neg.	A.M. by a ſhower of ſleet, which for a ſhort time occaſioned the rod to emit bright ſparks at the ball L.
8	NW	29 9	40	0	—	neg.	A.M. by the night bottle.
9	W	29.29	30	0	—	neg.	
10	W	29 90	37	ſmall	pof.	—	A.M. by means of a fog.
11	W	29 93	39	ſmall	—	neg.	A.M.
12	SW	29.70	45	0	pof.	—	P.M. This was obtained in the following manner. Soon after ſunſet I perceived a light-coloured dewy vapour ariſe 20 or 30 inches above the ground in the park; the evening being ſerene and fair, I ſtood upon an inſulated ſtool, and waved my exploring rod among the dew *, and with my finger touched a ſenſible electrometer, which inſtantly opened with poſitive electricity. As the evening advanced, a ſtrong fog filled the air; when it was of ſufficient height for the high or fixed rod, this alſo became electrified with the ſame kind of electricity which I had received near the earth's ſurface.
13	W	29.73	40	0	—	—	Weak ſigns only; balls not open.
14	SW	29.63	46	0	—	—	
15	SW	29.42	47	0	—	—	Notwithſtanding all my care and attention to my pointed rod, this day it has been intirely fruſtrated; for I have not perceived any ſigns of the electric fluid. However, it is the firſt day it has wholly failed me. A moiſt air

* When I find that the moiſture in the air has ſo far injured the inſulation of my high-pointed rod, that it will not retain the electric fluid; in that caſe, I make uſe of a ſmall rod which I hold in my hand, and project through an upper window; having firſt *warmed* the ſtool legs, I place myſelf upon it, &c. I find this method to be a good ſubſtitute in damp weather. The rod is about the length and ſtrength of a fiſhing-rod, with plenty of ſmall wire twined round it.

Days.	Wind.	Barom.	Ther.	Sparks.	Pos.	Neg.	
		Inches.	°				has prevailed for many days; and there have been six hours drizzling rain to-day, which must lessen the exactness of the insulation of the rod, and air also.
Nov. 16	SW	29.43	43	0	—	—	Weak signs; balls not open.
17	S	29.54	47	small	—	neg.	A.M. small sparks were in the rod, the greatest part of this day.
18	S	29.51	40	0	pos.	—	A.M. by means of a fog.
19	S	29.69	43	0	pos.	—	
20	W	29.65	47	0	pos.	—	
21	NW	29.86	44	0	—	neg.	A.M. by the night bottle, which I found well charged.
22	N	29.89	44	0	—	—	Weak signs only.
23	NE	30.12	42	0	—	—	No signs. This is the second failure. A moist atmosphere.
24	NE	30.29	41	small	pos.	—	A fog. There were visible sparks during the greatest part of the day.
25	N	30.20	37	0	pos.	—	A.M. by means of a fog.
26	N	30.30	37	small	pos.	—	A.M. and P.M. positive, with fine sparks.
27	SE	30.45	50	0	—	neg.	At break of day. Afterwards positive, by means of a fog.
28	W	30.43	33	0	pos.	—	All day, by a continued fog.
29	NW	30.30	35	small	pos.	—	All day. The fog still continues. I have observed, during these three days and three nights (abating a little time for sleep, and which I curtailed for so noble a purpose), that a foggy vapour was constantly electrified positively. The pith balls diverged from a quarter to three quarters of an inch, except when the fog (which was generally moderate) suddenly became thick and dark; then the balls would open to near two inches; at those times I received the electric fluid into bottles. This fog began strongly negative.
30	S	30.10	38	6	pos.	—	Nearly all day. The fog is entirely gone.
Dec. 1	SE	29.71	38	0	—	—	Weak signs; balls not open.
2	S	29.75	50	0	pos.	—	P.M. A moist air to-day.
3	S	29.90	39	strong	pos.	—	All day (I mean 16 hours out of 24), the weather foggy.

Days.	Winds.	Barom.	Ther.	Sparks.	Pos.	Neg.	
		Inches.	°				
Dec. 4	S	30.10	43	o	—	—	Weak signs; balls not open.
5	SW	30.38	52	o	—	—	
6	SW	30.40	48	o	—	—	No signs of electricity. The third failure. A very moist air.
7	SW	30.49	42	o	pos.	—	A.M. a slight fog.
8	SW	30.54	40	small	pos.	—	P.M.
9	SW	30.55	34	o	pos.	—	A.M.
10	W	30.52	39	o	—	—	Weak signs; balls not open.
11	SW	30.54	42	o	pos.	—	A.M. and P.M. The weather serene
12	SW	30.41	44	o	pos.	—	and fair.
13	SW	30.30	47	o	—	—	Weak signs.
14	S	29.92	44	o	—	neg.	A.M. from a little rain.
15	SW	28.95	45	o	—	—	No signs of electricity. The fourth day's failure.
16	SW	29.40	40	strong	pos.	—	P.M. a short shower of snow, by which the rod was strongly electrified.
17	W	29.15	38	o	—	—	No signs of electricity. The fifth day's failure.
18	W	30.	50	small	pos.	—	A.M. a fine serene morning.
19	SW	29.58	44	o	—	neg.	P.M. by a small rain. On an increase of the fall of rain, the electricity changed to positive.
20	SW	29.72	44	o	—	—	Weak signs.
21	SW	29.78	45	o	—	—	No signs of electricity. The sixth day's failure. A very damp air.
22	SW	29.60	56	o	—	—	No signs. The seventh day's failure. A moist air.
23	W	29.78	49	o	—	—	Weak signs of electricity.
24	W	29.27	52	small	—	neg.	At eight o'clock, A.M. began a small shower of rain, which was weakly electrified negatively; as the fall of rain increased, so did the intensity of the electric fluid increase with it, and after continuing for two hours, they both disappeared together.
25	SW	29.41	42	strong	—	neg.	A.M. by a shower of rain. Ten o'clock P.M. a fog strongly electrified positive.
26	W	29.89	46	small	pos.	—	A.M.
27	SW	29.80	46	o	—	—	Weak signs.
28	W	29.88	47	o	pos.	—	P.M. from a very thin fog.
29	SW	29.75	54	o	—	—	
30	SW	29.76	42	o	—	—	Only weak signs of electricity.
31	S	29.73	48	o	—	—	

Jan.

Days.	Wind.	Barom.	Ther.	Sparks.	Pos.	Neg.	
1790		Inches.	°				
Jan. 1	SW	29.90	35	o	pos.	—	Both A.M. and P.M. positive.
2	S	30.30	36	small	pos.	—	All day.
3	S	29.98	46	o	pos.	—	All the day, by a fog.
4	W	30.20	43	small	pos.	—	All the day, and bright sparks several times to-day.
5	N	30.25	45	small	pos.	—	All the day, with fine sparks.
6	SE	30.19	41	o	pos.	—	A.M.
7	W	30.40	41	small	pos.	—	A.M. weather dry and foggy.
8	W	30.45	36	small	pos.	—	Nearly all day in charge.
9	E	30.28	36	o	pos.	—	A.M. balls open near one inch.
10	SE	30.24	39	o	pos.	—	A.M. and P.M.
11	S	30.11	41	o	pos.	—	P.M.
12	W	30.20	45	o	—	—	Weak signs.
13	SW	30.34	42	o	pos.	—	A.M.
14	S	29.95	49	small	—	neg.	A.M. a small rain, by which the rod was electrified negatively.
15	NW	29.91	43	o	pos	—	A.M.
16	SW	29.95	42	small	pos.	—	A.M. weather very mild and fair.
17	N	30.28	40	o	pos.	—	A M.
18	NE	30.20	39	o	pos.	—	A.M. and P.M.
19	E	30.13	33	o	pos.	—	A.M. and P.M.
20	SE	30.24	32	small	pos.	—	A.M.
21	E	30.40	31	small	pos.	—	A.M. and P.M.
22	W	30.38	35	small	pos.	—	All the day, from a fog. The strength of the electric charge in the rod was much governed by the occasional intensity of the fog; for as one abated in strength so did the other, and the contrary. At ten o'clock P.M. I received bright sparks at the ball L.
23	W	30.33	40	small	pos.	—	Seven o'clock A.M. I found the rod electrified, sufficiently strong to emit visible sparks, and often afterwards the same day.
24	W	30.16	46	small	pos.	—	A.M.
25	NE	30.11	42	small	—	neg.	There was a moderate shower of rain this morning, by which the rod was electrified negatively.
26	W	30. 2	40	o	pos.	—	A.M. balls open from two to seven tenths of an inch.
27	W	29.30	42	o	pos.	—	
28	S	29.50	44	small	pos.	—	A.M. at noon fell a small rain, which was void of electricity. Near ten

Days.	Winds	Barom.	Ther.	Sparks.	Pof.	Neg.	
		Inches.	°				ten o'clock P.M. a ſudden ſhower of rain, ſtrongly negative
Jan. 29	W	29.26	40	0	pof.	—	P.M. A delightful clear fine day, but the atmoſpheric electricity was very weak till night came on.
30	N	29.50	41	0	pof.	—	A.M. And P.M. negative.
31	S	29.68	46	0	pof.	—	
Feb. 1	N	29.87	40	0	pof.	—	
2	SW	30.10	40	0	pof.	—	
3	W	30.30	44	0	pof.	—	Balls open from one to nine tenths of an inch, weather fair and ſerene.
4	W	30 60	46	0	pof.	—	
5	SW	30.61	43	0	pof.	—	
6	W	30.63	44	0	pof.	—	
7	NW	30.35	39	0	pof.	—	
8	N	30.22	36	0	pof.	—	
9	W	30. 5	40	ſmall	pof.	—	Sparks juſt viſible.
10	NE	30. 7	43	0	pof.	—	
11	W	30.20	41	0	pof.	—	Serene weather ſtill continues.
12	W	30.20	45	0	pof.	—	
13	SW	30.34	42	ſmall	pof.	—	Nearly all the day, with fine bright ſparks.
14	SW	30.15	44	0	pof.	—	Balls open from two to ſix tenths of an inch.
15	W	30.22	40	0	pof.	—	
16	S	29 95	44	0	pof.	—	A.M. A ſmall rain P.M. which did not occaſion any change in the electric fluid.
17	W	30.20	39	0	pof.	—	Balls open half an inch, weather ſtill ſerene.
18	W	30.42	42	0	pof.	—	
19	S	30.48	45	0	pof.	—	
20	SE	30.42	40	ſmall	pof.	—	A.M. A foggy day.
21	SW	30.38	42	0	pof.	—	
22	SW	30.43	43	0	pof.	—	Balls open from one to nine tenths of an inch, weather ſtill mild.
23	S	29.97	46	0	pof.	—	
24	SW	30.15	44	0	pof.	—	
25	S	29 99	51	0	pof.	—	
26	W	29.88	51	0	pof.	—	P.M. I have often obſerved that impetuous winds leſſen the intenſity of atmoſpheric electricity in clear weather, which has been verified this day; for not the leaſt ſign of electricity could be obtained from the rod till after ſunſet, at which time the high weſt wind hav-

Day.	Winds.	Barom.	Ther.	Sparks.	Pos.	Neg.	
		Inches.	°				ing subsided, a little low vapour sprung up; I then received the fluid in great plenty, and of the same kind that it has continued to be for twenty-seven days past.
Feb. 27	W	30.20	42	0	—	—	Weak signs: balls not open.
28	NW	30.22	45	0	—	—	
Mar. 1	N	30.32	47	0	pos.	—	A.M.
2	NW	30.49	50	small	pos.	—	A.M. sparks just visible.
3	N	30.33	49	0	pos.	—	Very mild serene weather.
4	NW	30.27	40	0	pos.	—	
5	NE	30.31	43	0	pos.	—	
6	NW	30.47	40	0	pos.	—	
7	NW	30.45	42	0	pos.	—	
8	SW	30.44	42	small	pos.	—	A.M. visible sparks. Still no change of kind.
9	SW	30.20	41	0	pos.	—	A.M.
10	SW	29.90	46	0	pos.	—	P.M. A strong gale of wind to-day. The atmosphere is extremely dry. The sun appeared bright all day. No atmospheric electricity could be obtained till near ten o'clock at night. This day, at Hallifax, fell a shower of snow, accompanied with one flash of lightning and one clap of thunder.
11	SW	30.20	41	0	pos.	—	A.M. Still moderate weather.
12	S	30.31	51	0	pos.	—	
13	SW	30.20	41	0	pos.	—	A.M. Nine o'clock PM. to eleven o'clock there was a moderate shower of rain, which was electrified negatively. I have not till this perceived a negative charge since January the 28th.
14	NW	30.50	49	0	pos.	—	A.M. At Thurso, in Scotland, hail, lightning, and thunder.
15	N	30.57	51	0	pos.	—	P.M.
16	N	30.60	40	small	pos.	—	A.M. visible sparks.
17	NE	30.60	40	0	pos.	—	A.M. a very dry atmosphere.
18	NE	30.50	45	0	pos.	—	
19	E	30.55	44	0	pos.	—	
20	E	30.44	43	0	pos.	—	A.M. there was an hoar frost upon the grass this morning.
21	E	30.39	44	0	pos.	—	A.M.

Days.	Wind.	Barom.	Ther	Sparks.	Pof.	Neg.	
		Inches.	°				
Mar. 22	E	30.15	42	0	pof.	—	
23	SW	29.81	46	0	pof.	—	
24	S	29.80	51	0	pof.	—	A.M. divergency of the pith balls from one to nine tenths of an inch.
25	NW	30.	46	0	pof.	—	
26	E	30. 4	48	0	pof.	—	
27	E	30.	46	0	—	neg.	A.M. from a ſmall ſhower of rain. P.M. the fluid was poſitive.
28	NE	29.98	44	0	pof.	—	P.M.
29	N	30.	48	0	pof.	—	P.M. the air a little foggy.
30	E	29.95	44	0	pof.	—	A.M. not a beam of ſun has appeared this day.
31	E	29.93	47	0	pof.	—	A.M. balls half an inch open.
April 1	E	30. 5	45	0	pof.	—	A.M.
2	E	30.20	41	ſmall	pof.	—	A.M. viſible ſparks.
3	E	30.25	40	0	pof.	—	
4	E	30.19	40	0	pof.	—	Balls open from one to ſeven tenths of an inch.
5	NE	30.23	47	0	pof.	—	
6	E	29.95	36	0	pof.	—	
7	E	29.75	45	0	—	—	Weak ſigns of electricity. This weak ſtate has not happened ſince the 31ſt of December. There has been for many days a ſtrong eaſt dry wind, which ſeems hitherto nearly void of the electric fluid.
8	E	29.75	52	0	—	—	
9	E	29.57	66	ſmall	—	neg.	P.M. from a fine ſhower of rain.
10	E	29.50	40	ſmall	—	neg.	A.M. the rain continues, ſo does its negative electricity.
11	E	29.36	35	ſmall	pof.	—	A.M. from a little fall of ſnow. P.M. ſome ſnow mixed with rain, on which the rod became charged much more ſtrongly poſitive. The rod has been charged full four hours to-day.
12	E	29.35	39	0	—	—	Weak ſigns; balls not open.
13	E	29.70	38	ſmall	—	neg.	A.M. a moderate rain, but ſtrongly electrified, and continued ſo full two hours. There were two gradual changes of electricity.
14	NE	29.81	41	0	—	neg.	P.M. but after ſunſet the rod was electrified poſitively.
15	E	29.68	35	ſmall	—	neg.	Six o'clock A.M. a little rain fell. Half after eight o'clock, a fine ſhower of

April

Days.	Wind.	Barom.	Ther.	Sparks.	Pos.	Neg.	
		Inches.	°				snow; the rod now became strongly electrified positively. Dense sparks were now received at the ball L; half after nine o'clock, the electricity changed to negative. I caught some of the snow in the apparatus mentioned the 20th of July in this journal, and I found it weakly electrified.
April 16	NE	29.85	41	0	pos.	—	
17	N	29.98	42	0	pos.	—	
18	NE	29.80	40	0	pos.	—	Divergency of the pith balls was from a tenth to seven-tenths of an inch. Fine serene weather.
19	NE	29.69	41	0	pos.	—	
20	SE	30.26	45	0	pos.	—	
21	S	30.23	47	0	pos.	—	
22	S	29.94	51	0	pos.	—	
23	SW	29.70	55	0	pos.	—	
24	W	29.55	52	0	pos.	—	A M. and P.M. negative from a little fall of rain.
25	SW	29.60	51	strong	pos.	—	Half past nine o'clock A.M. A distinct black cloud approached the rod, and some heavy drops of rain fell; the electricity of the rod then changed to negative. About half past three o'clock P.M. a very large low cloud passed over the rod, and rained a little, on which the rod became strongly electrified positively.
26	NE	29.85	53	small	pos.	—	A.M. P.M. on the fall of some rain, the rod was charged negatively.
27	N	29.98	47	0	pos.	—	The divergency of the balls from two to six tenths of an inch. Serene fine weather.
28	W	29.80	50	0	pos.	—	
29		29.69	51	0	pos.	—	
30		29.58	53	0	pos.	—	
May 1	E	29.75	50	small	pos.	—	Nine o'clock A.M. a shower of rain; the electricity now became negative. The rod has been electrified to-day from six o'clock A.M. to ten o'clock P.M. and I suppose all night also.
2	N	30.22	41	0	pos.	—	Divergency of the pith balls from three to seven tenths of an inch.
3	NE	29.85	49	0	pos.	—	
4	SE	29.63	51	small	pos.	—	A.M.

Days.	Wind.	Barom.	Ther.	Sparks.	Pof.	Neg	
		Inches.	°				
May 5	S	29.73	52	ſtrong	pof.	—	Six o'clock, A.M. very cloudy. Eight o'clock it rained, the electricity now became ſtrongly negative, with fine ſparks at the ball L. This ſhower having ceaſed, another ſoon followed, which electrified the rod poſitively. The rod was charged 7 hours to-day.
6	SW	29.75	50	ſmall	—	neg.	Every appearance at the rod to-day was nearly as during the preceding one.
7	W	30.10	56	0	pof.	—	Balls open from one to five tenths of an inch. Such weak ſigns of electricity, as have been obſerved for theſe two days, are the uſual effects of a very ſtrong and dry weſterly wind; and in general, let a ſtrong dry wind blow from what point of the compaſs it may, it is attended with weak ſigns of electricity.
8	W	29.84	50	0	pof.	—	

The above-mentioned eighth day of May completes this journal of one whole year, which I give to the curious in atmoſpheric electricity as a faithful narrative of facts, having never once deputed another perſon to make obſervations for me.

JOHN READ,
of Knightſbridge, near London.

A monthly

A monthly account of electrical ſparks, and of poſitive and negative electricity, as indicated by the pith-ball electrometer, and ſometimes by only flaxen threads without balls to them.

		Times.		Times.	Number of days in each month in which ſparks were perceived. Days.
23 days of May, 1789, 8 days of May, 1790,	Poſitive	17	Negative	18	9
June	Poſitive	32	Negative	36	12
July	Poſitive	13	Negative	22	12
Auguſt	Poſitive	19	Negative	19	9
September	Poſitive	9	Negative	23	7
October	Poſitive	17	Negative	7	7
November	Poſitive	12	Negative	8	8
December	Poſitive	12	Negative	6	7
January	Poſitive	26	Negative	4	13
February	Poſitive	26	Negative	0	3
March	Poſitive	30	Negative	1	3
April	Poſitive	28	Negative	12	8
		241		156	98

It appears from this journal, that there were only *ſeven* days throughout the year in which no ſigns of electricity were perceived; *viz.* the 15th and 23d of November, and the 6th, 15th, 17th, 21ſt, and 22d of December.

Remarks on the phænomena exhibited by the rod on the 31ſt of Auguſt.

I was for a long time extremely puzzled to account for the rapid changes which the pith balls on ſome days ſo frequently

exhibited; being poſitive one minute, then negative for another, and the next returning again to poſitive. From often conſidering this apparently whimſical changeableneſs in nature, I was at length induced to ſuſpect, what indeed was afterwards confirmed by actual experiment, *viz.* that ſome of theſe changes are only apparent, and not real, they being occaſioned not by the actual communication of a different ſort of electricity, but merely by the action of electrical atmoſpheres; thus, when an electrified cloud comes within a certain diſtance of the rod, and before it comes near enough to impart to it ſome of its own electricity, the electrical atmoſphere of the former, agreeable to the well known laws of electricity, will diſturb the electric fluid naturally belonging to the rod, and will conſequently occaſion ſeveral apparent changes in the electrometer, which changes an unexperienced obſerver would attribute intirely to the change of electricity in the clouds.

This obſervation was evidently confirmed by the phænomena obſerved on the 31ſt of Auguſt; and thence it appears, that the real number of changes from poſitive to negative, or from negative to poſitive electricity, cannot be ſo great as it is ſhewn by the electrometer affixed to the rod.

I cannot help lamenting with Signor BECCARIA, that there are ſo few high pointed rods erected to aſcertain the electrical ſtate of the earth and atmoſphere at all times; but more particularly during thunder ſtorms. If there had been pointed rods, for inſtance, at Whitehaven and Lancaſter on the 6th of October, and well attended to at the time of the ſtorm of lightning and thunder, which happened at both places nearly at the ſame time, it would then have been known, whether the apparatus might not be poſitive at one place when it is negative at the other.



Fig. 1.

XIII. *Farther Experiments relating to the Decompoſition of dephlogiſticated and inflammable Air.* By Joſeph Prieſtley, *LL.D. F. R. S.*

Read April 7, 1791.

THE doctrine of *phlogiſton*, and that of the *decompoſition of water*, have long engaged the attention of philoſophical chemiſts, and experiments have ſometimes ſeemed to favour one concluſion, and ſometimes an oppoſite one. I have myſelf been very differently inclined at different times, as appears in my publications on the ſubject; and I am hardly ſenſible of a wiſh which way this important controverſy, as it may be called, be decided, notwithſtanding the part that I have taken in it. I cannot help thinking, however, that the experiments, an account of which I ſhall now lay before the Society, are deciſive in favour of the compoſition of an acid from dephlogiſticated and inflammable air; and, therefore, that the opinion of theſe two kinds of air neceſſarily compoſing *water* cannot be well founded. It is, indeed, ſufficiently evident, that the ſame elements likewiſe compoſe *fixed air*, and therefore it is the leſs extraordinary that they ſhould compoſe another acid.

The doctrine of phlogiſton I would, however, obſerve, will not be affected by the moſt deciſive proof of the compoſition of water from dephlogiſticated and inflammable air; ſince this would

would only prove, that phlogiſton is one conſtituent part of water; which is an opinion that I have advanced, and mentioned on ſeveral occaſions; and it is the leſs extraordinary, as water reſembles metals in the remarkable property of being a pretty good conductor of electricity. What I ſhall now allege, however, will make it very doubtful, whether pure water be ever formed by the union of dephlogiſticated and inflammable air; and perhaps make it more probable, that water, as I have lately advanced, is only the *baſis* of thoſe kinds of air, as well as of every other kind.

It was objected to my former experiments on the decompoſition of dephlogiſticated and inflammable air, by firing them together in a copper veſſel, which always produced an acid liquor, that this acid came from the *phlogiſticated air* with which the dephlogiſticated air that I made uſe of was neceſſarily more or leſs diluted; or from that which I could not wholly exclude, as a part of atmoſpherical air, when I exhauſted the copper veſſel by means of an air-pump.

To obviate this objection, I then obſerved, that I not only conſtantly found that the more phlogiſticated air was contained in the two other kinds of air (mixed in the proportion of two meaſures of inflammable air to one of dephlogiſticated) the leſs acid I got; but that, when I purpoſely mixed any given quantity of phlogiſticated air with them, it appeared not to have been at all affected by the proceſs, but remained the very ſame, in quantity and quality, as before. Still, however, becauſe Mr. CAVENDISH, though in a very different proceſs, had found nitrous acid to reſult from the decompoſition of phlogiſticated and dephlogiſticated air; and becauſe M. LAVOISIER and his friends had found nothing but pure water after the ſlow burning of dephlogiſticated and inflammable air; it was maintained

maintained by the favourers of their ſyſtem, that the *water* only in the liquor which I procured came from the union of the two kinds of air, and the *acid* from the *phlogiſticated air* which I had not been able to exclude.

But let any perſon only conſider the very ſmall quantity of nitrous acid which was procured by Mr. CAVENDISH from the certain decompoſition of 3194 grain meaſures of atmoſpherical air, amounting to more than 1¼ ounce meaſures in one caſe, and of 2710 grain meaſures, amounting to 5½ ounce meaſures in another caſe (Phil. Tranſ. Vol. LXXVIII. p. 264. 268.), three-fourths of which was phlogiſticated air; and the vaſtly greater quantity which I procured (Ibid. p. 324.), when it could not be proved, that a particle of phlogiſticated air was decompoſed, and think whether it was at all probable, that the acid came from this kind of air, and not from the union of the dephlogiſticated and inflammable air, which evidently diſappeared in very great quantities. This circumſtance alone might have ſatisfied thoſe who intereſt themſelves in this queſtion; but it does not ſeem to have been attended to.

I have now, however, effectually removed the objection above mentioned, by intirely excluding all phlogiſticated air from the proceſs; the dephlogiſticated air which I at preſent uſe being ſo pure, that it contains no ſenſible quantity of phlogiſticated air. I alſo make uſe of no air-pump, but firſt fill the copper veſſel with water, and then diſplace it by the mixture of the two kinds of air; yet, in theſe circumſtances, in which all phlogiſticated air is excluded, I procure even a ſtronger acid than before.

The paper that I ſend along with this article contains the dry reſiduum of the turbid green liquor, produced by a ſingle exploſion of a mixture of two parts inflammable and ſomething

thing more than one part of dephlogiſticated air, in a copper veſſel which holds thirty-ſeven ounces of water; and a little more muſt have have remained in the veſſel, which I could not get out by draining or ſhaking it. It is moſt evident, therefore, that the acid neceſſary to diſſolve ſo much copper muſt have come from the union of the dephlogiſticated and inflammable air, becauſe there was nothing elſe in the veſſel. The inflammable air was procured from iron by means of ſteam.

This very pure dephlogiſticated air I firſt imagined could only be got by the proceſs in which I obſerved (Experiments on Air, Vol. II. p. 170.) that I once before procured it, though I then ſuppoſed the extraordinary reſult to be accidental; becauſe in other circumſtances I have ſometimes had it very pure when I could not ſucceed in a ſecond attempt of the ſame kind. It was by heating the yellow product of the ſolution of mercury in ſpirit of nitre, without ſuffering the red precipitate into which it is converted by heat to come into contact with the external air, from which I thought it probable that it might attract ſome phlogiſton. Afterwards, however, I found that this circumſtance makes no difference whatever; and that the air ſo procured *appeared* to be purer, aroſe from the greater purity of the *nitrous air* which I made uſe of as a teſt, and which I got from mercury, and not from copper, the nitrous air from which I find to be much leſs pure. For trying the dephlogiſticated air yielded by ſome red precipitate which had been prepared many months by the nitrous air from mercury, it appeared to be as pure as that which was procured in the manner above deſcribed.

That the dephlogiſticated air which I now made uſe of was ſufficiently pure for my purpoſe, appeared from mixing one meaſure of it with two of nitrous air, when the whole quan-

tity was reduced to leſs than four hundredth parts of one meaſure; ſo that it is probable that, by a more accurate proportion of the two kinds of air, and greater addreſs in mixing them, they might have almoſt intirely diſappeared. There is beſides ſome reaſon to think, from the great variety in nitrous air, that the greater part of this very ſmall reſiduum comes from the nitrous air, and not from the dephlogiſticated.

It will be ſaid, how is it poſſible to reconcile the reſult of this experiment with that of M. Lavoisier and his friends? which I was by no means diſpoſed to queſtion after the publication of the *Extract from the Regiſter of the Academy of Sciences for Auguſt* 28, 1790, in the ſeventh volume of the *Annales de Chimie*, in which a diſtinct account is given of a large quantity of very pure water procured from the ſlow combuſtion of the two kinds of air above mentioned: for before this it was acknowledged, that ſome little acid was always found in the water ſo procured.

But my late experiments, beſides aſcertaining the fact of the production of nitrous acid from the decompoſition of dephlogiſticated and inflammable air, throw ſome farther light on the ſubject, and may in ſome meaſure explain their reſult; for I am now able to procure, in my own proceſs, either nitrous acid or pure water, from the ſame materials.

I conſtantly obſerve, that if there be a ſurplus of dephlogiſticated air, the reſult of the exploſion is always the acid liquor; but that if there be a ſurplus of inflammable air, the reſult is ſimply water. That phlogiſticated air is not in all caſes affected by this proceſs, I completely aſcertained, by admitting a little common air into that mixture of the two kinds of air which always produced water, and finding nothing but water in the reſult.

I find, however, that, agreeably to the experiments of Mr. Cavendish, phlogisticated air is decomposed in this process, when there is not enough of inflammable air to saturate the dephlogisticated air; though when there is a redundancy of inflammable air, there is even a production of phlogisticated air. Putting 0.5 oz. m. of phlogisticated air to a mixture of two ounce measures of inflammable air and 1.5 oz. m. of dephlogisticated air, the whole was reduced by explosion to 1.05 oz. m. of the standard of 1.1, with two measures of dephlogisticated air, which appears by computation to contain no more than 0.388 oz. m. of phlogisticated air; so that 0.112 oz. m. had been decomposed in the process. When there is a sufficient quantity of inflammable air, the phlogisticated air always remains unaffected in this process, as appears by mixing any quantity of it with the two kinds of air to be exploded, and finding the very same quantity, as I have repeatedly done, in the residuum.

That when there was a sufficiency of inflammable air for the purpose, phlogisticated air is even *produced* in this process, was evident from my never being able to diminish any quantity of dephlogisticated air by inflammable air so far as by good nitrous air, and the residuum always containing phlogisticated air. Having exploded two measures of inflammable air with one of dephlogisticated air, which by a mixture of two measures of nitrous air was reduced to 0.04, there was a residuum of 0.1, of the standard of 1.3, which appears by computation to contain 0.0767 oz. m. of phlogisticated air.

The reason why, in my former experiments, I always procured more or less acid, must have been that, without any intention, or suspecting that any thing depended upon it, I must have had some surplus of dephlogisticated air. M. Lavoisier I also perceive to have taken it for granted, as I did, that after

after either of our proceſſes, any ſurplus of either of the two kinds of air would only have remained unſaturated, and have been found unchanged in the reſiduum.

I claim no merit whatever in this obſervation. It was in conſequence of accidentally finding pure water in what I then imagined to be the ſame circumſtances in which I had always before found acid, and which ſurprized me not a little at the time, that I was led to vary the proportions of the two kinds of air, till at length I ſucceeded in aſcertaining the circumſtances on which this remarkable difference in the reſult depends; but I am by no means able to aſſign any reaſon for this difference.

In this ſtate of my experiments I concluded, that *nitrous acid*, though conſiſting of the ſame elements with pure water, contains a greater proportion of dephlogiſticated air; and in the laſt edition of my *Obſervations on Air*, Vol. III. p. 543. I obſerved, that "ſubſtances, poſſeſſed of very different properties, may be compoſed of the ſame elements, in different proportions, and different modes of combination. It cannot "therefore be ſaid to be abſolutely impoſſible, but that water "may be compoſed of theſe elements," *viz.* dephlogiſticated and inflammable air.

When I firſt prepared an account of my late experiments for the Royal Society, I entertained this idea; but I now conſider it as at leaſt uncertain, becauſe when I mix the two kinds of air in ſuch proportions as to produce *water*, I find in the reſiduum much more *phlogiſticated air* than I do when *acid* is produced, which affords a ſuſpicion that, in this caſe, *the principle of acidity* goes wholly into the phlogiſticated air, which, as my former experiments ſhew, actually contains it, though it is not eaſy to aſcertain in what proportion.

Having exploded three ounce measures of a mixture of something more than two parts inflammable air, and one of dephlogisticated, and another equal quantity in which the inflammable air bore a less proportion to the dephlogisticated, the former of which I knew would yield water, and the latter acid, I found the residuum of the former to be 0.57 oz. m. not affected by nitrous air, and weakly inflammable; and in order to find how much phlogisticated air it contained, I mixed different proportions of phlogisticated and inflammable air, and concluded, from the manner of firing *them*, and this *residuum*, that it could not consist of less than one-third of phlogisticated air, *viz.* 0.19 oz. m. But the residuum of the mixture which would have produced acid was 0.62 oz. m. of the standard of 1.0, which I find by computation to contain not more than 0.062 oz. m. of phlogisticated air. I repeated this experiment very many times, and never failed to have a similar result; so that it is very possible that the *pure water* we find may be nothing more than the basis of the two kinds of air; and the principle of acidity in the dephlogisticated air, and the phlogiston in the inflammable air, may combine to form a superfluous acid in the one case, and the phlogisticated air in the other.

This supposition is strengthened by finding that whether the produce be acid, or pure water, the two kinds of air unite in nearly the same proportions. But since water has an affinity to almost every substance in nature, and a peculiarly strong one to the acid and alkaline principles, it may be impossible that it should be wholly free from them; and if they be in proper proportions to saturate one another, and in the same quantities, their presence may never appear.

As the reason why, in my former experiments, I always produced an acid liquor, and never pure water, was my using

too

too great a proportion of dephlogiſticated air; ſo the reaſon why M. LAVOISIER and his friends generally produced but little acid, and at laſt not at all, muſt have been, that the *ſlow combuſtion* which they made uſe of gave the principle of acidity in the dephlogiſticated air, and the phlogiſton in the inflammable air, a better opportunity of eſcaping, and forming the phlogiſticated air in their reſiduum, of which they have not publiſhed any ſatisfactory account *; and it is probable, that the weight of theſe elements compared with that of the water which forms the baſis of the two kinds of air, may be very ſmall. That excellent philoſopher M. DE LUC ſuppoſes that they have even no weight at all.

M. LAVOISIER himſelf, I obſerve, lays particular ſtreſs, (p. 262.) on the *ſlowneſs* of the combuſtion, as if he ſuſpected it to be neceſſary to his reſult. This circumſtance may alſo account for my want of ſucceſs in the attempts that I made to repeat his experiment: for whenever I made a ſtream of inflammable air to burn in a veſſel of dephlogiſticated air (which I contrived to do by means of a leſs expenſive, but I own a leſs accurate, apparatus than his) I always got ſome acid, though leſs than in my own proceſs; but I made a larger and ſtronger flame than I imagine M. LAVOISIER choſe to produce.

In the courſe of theſe experiments, I found, that when the inflammable air I made uſe of was from turnings of *caſt iron*, there was always a conſiderable quantity of *fixed air* in the reſiduum, not leſs than one-tenth of a meaſure, after the exploſion of two

* Since this was written, Meſſ. FOURCROY, VAUQUELIN, and SEGUIN, have publiſhed a very particular account of their experiment; from which it appears, that, after the combuſtion of the two kinds of air, there was a pretty large reſiduum of phlogiſticated air, more than was contained in the airs before combuſtion. See *Annales de Chimie*, for April 1791, p. 35.

meaſures

meaſures of inflammable air and one of dephlogiſticated; whereas there was either no fixed air at all, or the ſlighteſt appearance of it imaginable, when I made uſe of inflammable air from *malleable iron*, extracted either by means of ſteam or acids.

The principal of theſe experiments, as well as thoſe in my former Papers on this ſubject, will be found to confirm the ſimilar ones of Mr. CAVENDISH; but they prove the ſource of the *acid* in the reſults not to be what he imagined, *viz. phlogiſticated air*, but the union of the dephlogiſticated and inflammable air; and they alſo make it at leaſt doubtful, whether theſe two kinds of air compoſe *pure water*.

XIV. *Experiments on Human Calculi. In a Letter from Mr.* Timothy Lane, *F. R. S. to* William Pitcairn, *M. D. F. R. S.*

Read May 5, 1791.

SIR,

IT will give me much ſatisfaction ſhould the following experiments, made twenty years ago, ſo far meet your approbation, as to be thought worthy the notice of the Royal Society.

The Lixivium Saponarium of the late Pharmacopœia, prepared with the addition of ſo much lime as nearly to free the ſalt of tartar of its fixed air *, having been uſed as a medicine for the ſtone and gravel ſome years before, and its effects found very unequal, I thought it neceſſary to examine different calculi, then collected, both as to the effect of the above lixivium, and of fire, upon them.

Great diſparity was obſerved; ſome being diſſolved, and others ſcarcely altered in their figure.

When tried by fire, ſome were nearly evaporated by a red heat, and others retained their form.

Different parts even of the ſame calculus, varied conſiderably.

* See Letter to Dr. HEBERDEN, Medical Tranſactions, Vol. I. p. 112.

That I might be better informed of the above, the experiments were repeated both by fire and lixivium, with greater accuracy, as follows.

Fourteen ſpecimens were ſelected, ſome of which were parts of the ſame calculus, and others different calculi.

In the experiments by fire I was favoured with the aſſiſtance of Mr. STANESBY ALCHORNE, of the Tower, to whom were ſent ten grains of each, in ſeparate papers, which were numbered.

The contents of each paper were placed in ſeparate cupels, under a muffle, the ſame as is uſed by him for aſſaying gold and ſilver. The fire was raiſed gradually, till the furnace was fully heated: the time from raiſing the fire to the taking them out again was three hours, when it was concluded, that whatever volatile matter they contained was expelled.

The ſame quantity as above, of each ſpecimen, being put into ſeparate numbered phials, with one ounce meaſure of the lixivium in each, continued forty-eight hours; the phials were frequently ſhaken to forward the ſolution.

The clear liquor of each phial was decanted into freſh phials, and a quarter of an ounce more lixivium was added to ſuch as were undiſſolved; after twenty-four hours they were poured out of the phials into ſeparate filtering papers, each numbered, and the phials waſhed with diſtilled water, which was alſo poured into the papers, ſo that all that remained undiſſolved might be detained by the papers, which with their contents were carefully dried.

The

The remains of each

	Unsublimed.	Undissolved.
	Grains.	Grains.
N° 1.	1¼	¼
2.	2¼	2
3.	¼	¼
4.	1¼	2
5.	¼	0
6.	3¼	2¼
7.	3¼	6
8.	6	8¼
9.	6¼	6¼
10.	6¼	7¼
11.	¼	¼
12.	¼	0
13.	5¼	4
14.	6	5¼

Appearances of each after Calcination.

N° 1. 3. 7. 8. left a fine white and soft powder.

N° 4. 5. 11. 12. left a white and gritty powder.

N° 2. 6. 9. 10. 14. were partly in powder white and gritty, with some lumps of a dark colour, as if not fully calcined.

N° 13. Of this the figure was not greatly altered; it remained hard, and part of it appeared as if inclined to flux.

After being in the lixivium forty-eight hours.

N° 8. 9. 13. 14. were found soft.

N° 7. and 10. remained hard.

Theſe ſix were ſeparately taken out of the lixivium and put into a mortar, and rubbed or broken, and then carefully returned to their ſeparate phials before the ſecond addition of lixivium, in order to forward the ſolution.

Specimens deſcribed.

N° 1. The external part of a laminated calculus, of a light yellowiſh brown colour *.

N° 2. The external part of a calculus, in colour like dirty tobacco-pipe clay †.

N° 3. A light-brown laminated calculus.

N° 4. and 5. Two ſpecimens from one calculus; of which N° 4. is the external coat, of a dirty tobacco-pipe-clay colour.

N° 5. The internal part of N° 4., yellowiſh like N° 1.

N° 6. A calculus taken out of the urethra; a greyiſh white, inclining to yellow, of a porous texture.

N° 7. A calculus about the ſize of a nutmeg, taken from a child of a year old, given me by the late Mr. POTT; aſh-coloured, in waves of different ſhades, laminated and hard.

N° 8. A dark-brown very hard calculus, of the mulberry kind.

N° 9. and 10. Two ſpecimens from one calculus; of which N° 9. is the external whitiſh part, which appeared like a coat of calcareous earth, covering an irregular mulberry calculus ‡.

* The nucleus, ſo called, being the central part, was of a much deeper colour, and had been found not ſo ſoluble in lixivium as the light-brown part.

† The nucleus was of a bright yellow, and more ſoluble in lixivium than the whitiſh part.

‡ The covering of this calculus induced me to ſuſpect that lime or lime-water might have been taken, and, by being decompounded by freſh urine, containing fixed air, form this covering. Other calculi have afforded the ſame ſuſpicions.

In future, an account of medicines taken might afford much information, joined with the examination of different parts of large calculi taken out of the bladder.

N° 10.

N° 10. The brown mulberry part covered by N° 9. The three following are parts of one large, laminated calculus; of which

N° 11. Is the external lamina, of a browniſh yellow.

N° 12. The central part, called the nucleus, of a pale orange colour.

N° 13. Some of the laminæ, between the nucleus and the external coat, of a ſparkling appearance.

N° 14. A whitiſh, porous, and eaſily broken calculus.

The experiments by fire explain the unequal accounts of authors, reſpecting the component parts of calculi.

In general, thoſe which contain the largeſt proportion of volatile parts were moſt ſoluble in lixivium.

The inſolubility of ſome explains the want of ſucceſs in ſeveral caſes, where lixivium, ſoap, and lime-water, have been given as remedies.

The ſolubility of others, joined with the teſtimony of reputable authors, and my own experience for near thirty years, confirm the ſalutary effects of lixivium in many caſes.

It frequently happens, in fits of the gravel and ſtone, that gravel or ſmall pieces of calculi are diſcharged, which ſhould be examined.

If perfectly ſoluble in lixivium *(Aq. kali puri)*, the remedy is obvious; if imperfectly, doubtful; if inſoluble, lixivium will only irritate, without benefit.

I am, &c.

T. LANE.

Alderſgate-ſtreet,
March 10, 1791.

XV. *Chermes Lacca.* *By* William Roxburgh, *M. D. of* Samulcotta. *Communicated by* Patrick Ruſſell, *M. D. F. R. S.*

Read May 19, 1791.

SOME pieces of very freſh-looking lac, adhering to ſmall branches of Mimoſa cinerea *, were brought me from the mountains, on the 20th of November, 1789. I kept them carefully in wide-mouthed cryſtal bottles, ſlightly covered; and this day, the 4th of December, fourteen days from the time they came from the hills, thouſands of exceeding minute red animals were obſerved crawling about the lac and the branches it adhered to, and ſtill more were iſſuing from ſmall holes on the ſurface of the cells. By the aſſiſtance of glaſſes, ſmall imperforated excreſcences were alſo obſerved, interſperſed among theſe holes, two, regularly, to each hole, crowned with ſome very fine white hairs, which being rubbed off, two white ſpots appeared. The animals, when ſingle, ran about pretty briſkly; but, in general, on opening the cells, they were ſo numerous as to be crowded over one another. The ſubſtance of which the cells were formed cannot be better deſcribed, with reſpect to appearance, than by ſaying it is like the tranſparent amber that beads are made of. The external covering of the cells may be about half a line thick, is remarkably ſtrong, and able to reſiſt injuries: the partitions are much thinner. The cells are in general irregular ſquares, pentagons, and hexagons, about an

* Lac, on this coaſt, is always found upon the three following ſpecies of Mimoſa; 1ſt, a new ſpecies, called by the Gentoos Conda corinda; 2d, Mimoſa glauca of Kœnig; and, 3dly, Mimoſa cinerea of Linnæus.

eighth

eighth of an inch in diameter, and a quarter of an inch deep: they have no communication with each other. All thoſe I opened, during the time the animals were iſſuing from them, contained in one ſide, and which occupied half the cell, a ſmall bag, filled with a thick red jelly-like liquor, replete with what I take to be eggs. Theſe bags, or utriculi, adhere to the bottom of the cells, and have each two necks, which paſs through perforations in the external coat of the cells, forming the before-mentioned excreſcences, ending in ſome very fine hairs.

The other half of the cells have a diſtant opening, and contain a white ſubſtance, like ſome few filaments of cotton rolled together, and a number of the little red inſects themſelves crawling about, ready to make their exit. Their portion of each cell is about a half; and, I think, muſt have contained near one hundred of theſe animals. Other cells, leſs forward, contained in this half with one opening, a thick, red, dark blood-coloured liquor, with numbers of exceedingly minute eggs, many times ſmaller than thoſe found in the ſmall bags which occupied the other half of the cells. Several of theſe inſects I obſerved to have drawn up their legs, and to lie flat; they did not move on being touched; nor did they ſhew any ſigns of life upon the greateſt irritation *.

Dec. 5. The ſame minute hexapodes continue iſſuing from their cells in numbers.

Dec. 6. The male inſect, I have found to-day, at leaſt what I think is ſuch. A few of them are conſtantly running about, and over the little red inſects, (which I ſhall now call the female) moſt actively: as yet they are ſcarce, not more, I imagine, than 1 to 5000 females, but they are four or five times their ſize.

* It will appear in the ſequel, that theſe were on the point of tranſformation into the pupa ſtate.

To-day the female insects continue issuing in great numbers, and move about as before.

Dec. 7. The small red or female insects, are still more numerous, and move about as before. The winged or male insects, are still very few, but continue active. There have been fresh leaves and bits of the branches of Mimosa cinerea, and Mimosa intsia, put in to them. They go over them indifferently, without shewing any preference or inclination to work, or to copulate. I opened a cell, from whence I thought the winged flies had come, and found several (eight or ten) struggling to shake off their incumbrances. They were in one of those utriculi mentioned before, which end in two mouths, shut up with fine white hairs; but one of them was open for the exit of the flies; the other would, no doubt, have opened in due time. This utriculus I found now perfectly dry; and could plainly see it was divided into minute cells, by exceedingly thin membraneous partitions. I imagine, before any of the flies made their escape, it might have contained about sixteen or twenty. In the minute cells, with the living flies, or from whence they had made their escape, were small dark-coloured compressed grains.

March 26, 1790, I found some branches of the same sort of Mimosa, with numbers of the minute red hexapodes, mentioned in December (seemingly in their pupa state), adhering to them. They are of various sizes, from half a line to a line and a half in length. I found many of the large ones empty. They have a round opening at the lower end, with a small round operculum, or lid, which now loosely covers the empty husk or shell: the inside of these is lined with a small white membrane; others were still shut, some were opening, and some half open, with the insects projecting more or less, and soon extricating themselves entirely.

I opened

I opened ſome of the middle-ſized, and found they contained a thick, deep blood-coloured liquid; others, ſtill larger, put on the appearance of the fly, which was ſoon to iſſue, retrograde.

Deſcription of the male lac inſect in its perfect ſtate.

It is then about the ſize of a very ſmall fly, and exceedingly active; the larva and pupa ſtate, I am as yet unacquainted with.

Head obtuſe; between the eyes a beautiful, ſhining green.

Eyes, black, very large in proportion to the animal.

Antennæ, clavated, feathered, about two-thirds the length of the body; below the middle, an articulation, ſuch as thoſe in the legs.

Mouth: I could not diſtinctly ſee it.

Trunk, oval, brown.

Abdomen, oblong, length of the trunk and head.

Extremities. Vide p. 232.

Legs, ſix; with them it runs briſkly, and jumps actively.

Wings, four, membranaceous, longer than the abdomen, incumbent; the anterior pair twice the ſize of the poſterior.

Tail, none.

Deſcription of the female lac inſect.

Larva, red, very minute, requiring a good lens to diſtinguiſh its parts.

Head, ſcarcely to be diſtinguiſhed from the trunk.

Antennæ, filiform, bifid, hairy, length of the inſect.

Eyes: in the back part of the trunk are two minute elevations, which may be they.

Mouth,

Mouth, on the middle of the breaſt, between the firſt pair of legs, which the little animal projects on being injured, otherwiſe it cannot be ſeen.

Trunk and *Abdomen*, oblong, compreſſed, tapering equally towards each end, croſſed with twelve annular ſegments, margins very flat, and ſeem to be marked with a double line.

Extremities.

Legs, ſix, running, does not jump.

Wings, none.

Tail, two ſlender white hairs, as long as the antennæ, with a white point, which may be called the rump, between them.

Pupa: the duration and peregrinations of the larvæ ſeem very ſhort and confined; for, in a few days after iſſuing from their cells they fix themſelves on the ſmall, but hard, woody branches of the tree they were produced on; it ſeeming impoſſible that they can in this ſtate tranſport themſelves to any other. About the end of December, or beginning of January, they have done iſſuing from their cells, and are ſticking faſt to the branches, regularly with their heads towards the extremity of the branch. The legs, antennæ, and tail, are now entirely gone. Their progreſs through this ſtate is ſlow, requiring about three months. Soon after they have ſettled themſelves, they become covered with a hard, brittle, garnet-coloured cruſt, ſimilar to the lac of which the cells are made, but of a brighter colour. They retain only a rude reſemblance of their former ſhape. About the end of March they have acquired three or four times their original ſize; a ſmall, round lid or cover is now obſerved at the lower part, which opens, but does not always fall off, and gives a retrograde paſſage for the fly, now in its perfect ſtate.

The

The insect in its perfect state is rather smaller than the male, of a brighter red colour, and less active.

Head, small in proportion to the body, pointed.

Eyes, very minute.

Antennæ, filiform, not articulated as in the male, spreading, somewhat shorter than the insect.

Mouth: I could not discover it distinctly.

Trunk, red, almost orbicular.

Abdomen, red, oblong, composed of twelve annular segments.

Extremities.

Legs, six, for running or jumping.

Wings, two, incumbent, longer than the abdomen, transparent.

Tail, two white hairs as long as the insect.

With regard to the œconomy of these little animals, I must, for the present, be silent; having little more than conjecture to offer on that head.

The eggs, and dark-coloured glutinous liquor they are found in, communicate to water a most beautiful red colour, while fresh. After they have been dried, the colour they give to water is less bright; it would therefore be well worth while for those, who are situated near places where the lac is plentifully found, to try to extract and preserve the colouring principles by such means as would prevent them from being injured by keeping. I doubt not but in time a method may be discovered to render this colouring matter as valuable as cochineal.

Mr. Hellot's process for extracting the colouring matter from dry lac deserves to be tried with the fresh lac in the month of October, or beginning of November, before the insects have acquired life; for I found the deepest and best colour was procured from the eggs while mixed with their nidus.

His proceſs is as follows. Let ſome powdered gum lac be digeſted two hours in a decoction of comfrey root, by which a fine crimſon colour is given to the water, and the gum is rendered pale or ſtraw coloured. To this tincture, poured off clear, let a ſolution of alum be added; and when the colouring matter has ſubſided, let it be ſeparated from the clear liquor and dried; it will weigh about one fifth of the quantity of lac employed. This dried fecula is to be diſſolved or diffuſed in warm water; and ſome ſolution of tin is to be added to it, by which it acquires a vivid ſcarlet colour. This liquor is to be added to a ſolution of tartar in boiling water; and thus the dye is prepared.

In India comfrey roots are not to be had; but any other mucilaginous root, gum, or bark, would probably anſwer equally well. On ſome parts on the Coromandel coaſt, if not over it all, a decoction of the ſeeds of a very common plant (Caſſia tora of LINNÆUS), which is exceedingly mucilaginous, is uſed by the dyers of cotton cloth blue, to help to prepare the blue vat. It ſuſpends the indigo till a fermentation takes place to diſſolve it, and alſo helps to bring about that fermentation earlier than it otherwiſe would. The gum lac (or rather reſin) itſelf is known to be perfectly ſoluble in ſpirits of wine. The empty huſks which covered the pupa are alſo ſoluble in ſpirits, but without a very large proportion of the ſpirits is uſed, it ſoon becomes thick, like a jelly. Four grains communicated that quality to three drams of rectified ſpirits of wine. This jelly is very difficult of ſolution in ſpirits; a month has not effected it in a heat of from 80 to 90 degrees of FAHRENHEIT's ſcale. The ſubſtance of which theſe huſks are compoſed, is an exudation from the larvæ themſelves, which becomes hard by expoſure to the air. The cells ſeem

to be made of a very different ſubſtance; what that is, and the manner in which they are made, remains ſtill to be diſcovered.

Explanation of the figures. Tab. VI.

1. A piece of lac on a ſmall branch of Mimoſa cinerea, natural ſize.

2. The outſide of the top of a cell, with its three openings; the white one with the hairs is ſtill unopened.

3. One of the utriculi for the male flies, with its two necks, which correſpond with the two upper apertures in fig. 2.

4. One of the eggs found in the utriculus, fig. 3. which produces the male flies.

5. The male fly in its perfect ſtate.

6. Small compreſſed dry grains, found in the cellulæ with the male flies. The laſt five figures are all much magnified.

7. A ſmall bit of a branch of Mimoſa cinerea, with the female inſects in their pupa ſtate, natural ſize.

8. One of the eggs which produce the female larva. They are always in that portion of the cell from whence the larva iſſues.

9. The female larva.

10. ——— pupa.

11. ——— —— with the lid opening, and the inſect protruding.

12. The female fly in its complete ſtate.

The laſt five figures are much magnified.

XVI. *The Longitudes of* Dunkirk *and* Paris *from* Greenwich, *deduced from the Triangular Measurement in* 1787, 1788, *ſuppoſing the Earth to be an Ellipſoid. By Mr.* Iſaac Dalby; *communicated by* Charles Blagden, *M. D. Sec. R. S.*

Read May 19, 1791.

IN the account of the Trigonometrical Operation in 1787, 1788, which is given in the Philoſophical Tranſactions, Vol. LXXX. after the diſtance of Dunkirk from the meridian of Greenwich has been determined on a parallel to the perpendicular at Greenwich, its longitude is found by ſpherical computation, on a ſuppoſition, that the ſurface of a ſphere nearly coincides with that of the earth in an eaſt and weſt direction, where the operation was performed; and the magnitude of this ſphere, or which amounts to the ſame thing, the value in parts of a degree, &c. of a meaſured arc on its ſurface (for as ſuch the arc between the meridians of Botley Hill and Goudhurſt may be conſidered) has been determined by actual obſervation at two ſtations nearly in the latitude of Dunkirk; and this independent of any hypotheſis which can ſenſibly affect the concluſion. The principles, though not ſtrictly geometrical, admit of little objection; and therefore, as much care was taken in obſerving the angles at theſe ſtations, upon which the directions of the meridians depend, the longitude of Dunkirk (and conſequently that of Paris) as given in the Table, Vol. LXXX.


1
2
7
8
9



Hayter Sc. 1791.

p. 232. must be nearly true, whatever may be the real figure of the earth. But, it may be ſaid, that the arc between the meridians of Botley Hill and Goudhurſt (17′¼) is too ſhort to infer from obſervation the value of the arc between the meridians of Greenwich and Dunkirk (amounting to near a degree and a half), ſufficiently accurate for finding the longitude to great preciſion; becauſe it has been remarked in the Appendix to the ſame Volume, that an error of 1″, in either of the horizontal angles at the above ſtations, would cauſe a variation of near 6″ of a degree in the longitude of Dunkirk or Paris.

M. BOUGUER's ſpheroid agreeing nearly with the meridional meaſurements, it was adopted for the purpoſes of latitude. But the degree perpendicular to the meridian in latitude 51° 6′ 53″ is found to be 61248 fathoms (Vol. LXXX. p. 215.) which falls ſhort of M. BOUGUER's degree about 22 fathoms; therefore, ſuppoſing the directions of the meridians to have been very accurately determined, the earth cannot be this ſpheroid, notwithſtanding the ingenious hypotheſis reſpecting the curve of the meridian. But it is alſo well known, that the meaſured degrees of latitude in different places are inconſiſtent with an elliptical meridian: for, ſuppoſe an ellipſoid to be determined with the degrees found at the equator and polar circle, the computed degrees in middle latitudes will be much longer than the meaſured ones, as it is well known; and the whole meridional arc between Greenwich and Paris will, on ſuch an ellipſoid, exceed the meaſured arc by a quantity anſwering to about 21″ of latitude. It is evident, however, that if we ſuppoſe ſmall errors to have taken place in determining the celeſtial arcs, or differences of latitude in ſome of the operations (for there is little doubt but the terreſtrial menſurations in general have been

The length of the whole meridional arc between Greenwich and Paris on this ellipsoid is six fathoms greater than the measured arc; the degree in latitude 48° 43′, 16 fathoms less; in latitude 45°, 10 fathoms less; in 43°, 13 fathoms greater; and that in latitude 39° 12′, 54 fathoms greater. The degrees at the equator and polar circle are considerably less than the measured ones, conformable to the hypothesis.

Suppose CE, CP (fig. 1.), are the greater and less semi-axes of the ellipsoid; G Greenwich; PGE its meridian; PD the meridian of Dunkirk; and let GBA be perpendicular to the curve of the meridian at G; then GA will be the shorter axis of the elliptical section which is the perpendicular to the

at P, G, E, will be as PC^3, GB^3, and $\left(\frac{PC^2}{CE}\right)^3$, because at the point E (or equator) the line so drawn will become the radius of curvature itself, or $\frac{PC^2}{CE}$. Therefore $GB^3 : \left(\frac{PC^2}{CE}\right)^3$ :: *rad. curv. at* G : *rad. curv. at* E :: *length of a deg. in the lat. of* G : *length of a deg. at* E, the equator. Let the arc ERL be described with the radius CE; draw CR parallel to GB, RS parallel to PC, and join CK; then, by the nature of the ellipse, CR (CE) : CK :: GB : *half the parameter*, or $\frac{CP^2}{CE}$; therefore $CE^3 : CK^3 :: GB^3 : \left(\frac{CP^2}{CE}\right)^3$ :: 60844 : 60344 (*supposing the lat. of the point* G *to be* 50° 9′½), or CE (CR) : CK :: $\overline{60844}^{\frac{1}{3}} : \overline{60344}^{\frac{1}{3}}$; but CR : CK :: *sine* SKC : *sine* KRC (*co-lat.*); therefore, $\left(\frac{60844}{60344}\right)^{\frac{1}{3}} \times$ *cosine lat.* = *sine* SKC; hence the angle SCK is given (50° 1′ 35″½); therefore, as *tang.* SCK : *tang. lat.* (SCR) :: SK : SR :: *lesser semi-axis* CP : *greater* CE. And putting $d = 57.295779$, &c. *the degrees in the circular arc which is equal to the radius)* we have $\left(\frac{tang.\ lat}{tang.\ SCK}\right)^2 \times 60344\,d = 3489932$ fathoms the longer semi-axis; and $\frac{tang.\ lat.}{tang.\ SCK} \times 60344\,d = 3473656$, the shorter.

meridian

meridian at Greenwich, and the angle EBG will be the latitude of Greenwich, or 51° 28′ 40″. Let HO (parallel to GA) be the ſection of the parallel to that perpendicular, paſſing through Dunkirk. Then by the Table, p. 232. (Vol. LXXX.) the arc GH is 152549 feet; but this arc exceeds the real diſtance of the parallels GA, HO, not more than a fathom; therefore this diſtance may be taken = 25424 fathoms. Now the ſections GA, HO, of the ellipſoid being ſimilar, from the known properties of the figure, we ſhall get HO the ſhorter axis of the ſection of the parallel = 6959396, its longer axis = 6979374, and HW = 3531757 fathoms, W being the point where HO cuts the axis PI of the ellipſoid. Hence, if D be Dunkirk, and the arc HD the meaſured arc of the parallel, we have given the length of this arc, or 547058 feet (Table, p. 232.) = 91176 fathoms, and alſo the point W in the leſſer axis of the ſection HO, to determine the angle HWD in the plane of this ſection. But reverting the ſeries which exhibits the length of an elliptical arc in terms of the *abſciſs* and *ordinate*, will be of little uſe in the preſent caſe, where the arc and its chord are very near of the ſame length: For, let HKOL (fig. 2.) be the ſection of the parallel, where HO = 6959396, and KL = 6979374, are the axes; and HW = 3531757, as in *fig.* 1.; alſo, ſuppoſe HS is the radius of curvature at H, or at the middle of HD; then, if we conceive the arc HD to be a right line, or deſcribed with the radius HW, or with HS (3499700) and thence determine the angle SWD from the two ſides SD, SW, and the included angle (the ſupplement of HSD); in either caſe we get the angle HWD the ſame, or 1° 28′ 44″,8 to within 1″. This angle being obtained, the inclination of the planes PHW, PDW (the planes of the meridians of

Greenwich and Dunkirk, *fig.* 1.), or the longitude of the point D, will be found by the common proportion which in a right-angled ſpherical triangle determines an angle when the legs are given: this will be obvious by conceiving a ſphere (of any magnitude) to be deſcribed about W as a center.

Hence, as *rad.* : *cotang.* angle HWD (1° 28′ 44″.8) :: *ſine* angle HWP (38° 31′ 20″) : *cotang.* 2° 22′ 26″$\frac{1}{2}$, the inclination of the planes of the meridians PH, PD, or longitude of Dunkirk on this ellipſoid. And as the difference of meridians of Paris and Dunkirk is 2′ 21″.9 (for this will not be materially affected by different hypotheſes) the longitude of Paris will be 9 m. 20$\frac{1}{2}$ s. in time. The longitude of Dunkirk from Paris (2′ 21″.9) is the mean longitude deduced at p. 223. (Vol. LXXX.), which is only 1″.1 leſs than that given in the *Connoiſſance des Temps*, 1788.

The method of computing the latitude of the point D (was it neceſſary) is thus: as *rad.* : *coſine* DWH :: *coſine* HWP : *coſine* DWP; and ſince the point W in the axis PW is given, and alſo the angle DWP in the plane of the meridian PD (by the foregoing proportion), the point D will be determined by the properties of the ellipſe; which in fact is nothing more than finding the inclination of the vertical at the point D with the given line DW, which inclination added to the angle DWP, gives the co-latitude of the point D. And hence may be evinced the truth of what is advanced at p. 199. (Vol. LXXX.), that *if the value of an arc on a ſpheroid, conſidered as an arc of a great circle perpendicular to the meridian, be given, the longitude may be found by ſpherical computation, but not the latitude.* For conceive the arc HD to be perpendicular to the meridian at H, then the angle HWP would be the co-latitude

of the point H; and the former proportion would give the longitude of D, whether the figure was a ſphere or ſpheroid; and the angle DWP (found by the latter proportion) would be the co-latitude of D ſuppoſing it a ſphere, in which caſe the point W becomes the center; but this will not hold in a ſpheroid, becauſe DW would not be perpendicular to the meridian at D.

The forgoing method of computing the longitude from the meaſured arc of a parallel on a given ellipſoid (though evidently the direct one), will be tedious, eſpecially when the lengths of the meaſured arcs (GH, HD) are very conſiderable. But when the latitude of the point H is determined from the meaſured arc GH (on the known meridian), and the extent of the other arc (HD), or rather the angle HWD, is not more than two or three degrees, the ſame concluſions, extremely near, may be obtained in the following manner, which is nearly the ſame as the method uſed in computing the longitudes in the Table of General reſults, p. 232. (Vol. LXXX.).

Suppoſe G and D (fig. 1.) to be Greenwich and Dunkirk; PH, PD, their meridians, as before; and let HD (inſtead of its being a parallel to the perpendicular at Greenwich) be an arc of an ellipſe cutting the meridian of Greenwich at right angles, ſuppoſe in the point H. Then the arc GH being = 152549 + 50 feet nearly (becauſe the ellipſe which paſſes through D, and is at right angles to the meridian PG, will fall about 50 feet to the ſouth of the point cut by the parallel), therefore the value of the arc GH, or 25433 fathoms, will, on this ellipſoid, be 25′ 4″.4, and conſequently the angle PWH, or the co-latitude of H, is 38° 56′ 24″.4. Now, the radius of curvature of this perpendicular ellipſe at H, the extremity of its leſſer

axis, will be 3499798 fathoms *, which, divided by 57.295779, &c. (the degrees in the circular arc which is equal to the radius), gives 61083 fathoms for a degree on this ellipſe, conſidered as a great circle perpendicular to the meridian at the point H on the ellipſoid; and ſince the length of this arc (HD) will be nearly the ſame as that of the parallel, or 91176 fathoms, its value will be 1° 29′ 33″.6 (the arc DH, or rather the angle DWH). Hence, as *rad.* : *cotang.* 1° 29′ 33″.6 (HWD) :: *ſine* 38° 56′ 24″.4 (HWP) : *cotang.* 2° 22′ 26″.8, the longitude of D, or Dunkirk, the ſame as before, very near; hence the longitude of Paris will be 2° 20′ 4″.9. But the ſame may be obtained from the mean diſtance of the meridians of Greenwich and Paris, or 537950 feet. See p. 599. in the Appendix to Vol. LXXX.

It appears from the foregoing hypotheſis, that the meaſured degrees of the meridian in middle latitudes will anſwer nearly on an ellipſoid whoſe axes are in the ratio aſſigned by Sir ISAAC NEWTON. But this will receive further confirmation from the fifth ellipſoid in the ſecond Table, p. 232. (Vol. LXXX.), where the near agreement between the computed and meaſured arc of the meridian between Greenwich and Perpignan (differing but about 52 fathoms in the extent of 8° 46′ 44″) would be ſomewhat extraordinary, were we certain that the latitude of Perpignan (42° 41′ 56″) is correct; but this is ſuſpected by M. DE LA

* It is not neceſſary to determine the axes of this ellipſe, becauſe when HW is perpendicular to the curve of the meridian, it will (by the nature of the figure) be the radius of curvature of the arc HD at the point H. Hence, if we put v for the *cotang.* and c for the *coſine* of the latitude of the point H, and let a denote the *ſine* of an arc whoſe *tang.* is $\frac{CE}{CP} \times v$; then $\frac{a}{c} \times CE = HW$, by the properties of the ellipſe.

CAILLE.



Fig. 1.



Fig. 2.

CAILLE. See *Mem. de l'Acad.* 1758. The computed arc, however, between Greenwich and Paris is 19 fathoms longer than the measured arc, which answers to a little more than 1″ of latitude.

The longitude of Paris on this ellipsoid is 9 m. $20\frac{4}{10}$ s.

If it be contended, that the operations at the equator and polar circle were as correct as those executed for the like purpose in middle latitudes; and that a kind of mean between the extreme results ought to be preferred; we shall still get an ellipsoid, whose axes are nearly as 229 to 230, by taking the degrees at the equator and polar circle each 70 fathoms less, and that in latitude 50° 9′ $\frac{1}{2}$ as much greater than the measured ones; and the longitude of Paris will be found 9 m. $19\frac{7}{10}$ s. But the computed meridional arc between Greenwich and Paris will exceed the measured one by a quantity answering to about 11″ of latitude.

It is almost needless to observe, that the longitude of Paris (9 m. 20 s.) deduced by Dr. MASKELYNE from the different results found by astronomical observations (Phil. Trans. 1787) agrees to less than half a second with either of the above determinations.

N. B. In the Table of General Results, p. 232. Vol. LXXX. *read* 18′ 46″ for the longitude of Wrotham Hill; this should have been corrected in the Appendix.

XVII. *On the Method of determining, from the real Probabilities of Life, the Values of contingent Reverſions in which Three Lives are involved in the Survivorſhip.* *By Mr.* William Morgan, *F. R. S.*

Read May 26, 1791.

HAVING been encouraged to the further purſuit of the doctrine of ſurvivorſhips by the very honourable manner in which my two former Papers on this ſubject were received by the Royal Society, I think it my duty to ſubmit the reſult of my labours to their conſideration. The ſolutions of ſome of the following problems might have been derived from thoſe which I have already communicated; but the direct inveſtigation of each ſeparate problem being certainly more ſatisfactory, and the rules obtained by this means in general more ſimple, I have conſidered no problem as connected with another, except the relation between them either immediately ariſes from the ſolution, or is neceſſary to prove the truth of it. Being anxious to render myſelf as conciſe as poſſible, I have been minute only in the inveſtigation of the firſt problem, and have done little more than ſtate the contingencies which will determine the ſurvivorſhip in the others. By the aſſiſtance, however, of theſe, and the operations which are detailed in my former Papers, the theorems which I have given may be deduced without much difficulty.

In

In order to prevent unneceſſary repetitions, it may not be improper to begin with explaining the different ſymbols which are uſed in the following pages.

A, B, C, } denote the value of an annuity on the reſpective lives of A, B, or C.

D, denotes the value of S on the contingency of C's ſurviving A (by my 2d prob. Vol. LXXVIII).

E, denotes the ſame value on the contingency of B's ſurviving A, found by the ſame problem.

F, denotes the value of an annuity on a life one year younger than B.

G, denotes the value of the abſolute reverſion of S after the death of A.

H, denotes the value of an annuity on a life one year younger than A.

K, denotes the value of an annuity on a life one year younger than C.

L, denotes the value of an annuity on the longeſt of the three lives of A, B, and C.

M, denotes the value of S by the firſt problem on the contingency that A's life ſhall be the *firſt* that fails.

N, denotes the value of an annuity on a life one year older than A.

P, denotes the value of an annuity on a life one year older than B.

R, denotes the value of S on the contingency of B's dying after A (by my 3d prob. Vol. LXXVIII).

S, denotes the given ſum.

T, denotes the value of an annuity on a life one year older than C.

V, denotes

V, denotes the perpetuity.

W, denotes the value of S on the contingency of C's dying after A (by my 3d prob. Vol. LXXVIII).

α and a denote the number of perſons living in a table of obſervations at the ages of H and A.

s, t, u, w, &c. denote the number of perſons living at the end of the 1ſt, 2d, 3d, &c. years from the age of A.

β and b, denote the number of perſons living at the ages of F and B.

m, n, o, p, &c. denote the number of perſons living at the end of the 1ſt, 2d, 3d, &c. years from the age of B.

$\varkappa$ and c denote the number of perſons living at the ages of K and C.

d, e, f, g, &c. denote the number of perſons living at the end of the 1ſt, 2d, 3d, &c. years from the age of C.

a', a'', a''', &c.
b', b'', b''', &c.
c', c'', c''', &c.
} denote the decrements of life at the end of the 1ſt, 2d, 3d, &c. years from the reſpective ages of A, B, and C.

r, denotes the value of £. 1 increaſed by its intereſt for a year.

The combinations of two or three of the ſeveral letters A, B, C, F, H, &c. denote the values of annuities on the joint continuance of two or three of thoſe reſpective lives.

PROBLEM I.

To determine the value of a given ſum, payable if A ſhould be the *firſt* that fails of the three lives A, B, and C.

SOLUTION.

In order to receive the given ſum in the firſt year, it is neceſſary that one or other of four events ſhould happen. 1ſt, That

That all the three lives ſhould fail, and that A ſhould die firſt. 2dly, That B ſhould die *after* A, and C live. 3dly, That C ſhould die *after* A, and B live. 4thly, That A only ſhould die, and B and C both live. Theſe ſeveral contingencies being expreſſed by $\frac{a' \cdot \overline{b-m} \cdot \overline{c-d}}{3abc}$, $\frac{a' \cdot \overline{b-m} \cdot d}{2abc}$, $\frac{a' \cdot \overline{c-d} \cdot m}{2abc}$, and $\frac{a' \cdot dm}{abc}$, reſpectively, their ſum will be $= \frac{1}{abc} \times \overline{\frac{a'bc}{3} + \frac{a'md}{3} + \frac{a'mc}{6} + \frac{a'bd}{6}}$. In the ſecond and following years one or other of the ſame events muſt take place in order to receive the given ſum; that is, they muſt either all three die, A dying firſt; or only A and B muſt die, A dying firſt; or only A and C muſt die, A dying firſt; or only A muſt die, and B and C both live. The different fractions expreſſing thoſe four contingencies for the ſecond year being $\frac{a'' \cdot \overline{m-n} \cdot \overline{d-e}}{3abc}$, $\frac{a'' \cdot \overline{m-n} \cdot e}{2abc}$, $\frac{a'' \cdot \overline{d-e} \cdot n}{2abc}$, and $\frac{a'' \cdot ne}{abc}$; for the third year $\frac{a''' \cdot \overline{n-o} \cdot \overline{e-f}}{3abc}$, $\frac{a''' \cdot \overline{n-o} \cdot f}{2abc}$, $\frac{a''' \cdot \overline{e-f} \cdot o}{2abc}$, and $\frac{a''' \cdot of}{abc}$, and ſo on, the whole value of the given ſum will be $=$ $\frac{S \cdot a'}{abcr} \times \overline{\frac{bc}{3} + \frac{md}{3} + \frac{mc}{6} + \frac{bd}{6}} + \frac{S \cdot a''}{abcr^2} \times \overline{\frac{me}{3} + \frac{ne}{3} + \frac{nd}{6} + \frac{me}{6}} + \frac{S \cdot a'''}{abcr^3} \times \overline{\frac{ne}{3} + \frac{of}{3} + \frac{oe}{6} + \frac{nf}{6}}$ &c. which are $= \frac{S}{3abc} \times \overline{\frac{a'bc}{r} + \frac{a''md}{r^2} + \frac{a''' \cdot en}{r^3}} +$ &c. $+ \frac{S}{6abc} \times \overline{\frac{a'mc}{r} + \frac{a''nd}{r^2} + \frac{a''' \cdot oe}{r^3}} +$ &c. $+ \frac{S}{6abc} \times \overline{\frac{a'db}{r} + \frac{a''em}{r^2} + \frac{a''' \cdot fn}{r^3}} +$ &c. $+ \frac{S}{3abc} \times \overline{\frac{a'md}{r} + \frac{a'' \cdot en}{r^2} + \frac{a''' fo}{r^3}} +$ &c. From the demonſtration of the problem in my laſt Paper * it appears, that the firſt of theſe ſeries is $= \frac{S}{3} \times \frac{\beta\alpha \cdot \overline{FK-AFK}}{bc} - \frac{S}{3r} \times \overline{BC-ABC}$; that the ſecond ſeries is $= \frac{S}{6} \times \frac{\alpha \cdot \overline{BK-ABK}}{c} - \frac{S}{6r} \times \frac{m \cdot \overline{PC-APC}}{b}$; that the

* See Phil. Tranſ. Vol. LXXIX.

third series is $= \frac{S}{6} \times \frac{\beta \cdot \overline{FC-AFC}}{b} - \frac{S}{6r} \times \frac{d \cdot \overline{BT-ABT}}{c}$; and that the fourth series is $= \frac{S}{3} \times \overline{BC-ABC} - \frac{S}{3r} \times \frac{md \cdot \overline{PT-APT}}{bc}$. These several expreſſions being added together will be found $= S$ into

$$\frac{\alpha}{3r} \times \frac{\overline{\beta \cdot FK-AFK}}{b} + \frac{\overline{BK-ABK}}{2} + \frac{\beta}{6b} \times \overline{FC-AFC} + \frac{\overline{-1}}{3r} \times \overline{BC-ABC} - \frac{m \cdot \overline{PC-APC}}{6br} - \frac{d}{3cr} \times \frac{\overline{BT-ABT}}{2} + \frac{m \cdot \overline{PT-APT}}{b}.$$

This theorem gives the exact value when either B or C are the oldeſt of the three lives; but when A is the oldeſt, it will be neceſſary to exchange the ſymbols a', a'', a''', &c. for $\overline{a-s}$, $\overline{s-t}$, $\overline{t-u}$, &c. and the ſymbols m, n, o, &c. for $\overline{c-c'}$, $\overline{c-c'+c'}$, $\overline{c-c'+c''-c'''}$, &c. In this caſe the value of the given ſum for the firſt year will be found $= \frac{S}{abcr}$ into $\frac{abc}{2} - \frac{bsc}{2} + \frac{amc}{2} - \frac{msc}{2} - \frac{abc'}{6} + \frac{bsc'}{6} - \frac{amc'}{3} + \frac{msc'}{3}$; for the ſecond year $= \frac{S}{abcr^2}$ into $\frac{msc}{2} - \frac{mtc}{2} + \frac{nsc}{2} - \frac{ntc}{2} - \frac{msc''}{6} + \frac{mtc''}{6} - \frac{nsc''}{3} + \frac{ntc''}{3} - \frac{msc'}{2} + \frac{mtc'}{2} - \frac{nsc'}{2} + \frac{ntc'}{2}$; for the third year $= \frac{S}{abcr^3}$ into $\frac{ntc}{2} - \frac{nuc}{2} + \frac{otc}{2} + \frac{euc}{2} - \frac{ntc'''}{6} + \frac{nuc'''}{6} - \frac{otc''}{3} + \frac{euc'''}{3} - \frac{nt \cdot \overline{c'+c''}}{2} + \frac{nu \cdot \overline{c'+c''}}{2} - \frac{ot \cdot \overline{c'+c''}}{2} + \frac{ou \cdot \overline{c'+c''}}{2}$; and ſo on for the remaining years of A's life. Hence the whole value of the given ſum will be $= \frac{S}{2ab} \times \overline{\frac{ab}{r} + \frac{ms}{r^2} + \frac{nt}{r^3}} + \&c. - \frac{S}{2ab} \times \overline{\frac{bs}{r} + \frac{mt}{r^2} + \frac{nu}{r^3}} + \&c. + \frac{S}{2ab} \times \overline{\frac{am}{r} + \frac{ns}{r^2} + \frac{ot}{r^3}} + \&c. - \frac{S}{2ab} \times \overline{\frac{ms}{r} + \frac{nt}{r^2} + \frac{ou}{r^3}} + \&c. - \frac{S}{6abc} \times \overline{\frac{abc'}{r} + \frac{msc''}{r^2} + \frac{mtc'''}{r^3}} + \&c. - \frac{S}{2abcr} \times \overline{\frac{msc'}{r} + \frac{nt \cdot \overline{c'+c''}}{r^3}} + \&c. + \frac{S}{6abc} \times \overline{\frac{bsc'}{r} + \frac{mtc''}{r^2} + \frac{nuc'''}{r^3}} + \&c. + \frac{S}{2abcr} \times$

mtc'

$\frac{mtc'}{r} + \frac{nu \cdot \overline{c'+c''}}{r^2} + \&c. - \frac{S}{3abc} \times \frac{amc'}{r} + \frac{ns \cdot c''}{r^2} + \frac{ot \cdot c'''}{r^3} + \&c. - \frac{S}{2abcr} \times \frac{nsc'}{r} + \frac{ot \cdot \overline{c'+c''}}{r^2} + \&c. + \frac{S}{3abc} \times \frac{msc'}{r} + \frac{nt \cdot c''}{r^2} + \frac{ouc'''}{r^3} + \&c. \times \frac{S}{2abcr} \times \frac{nt \cdot c'}{r} + \frac{ou \cdot \overline{c'+c''}}{r^2} + \&c.$ . . . The firſt four ſeries are reſpectively $=$ $\frac{a\beta \cdot HF}{2ab}$ (or $\frac{1+AB}{2r}$) $- \frac{\beta \cdot AF}{2b} + \frac{a \cdot HB}{2a} - \frac{AB}{2}$ . . . The firſt term of the fifth ſeries, or $\frac{abc'}{r}$, is $= \frac{a\beta}{abr} \times \frac{ab}{a\beta} - \frac{ab \cdot \overline{c-c'}}{a\beta c}$, the ſecond term or $\frac{msc''}{r^2}$, is $= \frac{a\beta}{abr^2} \times \frac{ms}{a\beta} - \frac{ms \cdot \overline{c-c'+c'}}{a\beta c} - \frac{msc'}{r^2}$ . . . . the third term, or $\frac{ntc'''}{r^3}$, is $= \frac{a\beta}{abr^3} \times \frac{nt}{a\beta} - \frac{nt \cdot \overline{c-c'+c''+c'''}}{a\beta c} - \frac{nt' \cdot \overline{c'+c''}}{r^3}$. Therefore the ſum of the fifth and ſixth ſeries is $= - \frac{a\beta \cdot \overline{HF-HFC}}{6ab} \left(- \frac{msc'}{3abcr^2} - \frac{nt \cdot \overline{c'+c''}}{3abcr^3} - \&c. = \right) - \frac{1}{3r} \times \overline{AB-ABC}$. In the ſame manner the ſeventh ſeries may be found $= \frac{\beta}{6b} \times \overline{AF-AFC} - \frac{mtc'}{6abcr^2} - \frac{nu \cdot \overline{c+c''}}{6abcr^3} - \&c.$; and conſequently the ſum of the ſeventh and eighth ſeries is $= \frac{\beta \cdot \overline{AF-AFC}}{6b} + \frac{1 \cdot \overline{BN-BNC}}{3ar}$. Again, the ninth and tenth ſeries may be found $= - \frac{a \cdot \overline{HB-HBC}}{3a} \left(- \frac{nsc'}{6abcr^2} - \frac{ot \cdot \overline{c'+c''}}{6abcr^3} - \&c. = \right) - \frac{m \cdot \overline{AP-APC}}{6br}$; and the eleventh and twelfth or two laſt ſeries, $= \frac{AB-ABC}{3} \left(+ \frac{ntc'}{6abcr^2} + \frac{uo \cdot \overline{c'+c''}}{6abcr^3} + \&c. = \right) + \frac{ms \cdot \overline{PN-PNC}}{6abr}$. If all theſe expreſſions be added together, we ſhall have the value of the given ſum $=$ S into $\frac{\beta}{3b} \times \frac{a \cdot HF + \frac{1}{2}HFC}{a} - \overline{AF + \frac{1}{2}AFC}$

$+\frac{1}{b}\times\frac{\alpha\cdot\overline{HB+2HBC}}{a}-\overline{AB+2ABC}+\frac{1}{3r}-\frac{s\cdot\overline{BN-BNC}}{a}-\overline{AB-ABC}$

$+\frac{m}{6br}\times\frac{s\cdot\overline{PN-PNC}}{a}-\overline{AP-APC}$.

It may be obſerved, that the firſt four ſeries are alſo = E, and therefore if this value be ſubſtituted inſtead of $\frac{\alpha\beta\cdot HF}{2ab}$, &c. the general rule will become = E + S into $-\frac{\alpha}{3a}\times\overline{HB-HBC}+\frac{\beta}{2b}$

$\times\overline{HF-HFC}+\frac{r-1\cdot\overline{AB-ABC}}{3r}+\frac{\beta\cdot\overline{AF-AFC}}{6b}-\frac{m\cdot\overline{AP-APC}}{6br}+\frac{s}{3ar}$

$\times\overline{BN-BNC}+\frac{2b}{m}\times\overline{PN-PNC}$ ..... Suppoſing the three lives to be equal, the firſt of theſe rules will become $=\frac{1+CC}{2r}-$

$\frac{xx\cdot\overline{KK-CKK}}{6\cdot cc}-\frac{1}{3r}\times\overline{CC-CCC}-\frac{s\cdot\overline{CK-CCK}}{6c}+\frac{d\cdot\overline{CT-CCT}}{6cr}-$

$\frac{CC+2CCC}{6}+\frac{dd\cdot\overline{TT-CTT}}{6\cdot cc\cdot r}$, and the ſecond $=\frac{r-1\cdot\overline{V-CC}}{2r}+$

$\frac{r-1\cdot\overline{CC-CCC}}{3r}-\frac{s\cdot\overline{CK-CCK}}{6c}-\frac{xx\cdot\overline{KK-CCK}}{6cc}+\frac{d\cdot\overline{CT-CCT}}{6cr}+$

$\frac{dd\cdot\overline{TT-CTT}}{6ccr}$: and alſo the rule denoting the value (when either B or C are the eldeſt of the three lives) will become $=$

$\frac{r-1\cdot\overline{CC-CCC}}{3r}+\frac{s}{3c}\times\overline{CK-CCK}+\frac{xx}{3\cdot cc}\times\overline{KK-CKK}-\frac{d}{c}\times$

$\frac{\overline{CT-CCT}}{3r}-\frac{dd.}{3ccr}\times\overline{TT-CTT}$. Let this laſt expreſſion be called Q and compared with the firſt of the preceding expreſſions. In this caſe we ſhall have $\frac{1+CC}{2r}=\frac{CC+2CCC}{6}-\frac{\overline{CC-CCC}}{3r}-\frac{Q}{2}+$

$\frac{r-1\cdot\overline{CC-CCC}}{6r}=Q$, from which Q may be eaſily found $=$

$\frac{S}{3}\times\frac{r-1\cdot\overline{V-CCC}}{r}$. Again, let the ſame expreſſion, denoted

by

by Q, be compared with the ſecond of the preceding expreſſions, and we ſhall then have $\frac{r-1 \cdot V-CC}{2r} - \frac{Q}{2} + \frac{r-1 \cdot CC-CC\bar{C}}{2r}$ $= Q$, and conſequently $Q = S \times \frac{r-1 \cdot V-CCC}{3r}$ as before * . . . . Now it is well known, that when the lives are all equal, the value of the reverſion muſt be *one third* the difference between the perpetuity and the three joint lives, and therefore a demonſtration ariſes of the truth of the whole ſolution. As a ſtill further proof of this, the foregoing theorem may be immediately deduced from the ſeries themſelves: thus, the value of the given ſum for the firſt year will in this caſe be $= \frac{S \cdot c^3 - d^3}{3c^3 \cdot r}$; for the ſecond year it will be $= \frac{S \cdot d^3 - e^3}{3c^3 r^2}$; for the third year it will be $= \frac{S \cdot e^3 - f^3}{3c^3 r^3}$. Hence the value of the whole reverſion will be $=$

$$\frac{S}{3} \times \frac{1}{r} - \frac{d^3}{c^3 r} - \frac{e^3}{c^3 r^2} - \&c. + \frac{S}{3} \times \frac{d^3}{c^3 r^2} + \frac{e^3}{c^3 r^3} + \frac{f^3}{c^3 r^4} + \&c. = \frac{S}{3} \times$$

$$\frac{1}{r} - CCC + \frac{CCC}{r} = \frac{S}{3} \times \frac{r-1 \cdot V-CCC}{r} \ldots \text{Q. E. D.}$$

PROBLEM II.

To determine the value of a given ſum, payable if A ſhould be the *ſecond* that fails of the three lives A, B, and C.

SOLUTION.

The ſum S may be received in the firſt year, provided either

* This expreſſion may alſo be obtained from either of the above general rules, independent of the two others, in like manner as in the ſolution of the fourth problem.

of three events ſhould happen; 1ſt, if the three lives ſhould become extinct, and A be the ſecond that fails; 2dly, if A ſhould die after B, and C live; and, 3dly, if A ſhould die after C, and B live. But in the ſecond and following years, the given ſum may be received, provided either of ſeven events ſhould happen. 1ſt, If the three lives ſhould fail in the year, A's life having been the ſecond that failed. 2dly, If A only ſhould die in the year, B having died before the beginning and C lived to the end of it. 3dly, If A only ſhould die in the year, C having died before the beginning and B lived to the end of it. 4thly, If A ſhould die after B in the year and C live. 5thly, If A ſhould die after C in the year and B live. 6thly, if A and C ſhould both die in the year (A dying firſt) and B's life ſhould have failed in one or other of the foregoing years; and, 7thly, if A and B ſhould both die in the year (A dying firſt) and C's life ſhould have failed in the foregoing years. From the ſeveral expreſſions denoting theſe contingencies the whole value of the reverſion may be found =

$$\frac{S}{2ac}\times\overline{\frac{d'c}{r}+\frac{a''d}{r^2}+\frac{a'''e}{r^3}+\&c.}+\frac{S}{2ac}+\overline{\frac{a'd}{r}+\frac{a''e}{r^2}+\frac{a'''f}{r^3}+\&c.}+$$

$$\frac{S}{2ab}\times\overline{\frac{a'b}{r}+\frac{a''m}{r^2}+\frac{a'''n}{r^3}+\&c.}+\frac{S}{2ab}\times\overline{\frac{a'm}{r}+\frac{a''n}{r^2}+\frac{a'''o}{r^3}+\&c.}-$$

$$\frac{2S}{3abc}\times\overline{\frac{a'bc}{r}+\frac{a''md}{r^2}+\frac{a'''ne}{r^3}+\&c.}-\frac{2.S}{3abc}\times\overline{\frac{a'md}{r}+\frac{a''ne}{r^2}+\frac{a'''of}{r^3}+\&c.}-$$

$$\frac{S}{3abc}\times\overline{\frac{a'mc}{r}+\frac{a''na}{r^2}+\frac{a'''oe}{r^3}+\&c.}-\frac{S}{3abc}\times\overline{\frac{a'bd}{r}+\frac{a''me}{r^2}+\frac{a'''nf}{r^3}+\&c.}$$

The two firſt of theſe ſeries are = D, .. the two next = E ... and it appears from the ſolution of the preceding problem that the four remaining ſeries expreſs double the value of the ſum S, depending on the contingency of A's dying firſt (when B or C are the oldeſt of the three lives) with a negative ſign.

The

The general rule, therefore, in this caſe will become $= D + E - 2M$.

But when A is the oldeſt life, recourſe muſt be had, as in the ſecond part of the preceding problem, to different ſymbols, and the value of the reverſion will then be found $=$

$$\frac{S}{ab} \times \overline{-\frac{ab}{r} + \frac{ms}{r^2} + \frac{nt}{r^3} + \&c.} + \frac{S}{ab} \times \overline{\frac{bs}{r} + \frac{mt}{r^2} + \frac{nu}{r^3} + \&c.} +$$

$$\frac{S}{3abc} \times \overline{\frac{abc'}{r} + \frac{msc''}{r^2} + \frac{nsc'''}{r^3} + \&c.} - \frac{S}{3abc} \times \overline{\frac{bsc'}{r} + \frac{mtc''}{r^2} + \frac{nuc'''}{r^3} + \&c.} +$$

$$\frac{2S}{3abc} \times \overline{\frac{amc'}{r} + \frac{nsc''}{r^2} + \frac{otc'''}{r^3} + \&c.} - \frac{2S}{3abc} \times \overline{\frac{msc'}{r} + \frac{ntc''}{r^2} + \frac{ouc'''}{r^3} + \&c.} +$$

$$\frac{S}{abc} \times \overline{\frac{msc'}{r} + \frac{nt \cdot c' + c''}{r^2} + \&c.} - \frac{S}{abc} \times \overline{\frac{mtc'}{r} + \frac{nu \cdot c' + c''}{r^2} + \&c.} +$$

$$\frac{S}{abc} \times \overline{\frac{nsc'}{r} + \frac{ot \cdot c' + c''}{r^2} + \&c.} - \frac{S}{abc} \times \overline{\frac{nt \cdot c'}{r} + \frac{ou \cdot c' + c''}{r^2} + \&c.} +$$

$$\frac{S}{2ab} \times \overline{\frac{a-1 \cdot b}{r} + \frac{1-t \cdot m}{r^2} + \frac{t-u \cdot n}{r^3} + \&c.} + \frac{S}{2ab} \times \overline{\frac{a-1 \cdot m}{r} + \frac{1-t \cdot n}{r^2} + \&c.} +$$

$$\frac{S}{2ac} \times \overline{\frac{a-1 \cdot c}{r} + \frac{1-t \cdot c-c'}{r^2} + \&c.} + \frac{S}{2ac} \times \overline{\frac{a-1 \cdot c-c'}{r} + \frac{1-t \cdot c-c'+c''}{r^2} +}$$

$$\overline{\&c.} - \frac{S}{ab} \times \overline{\frac{am}{r} + \frac{ns}{r^2} + \frac{ot}{r^3} + \&c.} + \frac{S}{ab} \times \overline{\frac{ms}{r} + \frac{nt}{r^2} + \frac{ou}{r^3} + \&c.}$$

The laſt two, and the firſt ten ſeries are $= -2M$, the eleventh and twelfth ſeries are $= D$, and the thirteenth and fourteenth ſeries $= E$; conſequently the general rule becomes $= D + E - 2M$, as before.

Suppoſing the lives were all equal, the above expreſſion would be $= \frac{S \cdot \overline{r-1} \cdot \overline{V - CC}}{r} - \frac{2 \cdot S}{3r} \times \overline{r-1} \cdot \overline{V - CCC} = \frac{S}{3} \times \frac{\overline{r-1}}{r} \times \overline{V - 3CC - 2CCC}$, which is known from other principles * to be the true value, and therefore the inveſtigation is right. As a further demonſtration, however, it may not be improper to

* See Phil. Tranſ. Vol. LXXIX.

obſerve,

obſerve, that this rule is immediately obtained from the different fractions which expreſs the ſeveral contingencies in each year. For in this caſe, the value of S for the firſt year becomes $= \frac{S}{r} \times \frac{1}{3} - \frac{dd}{cc} + \frac{2d^3}{c^3}$, for the ſecond year $= \frac{S}{r^2} \times \frac{dd}{cc} - \frac{ee}{cc} - \frac{2d^3}{c^3} + \frac{2e^3}{c^3}$ and ſo on for the other years. Theſe ſeries being added together will be found $= \frac{S}{3} \times \frac{1}{r} + \frac{2d^3}{c^3 \cdot r} + \frac{2e^3}{c^3 \cdot r^2} + \&c. - \frac{S}{3} \times \frac{3dd}{ccr} + \frac{3ee}{cc \cdot r^2} + \&c. + \frac{S}{3r} \times \frac{3dd}{ccr} + \frac{3ee}{cc \cdot r^2} + \&c. - \frac{S}{3r} \times \frac{2d^3}{c^3 \cdot r^2} + \frac{2e^3}{c^3 \cdot r^2} + \&c. = \frac{S}{3} \times \frac{r-1}{r} \times \overline{V - 3CC - 2CCC} \ldots\ldots$ Q. E. D.

PROBLEM III.

To determine the value of a given ſum payable on the death of A, if his life ſhould be the *laſt* that fails of the three lives A, B, C.

SOLUTION.

The given ſum can be received in the firſt year only upon the extinction of the three lives, A having died laſt. In the ſecond and following years it may be received provided either of four events ſhould happen: 1ſt, if all the three lives ſhould fail in the year, A dying laſt; 2dly, if A ſhould die *after* C in the year, B having died in either of the foregoing years; 3dly, if A ſhould die *after* B in the year, C having died in either of the foregoing years; 4thly, if only A ſhould die in the year, B and C having both died before the beginning of it. The value therefore of the reverſion (when B or C are older than

than A) will be $= \frac{S}{a} \times \overline{\frac{a'}{r} + \frac{a''}{r^2} + \frac{a'''}{r^3} + \&c.} + \frac{S}{3abc} \times \overline{\frac{a'bc}{r} + \frac{a''md}{r^2} + \frac{a'''ne}{r^3} + \&c.} + \frac{S}{3abc} \times \overline{\frac{a'md}{r} + \frac{a''ne}{r^2} + \frac{a'''of}{r^3} + \&c.} + \frac{S}{6abc} \times \overline{\frac{a'me}{r} + \frac{a''nd}{r^2} + \frac{a'''oe}{r^3} + \&c.} + \frac{S}{6abc} \times \overline{\frac{a'bd}{r} + \frac{a''me}{r^2} + \frac{a'''nf}{r^3} + \&c.} - \frac{S}{2ac} \times \overline{\frac{a'c}{r} + \frac{a''d}{r^2} + \frac{a'''e}{r^3} + \&c.} - \frac{S}{2ac} \times \overline{\frac{a'd}{r} + \frac{a''e}{r^2} + \frac{a'''f}{r^3} + \&c.} - \frac{S}{2ab} \times \overline{\frac{a'b}{r} + \frac{a''m}{r^2} + \frac{a'''n}{r^3} + \&c.} - \frac{S}{2ab} \times \overline{\frac{a'm}{r} + \frac{a''n}{r^2} + \frac{a'''o}{r^3} + \&c.} = G + M - \overline{D + E}.$

When the life of A is the oldeſt of the three lives, the ſymbols being changed as in the two preceding problems, the value of the reverſion will become $= -\frac{S}{6abc} \times \overline{\frac{abc'}{r} + \frac{msc''}{r^2} + \frac{ntc'''}{r^3} + \&c.} - \frac{S}{2abcr} \times \overline{\frac{msc'}{r} + \frac{nt \cdot c' + c''}{r^2} + \&c.} + \frac{S}{6abc} \times \overline{\frac{bsc'}{r} + \frac{mtc''}{r^2} + \frac{nuc'''}{r^3} + \&c.} + \frac{S}{2abcr} \times \overline{\frac{mt \cdot c'}{r} + \frac{nu \cdot c' + c''}{r^2} + \&c.} - \frac{S}{3abc} \times \overline{\frac{amc'}{r} + \frac{ns''c}{r^2} + \frac{ot \cdot c'''}{r^3} + \&c.} - \frac{S}{2abcr} \times \overline{\frac{ns \cdot c'}{r} + \frac{ot \cdot c' + c''}{r^2} + \&c.} + \frac{S}{3abc} \times \overline{\frac{msc'}{r} + \frac{nt \cdot c''}{r^2} + \frac{ou \cdot c'''}{r^3} + \&c.} + \frac{S}{2abcr} \times \overline{\frac{nt \cdot c'}{r} + \frac{ou \cdot c' + c''}{r^2} + \&c.} + \frac{S}{2ac} \times \overline{\frac{a-s \cdot c'}{r} + \frac{s-t \cdot c''}{r^2} + \frac{t-u \cdot c'''}{r^3} + \&c.} + \frac{S}{acr} \times \overline{\frac{s-t \cdot c'}{r} + \frac{t-u \cdot c' + c''}{r^2} + \&c.}$

By the ſecond part of the ſolution of the firſt problem, the firſt eight ſeries may be found $= M - E$; and the laſt two ſeries being eaſily reſolved into $\frac{S}{a} \times \overline{\frac{a-s}{r} + \frac{s-t}{r^2} + \frac{t-u}{r^3} + \&c.} - \frac{S}{2ac} \times \overline{\frac{a-s \cdot c-c'}{r} + \frac{s-t \cdot c-c' + c''}{r^2} + \&c.} - \frac{S}{2ac} \times \overline{\frac{a-s \cdot c}{r} + \frac{s-t \cdot c-c'}{r^2} + \&c.}$ are $= G - D$. Hence the general rule, as in the former caſe, becomes $= G + M - \overline{D + E}$.

When the lives are all equal, the above expreſſion will be changed into $\frac{S}{r} \times \overline{r-1} \times \overline{\overline{V-C} + \frac{V-CCC}{3}} - \overline{V-CC} = \frac{S \cdot \overline{r-1}}{r} \times \frac{\overline{V-L}}{3}$; or "*one-third* the value of the abſolute reverſion after "the extinction of the longeſt of the three lives." This is known from ſelf-evident principles to be the true value, and therefore the foregoing ſolution is correct. The ſame rule may alſo be obtained immediately from the ſeries; for the value in this caſe for the firſt year will be $= \frac{S}{r} \times \overline{\frac{1}{3} - \frac{d}{c} + \frac{dd}{cc} - \frac{d^3}{3c^3}}$, for the ſecond year $= \frac{S}{r^2} \times \overline{\frac{d^3}{3c^3} - \frac{e^3}{3^3} - \frac{dd}{cc} + \frac{d}{c} + \frac{ee}{cc} - \frac{e}{c}}$, and ſo on for the remaining years. Hence the whole value of the reverſion will be $= \frac{S}{3r} \times L + \frac{S}{3} \times \overline{\frac{1}{r} - L} = \frac{\overline{r-1} \cdot S}{r} \times \frac{\overline{V-L}}{3}$ . . . . Q. E. D.

PROBLEM IV.

To determine the value of a given ſum payable on the extinction of the lives of A and B, ſhould they be the *firſt* that fail of the three lives A, B, and C.

SOLUTION.

The given ſum may be received in the firſt year, either on the extinction of the three lives, C having died laſt; or on the extinction of only the two lives A and B, C having ſurvived the year. In the ſecond and following years the given ſum may be received, provided either of ſix events ſhould happen. 1ſt, If the three lives ſhould become extinct in the year, C having been the laſt that failed. 2dly, If A only ſhould

ſhould die in the year, B having died before the beginning, and C lived to the end of it. 3dly, If B only ſhould die in the year, A having died before the beginning, and C lived to the end of it. 4thly, If A and B ſhould both die in the year, and C ſurvive it. 5thly, If C ſhould die after A in the year, B having died in either of the foregoing years. 6thly, if C ſhould die after B in the year, A having died in either of the foregoing years. The fractions denoting theſe ſeveral contingencies being added together will be found =

$$\frac{S}{2ac}\times\overline{\frac{a'c}{r}+\frac{a''\cdot d}{r^2}+\frac{a'''\cdot e}{r^3}+\&c.}+\frac{S}{2ac}\times\overline{\frac{a\,d}{r}+\frac{a''e}{r^2}+\frac{a'''\cdot f}{r^3}+\&c.}-$$

$$\frac{S}{6abc}\times\overline{\frac{a'bc}{r}+\frac{a''md}{r^2}+\frac{a'''ne}{r^3}+\&c.}+\frac{S}{2abcr}\times\overline{\frac{md\cdot a'}{r}+\frac{ne\cdot a'+a''}{r^2}+\&c.}-$$

$$\frac{S}{3abc}\times\overline{\frac{a'mc}{r}+\frac{a''nd}{r^2}+\frac{a'''oe}{r^3}+\&c.}-\frac{S}{2abcr}\times\overline{\frac{nd\cdot a'}{r}+\frac{oe\cdot a'+a''}{r^2}+\&c.}+$$

$$\frac{S}{6abc}\times\overline{\frac{a'bd}{r}+\frac{a''me}{r^2}+\frac{a'''nf}{r^3}+\&c.}+\frac{S}{2abcr}\times\overline{\frac{me\cdot a'}{r}+\frac{nf\cdot a'+a''}{r^2}+\&c.}-$$

$$\frac{2S}{3abc}\times\overline{\frac{a'md}{r}+\frac{a''ne}{r^2}+\frac{a'''of}{r^3}+\&c.}-\frac{S}{2abcr}\times\overline{\frac{ne\cdot a'}{r}+\frac{of\cdot a'+a''}{r^2}+\&c.}$$

The firſt two ſeries are = D; the third and fourth ſeries are = $\frac{\alpha}{3r}\times\overline{BC-ABC}-\frac{\beta\alpha\cdot\overline{FK-AFK}}{6\cdot bc}$; the fifth and ſixth ſeries are = $-\frac{\alpha\cdot\overline{BK-ABK}}{3c}-\frac{m\cdot\overline{FC-APC}}{6br}$; the ſeventh and eighth ſeries = $\frac{\beta\cdot\overline{FC-AFC}}{6b}+\frac{d\cdot\overline{BT-ABT}}{3cr}$; and the ninth and tenth ſeries = $\frac{md\cdot\overline{PT-APT}}{6bcr}-\frac{2\cdot\overline{BC-ABC}}{3}$. Hence the whole value of the reverſion will be = $D-\frac{S\cdot\alpha}{3c}\times\overline{\frac{\beta\cdot\overline{FK-AFK}}{2b}+\overline{BK-ABK}}+\frac{S\cdot\beta}{6b}\times\overline{FC-AFC}-\frac{2\cdot S\cdot r-1}{3r}\times\overline{BC-ABC}-\frac{S\cdot m}{6br}\times\overline{PC-APC}+\frac{S\cdot d}{3cr}\times\overline{BT-ABT}+\frac{m\cdot\overline{PT-APT}}{2b}$.

The preceding theorem expreſſes the value of the given ſum whether C be the oldeſt, or B one of the two lives to be ſurvived, and will therefore be ſufficient in all caſes. In order, however, ſtill more fully to prove this, let A be ſuppoſed the oldeſt life, and inſtead of $\overline{b-m}$, $\overline{m-n}$, $\overline{n-o}$, &c. and a', a'', a''', &c. let b', b'', b''', &c. and $\overline{a-s}$, $\overline{s-t}$, $\overline{t-u}$, &c. be ſubſtituted; then will the value of the reverſion be found =

$$\frac{S}{2bc}\times\overline{\frac{cb'}{r}+\frac{db''}{r^2}+\frac{eb'''}{r^3}+\&c.}+\frac{S}{2bc}\times\overline{\frac{db'}{r}+\frac{eb''}{r^2}+\frac{fb'''}{r^3}+\&c.}-$$

$$\frac{S}{6abc}\times\overline{\frac{acb'}{r}+\frac{dsb''}{r^2}+\frac{etb'''}{r^3}+\&c.}+\frac{S}{2abcr}\times\overline{\frac{dsb'}{r}+\frac{et\,.\,b'+b''}{r^2}+\&c.}-$$

$$\frac{S}{3abc}\times\overline{\frac{rsb'}{r}+\frac{drb''}{r^2}+\frac{eub'''}{r^3}+\&c.}-\frac{S}{2abcr}\times\overline{\frac{drb'}{r}+\frac{eu\,.\,b'+b''}{r^2}+\&c.}+$$

$$\frac{S}{6abc}\times\overline{\frac{adb'}{r}+\frac{aeb''}{r^2}+\frac{ftb'''}{r^3}+\&c.}+\frac{S}{2abcr}\times\overline{\frac{aeb'}{r}+\frac{tf\,.\,b'+b''}{r^2}+\&c.}-$$

$$\frac{2S}{3abc}\times\overline{\frac{dsb'}{r}+\frac{et\,.\,b''}{r^2}+\frac{fu\,.\,b'''}{r^3}+\&c.}-\frac{S}{2abcr}\times\overline{\frac{et\,.\,b'}{r}+\frac{fu\,.\,b'+b''}{r^2}+\&c.}$$

Let Σ denote the value of S on the contingency of C's ſurviving B, and the general rule deduced from the preceding ſeries will become $=\Sigma-\frac{S\,.\,a}{3}\times\overline{\frac{a\,.\,\overline{HK-HBK}}{2a}+\overline{AK-ABK}}+$

$$\frac{S\,.\,a}{6a}\times\overline{HC-HBC}-\frac{2S\,.\,r-1}{3r}\times\overline{AC-ABC}-\frac{S\,.\,s}{6ar}\times\overline{NC-NBC}+$$

$$\frac{S\,.\,d}{3cr}\times\overline{AT-ABT}+\frac{s\,.\,\overline{NT-NBT}}{2a}$$, which appears to be exactly the ſame rule with the foregoing, if the ſymbols of A and B be only exchanged for each other.

If the three lives be of the ſame age, both thoſe rules will ſeverally become $=S$ into $\frac{r-1}{2r}\times\overline{V-CC}-\frac{xx}{6cc}\times\overline{KK-CKK}-$

$$\frac{x}{6c}\times\overline{CK-CCK}-\frac{2\,.\,r-1}{3r}\times\overline{CC-CCC}+\frac{d}{6cr}\times\overline{CT-CCT}+\frac{dd}{6ccr}\times$$

$\overline{TT-CTT}$. The two expreſſions $-\frac{nn\cdot\overline{KK-CKK}}{6cc}+\frac{d}{6cr}\times$ $\overline{CT-CCT}$, by reſolving them into their reſpective ſeries, will be found $=\frac{1-CC}{6r}+\frac{d}{6cr}+\frac{dd}{6c^2r^3}+\frac{ef}{6.2.r^3}+\frac{fg}{6c^2r^4}+$ &c., and the two expreſſions $\frac{dd\cdot\overline{TT-CTT}}{6ccr}-\frac{n}{c}\times\overline{CK-CCK}=\frac{CC}{6r}-\frac{d}{6cr}-$ $\frac{dd}{6ccr^3}+\frac{ef}{6ccr^3}+\frac{fg}{6ccr^4}+$ &c.; hence the ſum of theſe four expreſſions will be $=-\frac{r-1\cdot V-CC}{6r}$, and conſequently the general rule in this caſe will be $=\frac{S\cdot r-1}{3r}\times\overline{V-3CC-2CCC}$, which is known to denote the true value from other principles *.

As a further proof of the accuracy of the preceding inveſtigation, it may not be improper to obſerve, that this rule may be immediately obtained from the different fractions which expreſs the value of the given ſum in each year. For in this caſe the value of S in the firſt year is $=\frac{S}{3}\times\overline{\frac{1}{r}-\frac{3d}{ccr}+\frac{2d^3}{c^3\cdot r}}$, in the ſecond year $=\frac{S}{3}\times\overline{\frac{2r^3}{c^3r^3}-\frac{3er}{ccr^3}-\frac{2d^3}{c^3r^3}+\frac{3dd}{ccr^3}}$, and ſo on in the other years, which expreſſions may be eaſily found $=$ $\frac{S\cdot r-1}{3r}\times\overline{V-3CC-2CCC}$. Q. E. D.

PROBLEM V.

To find the value of a given ſum, payable on the death of A, if his life ſhould be the *firſt* or *ſecond* that fails of the three lives A, B, and C.

* See my Paper in the Phil. Tranſ. Vol. LXXIX.

SOLU-

SOLUTION.

In the first year the payment of the given sum depends upon either of four events. 1st, That all the three lives shall become extinct, the life of A having been the first or second that failed. 2dly, That A and B shall both die, and C live to the end of the year. 3dly, That A and C shall both die, and B live to the end of the year. 4thly, That only A shall die, and B and C both live to the end of the year. In the second and following years the payment of the given sum will depend upon either of eight events happening. 1st, That all the three lives shall drop in the year, A having been the first or second that failed. 2dly, That C survives, and only A and B die in the year. 3dly, That B survives, and only A and C die in the year. 4thly, That both B and C survive, and A only dies in the year. 5thly, That A dies in the year, B having died before the beginning, and C lived to the end of it. 6thly, That A in like manner dies in the year, C having died before the beginning, and B lived to the end of it. 7thly, That C dies after A in the year, B's life having failed in either of the preceding years. 8thly, That B dies after A in the year, C's life having failed in either of the preceding years. The fractions denoting these several contingencies are $= \frac{S}{2ac} \times \overline{\frac{a'c}{r} + \frac{a''.d}{r^2} + \frac{a'''.e}{r^3} + \&c.} + \frac{S}{2ac} \times \overline{\frac{a'd}{r} + \frac{a''.e}{r^2} + \frac{a'''f}{r^3} + \&c.} + \frac{S}{2ab} \times \overline{\frac{a'b}{r} + \frac{a''.m}{r^2} + \frac{a'''n}{r^3} + \&c.} + \frac{S}{2ab} \times \overline{\frac{a'm}{r} + \frac{a''n}{r} + \frac{a'''o}{r^3} + \&c.} - \frac{S}{3abc} \times \overline{\frac{a'bc}{r} + \frac{a''md}{r} + \frac{a'''ne}{r^3} + \&c.} - \frac{S}{6abc} \times \overline{\frac{a'md}{r} + \frac{a''nd}{r^2} + \frac{a''oe}{r^3} + \&c.} - \frac{S}{6abc} \times \overline{\frac{a'be}{r} + \frac{a''me}{r^2} + \frac{a'''nf}{r^3} + \&c.} - \frac{S}{3abc} \times \overline{\frac{a'md}{r} + \frac{a'ne}{r^2} + \frac{a'''of}{r^3} + \&c.} = D + E - M.$

This

This general rule gives the true value whether the life of A be older or younger than both or either of the lives of B and C. When the three lives are of equal age, the value of S for the firſt year will be $= \frac{S}{r} \times \overline{\frac{2}{3} + \frac{dd}{3^3} - \frac{dd}{cc}}$, for the ſecond year $= \frac{S}{r^2} \times \overline{\frac{ee}{3e^3} - \frac{ee}{cc} - \frac{dd}{3^3} + \frac{dd}{cc}}$, and ſo on for the other years. Hence the whole value in this caſe will be $= \frac{S \cdot \overline{r-1}}{3r} \times \overline{2V - 3CC - CCC}$, which expreſſion may alſo be derived from the general rule juſt given above, or D + E − M.

The ſolution of this problem may alſo be obtained either from the firſt and ſecond, or from the third problems. In the one caſe the value of S is evidently equal to the *ſum* of the two values determined by the two firſt-mentioned problems, or D + E − 2M + M = D + E − M. And in the other caſe its value is equal to the *difference* between the abſolute value of the reverſion after A (= G) and its value depending upon the contingency that A ſhall be the *laſt* life that ſhall fail, which being = G + M − D + E by the third problem, it follows, that the general rule on this ſuppoſition will be alſo = D + E − M. Q. E. D.

PROBLEM VI.

To find the value of a given ſum payable on the death of A, ſhould his life be the *ſecond* or *third* that ſhall fail of the three lives A, B, and C.

SOLUTION.

The payment of the given ſum in the firſt year will depend upon the contingency of either of three events. 1ſt, That all

all the three lives ſhall become extinct, A having been the ſecond or third that has failed. 2dly, That A ſhall die after B, and C live to the end of the year. 3dly, That A ſhall die after C, and B live to the end of the year. In the ſecond and following years the given ſum will become payable, provided either of eight events ſhould happen. 1ſt, If all the three lives ſhould fail in the year, A having been the ſecond or third that died. 2dly, If A ſhould die after B in the year, and C live to the end of it. 3dly, If A ſhould die after C in the year, and B live to the end of it. 4thly, If A and C ſhould both die in the year, B having died before the beginning of it. 5thly, If A and B ſhould both die in the year, C having died before the beginning of it. 6thly, If A only ſhould die in the year, the lives of B and C having become extinct in either of the preceding years. 7thly, If A ſhould die in the year, B having died before the beginning, and C lived to the end of it. 8thly, if A ſhould die in the year, C having died before the beginning, and B lived to the end of it. Hence the whole value of the reverſion will be found $=$

$$\frac{S}{a} \times \frac{a'}{r} + \frac{a''}{r^2} + \frac{a'''}{r^3} + \&c. - \frac{S}{3abc} \times \overline{\frac{a'bc}{r} + \frac{a''md}{r^2} + \frac{a'''ne}{r^3} + \&c.} -$$

$$\frac{S}{6abc} \times \overline{\frac{a'mc}{r} + \frac{a''nd}{r^2} + \frac{a'''oe}{r^3} + \&c.} - \frac{S}{6abc} \times \overline{\frac{a'bd}{r} + \frac{a''me}{r^2} + \frac{a'''nf}{r^3} + \&c.} -$$

$$\frac{S}{3abc} \times \overline{\frac{a'md}{r} + \frac{a''ne}{r^2} + \frac{a'''of}{r^3} + \&c.} = G - M.$$

This rule is correct in all caſes; but when the three lives are equal it becomes more ſimple, and is $= \frac{S \cdot r-1}{3r} \times \overline{2V - 3C - CCC}$; which expreſſion may likewiſe be obtained immediately from the ſeries given above.

The ſolution of this problem, like that of the foregoing one, may alſo be derived from the firſt three problems; for the value

value of S is either equal to the *difference* between the abſolute value of the reverſion after the death of A and its value depending on the contingency that A ſhall be the *firſt* that fails (found by Prob. 1.), or it is equal to the *ſum* of the two values depending on the contingencies that A ſhall be the *ſecond* or *third* that fails (found by Prob. 2. and 3.). In both caſes the general rule is = G − M. Q. E. D.

PROBLEM VII.

To find the value of a given ſum payable on the death of A, ſhould his life be the *firſt* or *laſt* that fails of the three lives A, B, C.

SOLUTION.

In order to receive the given ſum in the firſt year, it is neceſſary that either of four events ſhould happen. 1ſt, That all the three lives ſhould fail, A having been the firſt or third that died. 2dly, That B ſhould die after A, and C live. 3dly, That C ſhould die after A, and B live. 4thly, That A only ſhould die, and B and C both live. In the ſecond and following years the given ſum may be received, provided either of ſeven events ſhould happen. 1ſt, If the three lives ſhould fail, A having been the firſt or laſt that died. 2dly, if B ſhould die after A in the year, and C live to the end of it. 3dly, If C ſhould die after A in the year, and B live to the end of it. 4thly, If A only ſhould die in the year, and B and C both live to the end of it. 5thly, If A's life ſhould fail after that of B in the year, C's life having failed before the beginning of it. 6thly, If A ſhould fail after C in the year, B having failed before the beginning of it. 7thly, If A only

ſhould die in the year, B and C having died in either of the preceding years. From the fractions denoting theſe ſeveral contingencies the whole value of the reverſion will be found $=$

$$\frac{S}{a}\times\overline{\frac{a'}{r}+\frac{a''}{r^2}+\frac{a'''}{r^3}+\&c.}-\frac{S}{2ac}\times\overline{\frac{a'c}{r}+\frac{a''d}{r^2}+\frac{a'''e}{r^3}+\&c.}-\frac{S}{2ac}\times\overline{\frac{a'd}{r}+\frac{a''e}{r^2}+\frac{a'''f}{r^3}+\&c.}-\frac{S}{2ab}\times\overline{\frac{a'b}{r}+\frac{a''m}{r^2}+\frac{a'''n}{r^3}+\&c.}-\frac{S}{2ab}\times\overline{\frac{a'm}{r}+\frac{a''n}{r^2}+\frac{a'''o}{r^3}+\&c.}+\frac{2S}{3abc}\times\overline{\frac{a'bc}{r}+\frac{a''md}{r^2}+\frac{a'''ne}{r^3}+\&c.}+\frac{S}{3abc}\times\overline{\frac{a'mc}{r}+\frac{a''nd}{r^2}+\frac{a'''oe}{r^3}+\&c.}+\frac{S}{3abc}\times\overline{\frac{a'bd}{r}+\frac{a''me}{r^2}+\frac{a'''nf}{r^3}+\&c.}+\frac{2S}{3abc}\times\overline{\frac{a'md}{r}+\frac{a''ne}{r^2}+\frac{a'''ef}{r^3}+\&c.}=G-\overline{D+E}+2M.$$

This general theorem will give the exact value in all caſes; but when the lives are equal, it is rendered more ſimple, by ſubſtituting the ſeveral values of G, D, E, and M, and will then become $=\frac{S\cdot\overline{r-1}}{3r}\times\overline{2V-3C-3CC+2CCC}$; which expreſſion may alſo be derived in this particular caſe from the different ſeries denoting the value of S in each year.

The ſolution of this, like thoſe of the two preceding problems, may likewiſe be obtained by the aſſiſtance of the firſt three problems. For the value of this contingent reverſion is either equal to the *ſum* of the two values of S payable on the death of A, if his life ſhould be the firſt, or if it ſhould be the laſt that fails (found by Prob. 1. and 3.), or it is equal to the *difference* between the value of the abſolute reverſion after A's death, and the value of the contingent reverſion after A's death, provided he ſhould be the *ſecond* that fails of the three lives (found by Prob. 2.). In both caſes the general rule becomes $=G-\overline{D+E}+2M$. Q. E. D.

PRO

PROBLEM VIII.

To find the value of a given ſum payable on the death of A or B, ſhould *either* of them be the *firſt* that ſhall fail of the three lives A, B, and C.

SOLUTION.

In each year the payment of the given ſum will depend upon either of ſix events. 1ſt, If the three lives ſhould fail in the year, A or B having died firſt. 2dly, if A and B ſhould die in the year, and A live. 3dly, If C ſhould die after A in the year, and B live. 4thly, If C ſhould die after B in the year, and A live. 5thly, If A only ſhould die in the year, and B and C both live. 6thly, If B only ſhould die in the year, and A and C both live. The fractions denoting theſe ſeveral contingencies are $= \frac{2S}{3abc} \times \overline{\frac{bca'}{r} + \frac{md \cdot a''}{r^2} + \frac{nea'''}{r^3} + \&c.} - \overline{\frac{S}{6abc} \times \frac{mca'}{r} + \frac{nda''}{r^2} + \frac{oea'''}{r^3} + \&c.} + \overline{\frac{S}{3abc} \times \frac{bda'}{r} + \frac{mea''}{r^2} + \frac{nf \cdot a'''}{r^3} + \&c.} + \overline{\frac{S}{6abc} \times \frac{mda'}{a} + \frac{nea''}{r^2} + \frac{ofa'''}{r^3} + \&c.} + \overline{\frac{S}{2bc} \times \frac{b-m.c}{r} + \frac{m-n.d}{r^2} + \frac{n-o.e}{r^3}} + \&c. + \overline{\frac{S}{2bc} \times \frac{b-m.d}{r} + \frac{m-n.e}{r^2} + \frac{n-o.f}{r^3} + \&c.} - \overline{\frac{S}{2abc} \times \frac{bca'}{r} + \frac{md.a'+a''}{r^2}} + \&c. + \overline{\frac{S}{2abc} \times \frac{mca'}{r} + \frac{nd.a'+a''}{r^2}} + \&c. - \overline{\frac{S}{2abc} \times \frac{bda'}{r} + \frac{me.a'+a''}{r^2}} + \&c. + \overline{\frac{S}{2abc} \times \frac{md.a'}{r} + \frac{ne.a'+a''}{r^2}} + \&c. = S \text{ into } \frac{x}{3c} \times \overline{\frac{\beta \cdot FK - AFK}{2b}} + \overline{BK - ABK} - \overline{\frac{\beta \cdot FC - AFC}{6b}} + \overline{\frac{2 \cdot r-1 \cdot BC - ABC}{3r}} + \overline{\frac{m \cdot PC - APC}{6br}} - \frac{d}{3er} \times \overline{BT - ABT} + \overline{\frac{m \cdot PT - APT}{2b}} + \Sigma$ (Σ denoting the value of S on the contingency of C's ſurviving B, as in Prob. 4.)

The above rule gives the exact value when C is the oldeſt of the three lives. But if A be the oldeſt, the ſymbols muſt be changed as in ſome of the foregoing problems, and the value in this caſe will be expreſſed by the ſeries $\frac{2S}{3abc} \times \overline{\frac{abc'}{r} + \frac{msc''}{r^2} + \frac{mtc'''}{r^3} + \&c.} - \frac{S}{6abc} \times \overline{\frac{amc'}{r} + \frac{nsc''}{r^2} + \frac{otc'''}{r^3} + \&c.} - \frac{S}{6abc} \times \overline{\frac{bsc'}{r} + \frac{mtc''}{r^2} + \frac{nuc'''}{r^3} + \&c.} - \frac{S}{3abc} \times \overline{\frac{msc'}{r} + \frac{ntc''}{r^2} + \frac{ouc'''}{r^3} + \&c.} + \frac{S}{ab} \times \overline{\frac{ab}{r} + \frac{ms}{r^2} + \frac{nt}{r^3} + \&c.} - \frac{S}{ab} \times \overline{\frac{ms}{r} + \frac{nt}{r^2} + \frac{ou}{r^3} + \&c.} - \frac{S}{abc} \times \overline{\frac{abc'}{r} + \frac{ms \cdot c' + c''}{r^2} + \&c.} + \frac{S}{abc} \times \overline{\frac{mtc'}{r} + \frac{nt \cdot c' + c''}{r^2} + \&c.}$ From theſe ſeveral ſeries, the general rule expreſſing the value of the reverſion will be found = S into $\frac{r-1 \cdot \overline{V-AB}}{r} - \frac{a}{3a} \times \frac{\beta \cdot \overline{HF-HFC}}{b} + \frac{\overline{HB-HBC}}{2} - \frac{\beta \cdot \overline{AF-AFC}}{6b} + \frac{2 \cdot \overline{r-1} \cdot \overline{AB-ABC}}{3r} + \frac{m \cdot \overline{AP-APC}}{6br} + \frac{S}{3ar} \times \overline{\frac{BN-BNC}{2} + \frac{m \cdot \overline{PN-PNC}}{b}}$.

When the lives are all equal, the firſt rule becomes = $\frac{\overline{r-1} \cdot \overline{V-CC}}{2r} + \frac{2 \cdot \overline{r-1} \cdot \overline{CC-CCC}}{3r} + \frac{\alpha\alpha \cdot \overline{KK-CKK}}{6cc} + \frac{\alpha \cdot \overline{CK-CCK}}{6c} - \frac{d \cdot \overline{CT-CCT}}{6cr} - \frac{dd \cdot \overline{TT-CTT}}{6ccr}$, and the ſecond rule = $\frac{\overline{r-1} \cdot \overline{V-CC}}{r} + \frac{2 \cdot \overline{r-1} \cdot \overline{CC-CCC}}{3r} - \frac{\alpha\alpha \cdot \overline{KK-CKK}}{3cc} - \frac{\alpha \cdot \overline{KC-CCK}}{3c} + \frac{d \cdot \overline{CT-CCT}}{3cr} + \frac{dd \cdot \overline{TT-CTT}}{3ccr}$. In the one caſe the four laſt fractions are = $\frac{\overline{r-1} \cdot \overline{V-CC}}{6r}$; and in the other caſe thoſe fractions are = $-\frac{\overline{r-1} \cdot \overline{V-CC}}{3r}$; therefore, in both caſes the general rule becomes = $\frac{2 \cdot S \cdot \overline{r-1}}{3r} \times \overline{V-CCC}$, which is known to be the true value from

from ſelf-evident principles. This expreſſion may alſo be immediately derived from the different fractions which denote the value of S in each year. For in the firſt year theſe fractions will be reduced to $\frac{2S}{3} \times \frac{1}{r} - \frac{d^3}{c^3 r}$, in the ſecond year to $\frac{2S}{3} \times \frac{d^3 - c^3}{c^3 r^2}$, in the third year to $\frac{2S}{3} \times \frac{c^3}{c^3 r^3} - \frac{r^3}{c^3 r^3}$, and ſo on in the other years. Hence the whole value of the reverſion will, as above, be $= \frac{2 \cdot S \cdot r - 1}{3r} \times \overline{V - CCC}$. Q. E. D *.

PROBLEM IX.

To determine the value of a given ſum payable on the death of A or B, ſhould *either* of them be the *ſecond* that fails of the three lives A, B, and C.

SOLUTION.

The payment of the given ſum in the firſt year will depend upon either of four events happening. 1ſt, That the three lives ſhould fail, A or B having been the ſecond that failed. 2dly, That A ſhould die after C, and B live. 3dly, That B ſhould die after C, and A live. 4thly, That A and B ſhould both die, and C live. In the ſecond and following years the given ſum will become payable, provided either of eleven events ſhould happen. 1ſt, If the three lives ſhould drop in the year, A or B having been the ſecond that failed. 2dly, If A ſhould die after C in the year, and B live. 3dly, If B ſhould die after C in the year, and A live. 4thly, If A and B

* I do not know that any ſolution has been attempted before, either of this or of the two following problems.

ſhould

ſhould both die in the year, and C live. 5thly, if B only ſhould die in the year, A having died before the beginning, and C lived to the end of it. 6thly, If A only ſhould die in the year, B having died before the beginning, and C lived to the end of it. 7thly, If C ſhould die after A in the year, B having died in either of the foregoing years. 8thly, If C ſhould die after B in the year, A having died in either of the foregoing years. 9thly, If A and B ſhould both die in the year, C having died in either of the preceding years. 10thly, If B only ſhould die in the year, C having died before the beginning, and A lived to the end of it. And, laſtly, if A only ſhould die in the year, C having died before the beginning, and B lived to the end of it. The ſeveral fractions denoting theſe contingencies in each year being added together will become $= \frac{S}{abc} \times \overline{\frac{bca'}{r} + \frac{md \cdot a' + a''}{r^2} + \&c.} - \frac{S}{abc} \times \overline{\frac{mca'}{r} + \frac{nd \cdot a' + a''}{r^2} + \&c.}$

$+ \frac{S}{abc} \times \overline{\frac{bda'}{r} + \frac{me \cdot a' + a''}{r^2} + \&c.} - \frac{S}{abc} \times \overline{\frac{mda'}{r} + \frac{ne \cdot a' + a''}{r^2} + \&c.} -$

$\frac{4S}{3abc} \times \overline{\frac{bca'}{r} + \frac{mda''}{r^2} + \frac{nea'''}{r^3} + \&c.} + \frac{S}{3abc} \times \overline{\frac{mca'}{r} + \frac{nda''}{r^2} + \frac{oea'''}{r^3} + \&c.} -$

$\frac{2S}{3abc} \times \overline{\frac{bda'}{r} + \frac{mea''}{r^2} + \frac{nfa'''}{r^3} + \&c.} - \frac{S}{3abc} \times \overline{\frac{mda'}{r} + \frac{nea''}{r^2} + \frac{of \cdot a'''}{r^3} + \&c.} -$

$\frac{S}{2bc} \times \overline{\frac{b-m \cdot c}{r} + \frac{m-n \cdot d}{r^2} + \frac{n-o \cdot e}{r^3} + \&c.} - \frac{S}{2bc} \times \overline{\frac{b-m \cdot d}{r} + \frac{m-n \cdot e}{r^2} + \frac{n-o \cdot f}{r^3}}$

$\overline{+ \&c.} + \frac{S}{ab} \times \overline{\frac{ma'}{r} + \frac{n \cdot a' + a''}{r^2} + \&c.} - \frac{S}{ab} \times \overline{\frac{b \cdot a'}{r} + \frac{m \cdot a' + a''}{r^2} + \&c.} +$

$\frac{S}{ab} \times \overline{\frac{ba'}{r} + \frac{ma''}{r^2} + \frac{na''}{r^3} + \&c.} + \frac{S}{b} \times \overline{\frac{b-m \cdot}{r} + \frac{m-n \cdot}{r^2} + \frac{n-o}{r^3} + \&c.} +$

$\frac{S}{2ac} \times \overline{\frac{ca'}{r} + \frac{da''}{r^2} + \frac{ea'''}{r^3} + \&c.} + \frac{S}{2ac} \times \overline{\frac{da'}{r} + \frac{ea''}{r^2} + \frac{fa'''}{r^3} + \&c.}$ Let Q denote the value of S by the firſt rule in the eighth problem, and the firſt ten ſeries will be $= \Sigma - 2Q$. The four next ſeries

are $= \frac{S \cdot \overline{r-1} \cdot \overline{V-AB}}{r}$; and the two laſt ſeries are $= D$; the whole value of the reverſion, therefore (when C is the oldeſt of the three lives), is $= \frac{S \cdot \overline{r-1} \cdot \overline{V-AB}}{r} + D + \Sigma - 2Q$.

When A is the oldeſt of the three lives, the different ſeries, by changing the ſymbols, as in ſome of the foregoing problems, will become $= \frac{2S}{abc} \times \overline{\frac{abc'}{r} + \frac{ms \cdot \overline{c'+c''}}{r^2}} + \&c. - \frac{2S}{abc} \times \overline{\frac{msc'}{r} + \frac{nt \cdot \overline{c'+c''}}{r^2}} + \&c. - \frac{4S}{3abc} \times \overline{\frac{abc'}{r} + \frac{msc''}{r^2} + \frac{nt \cdot c'''}{r^3}} + \&c. + \frac{2S}{3abc} \times \overline{\frac{ms \cdot c'}{r} + \frac{nt \cdot c''}{r^2} + \frac{ou \cdot c'''}{r^3}} + \&c. + \frac{S}{3abc} \times \overline{\frac{amc'}{r} + \frac{nsc''}{r^2} + \frac{otc'''}{r^3}} + \&c. + \frac{S}{3abc} \times \overline{\frac{bsc'}{r} + \frac{mtc''}{r^2} + \frac{nuc'''}{r^3}} + \&c. - \frac{S}{bc} \times \overline{\frac{bc'}{r} + \frac{m \cdot \overline{c'+c''}}{r^2}} + \&c. + \frac{S}{bc} \times \overline{\frac{mc'}{r} + \frac{n \cdot \overline{c'+c''}}{r^2}} + \&c. + \frac{S}{2bc} \times \overline{\frac{bc'}{r} + \frac{mc''}{r^2} + \frac{nc'''}{r^3}} + \&c. - \frac{S}{2bc} \times \overline{\frac{mc'}{r} + \frac{nc''}{r^2} + \frac{oc''}{r^3}} + \&c. - \frac{S}{ac} \times \overline{\frac{ac'}{r} + \frac{s \cdot \overline{c'+c''}}{r^2}} + \&c. + \frac{S}{ac} \times \overline{\frac{sc'}{r} + \frac{t \cdot \overline{c'+c''}}{r^2}} + \&c. + \frac{S}{2ac} \times \overline{\frac{ac'}{r} + \frac{sc''}{r^2} + \frac{tc'''}{r^3}} + \&c. - \frac{S}{2ac} \times \overline{\frac{sc'}{r} + \frac{tc''}{r^2} + \frac{uc'''}{r^3}} + \&c. + \frac{S}{b} \times \overline{\frac{b-m}{r} + \frac{m-n}{r^2} + \frac{n-o}{r^3}} + \&c. - \frac{S}{ab} \times \overline{\frac{b-m \cdot s}{r} + \frac{m-n \cdot t}{r^2} + \frac{n-o}{r^3} \cdot u} + \&c. + \frac{S}{abr} \times \overline{\frac{b-m \cdot s}{r} + \frac{m-n \cdot t}{r^2}} + \&c.$ Let Q repreſent the value of S by the ſecond rule in the eighth problem, Π and Δ the values of the ſame ſum on the contingency of B's dying after C, and on the contingency of A's dying after C reſpectively *; then will the general rule in this caſe become $= \frac{S \cdot \overline{r-1} \cdot \overline{3V-B-A-AB}}{r} - 2Q - \overline{\Pi + \Delta}$, or, becauſe $D + \Sigma$ are $= \frac{\overline{r-1} \cdot \overline{2V-B-A}}{r} - \overline{\Pi + \Delta}$ †,

* See my 3d Prob. in Phil. Tranſ. Vol. LXXVIII.

† See my 2d and 3d Prob. Vol. LXXVIII.

it

it will be $= \frac{S \cdot \overline{r-1} \cdot \overline{V-AB}}{r} + D + E - 2Q$, as above.

When the lives are equal, the rule may be found either immediately from the ſeries, or from the foregoing expreſſions, $= \frac{2S \cdot \overline{r-1}}{3r} \times \overline{V - 3CC - 2CCC}$. Q. E. D.

PROBLEM X.

To find the value of a given ſum payable on the deceaſe of B or C, ſhould *either* of them be the *laſt* that fails of the three lives A, B, and C.

SOLUTION.

The ſum S can be received in the firſt year only on the extinction of the three lives, reſtrained to the contingency of A's life having been the firſt or ſecond that failed. In the ſecond and following years it may be received provided either of ſix events ſhould happen. 1ſt, If the three lives ſhould fail in the year, A having been the firſt or ſecond that died. 2dly, If B and C ſhould both die in the year, A having died before the beginning of it. 3dly, If C only ſhould die in the year, A and B having died in either of the preceding years. 4thly, If B only ſhould die in the year, A and C having died in either of the preceding years. 5thly, If B ſhould die after A in the year, C having died before the beginning of it. 6thly, If C ſhould die after A in the year, B having died before the beginning of it. From the fractions expreſſing theſe ſeveral contingencies, the whole value of the reverſion will be found $=$

$$\frac{2S}{3abc} \times \overline{\frac{mda'}{r} + \frac{ne \cdot a' + a''}{r^2} + \&c.} - \frac{S}{3abc} \times \overline{\frac{bca'}{r} + \frac{md \cdot a' + a''}{r^2} + \&c.} - \frac{S}{6abc}$$

$\frac{S}{6abc} \times \overline{\frac{mca'}{r} + \frac{nda''}{r^2} + \frac{oea'''}{r^3} + \&c.} - \frac{S}{6abc} \times \overline{\frac{bda'}{r} + \frac{mea''}{r^2} + \frac{nf \cdot a'''}{r^3} + \&c.} +$
$\frac{S}{2ab} \times \overline{\frac{ba'}{r} + \frac{m \cdot a' + a''}{r^2} + \frac{n \cdot a' + a'' + a'''}{r^3} + \&c.} - \frac{S}{2ab} \times \overline{\frac{ma'}{r} + \frac{n \cdot a' + a''}{r^2} +$
$\&c.} + \frac{S}{2ac} \times \overline{\frac{ca'}{r} + \frac{d \cdot a' + a''}{r^2} + \&c.} - \frac{S}{2ac} \times \overline{\frac{da'}{r} + \frac{e \cdot a' + a''}{r^2} + \&c.} -$
$\frac{2S}{3abcr} \times \overline{\frac{mda'}{r} + \frac{ne \cdot a' + a''}{r^2} + \&c.} + \frac{S}{3abcr} \times \overline{\frac{nea'}{r} + \frac{of \cdot a' + a''}{r^2} + \&c.} +$
$\frac{S}{2abr} \times \overline{\frac{ma'}{r} + \frac{n \cdot a' + a''}{r^2} + \&c.} - \frac{S}{2abr} \times \overline{\frac{na'}{r} + \frac{o \cdot a' + a''}{r^2} + \&c.} +$
$\frac{S}{2acr} \times \overline{\frac{d \cdot a'}{r} + \frac{e \cdot a' + a''}{r^2} + \&c.} - \frac{S}{2acr} \times \overline{\frac{ea'}{r} + \frac{f \cdot a' + a''}{r^2} + \&c.}$, and

therefore the general rule (when either B or C are the oldeſt of the three lives) will be $= S$ into $\frac{2 \cdot r-1 \cdot BC-ABC}{3r} - \frac{x}{3c} \times$
$\frac{\beta \cdot FK-AFK}{b} + \frac{BK-ABK}{2} \times \frac{m}{3br} \times \frac{PC-APC}{2} + \frac{d \cdot PT-APT}{c} - \frac{\beta \cdot PC-APC}{6b}$
$+ \frac{d \cdot BT-ABT}{6cr} + \frac{\beta \cdot F-AF}{2b} - \frac{r-1 \cdot B-AB}{2r} - \frac{m \cdot P-AP}{2br} + \frac{x \cdot K-AK}{2c} -$
$\frac{r-1 \cdot C-AC}{2r} - \frac{d \cdot T-AT}{2cr}$; or (ſince the firſt five expreſſions are $=$
$\frac{r-1 \cdot BC-ABC}{r} - M$, and the ſix remaining ones $= R + W$) it will be $= \frac{S \cdot r-1 \cdot BC-ABC}{r} + R + W - M$.

When A *is the oldeſt of the three lives*, the ſame general rule may be obtained. In this caſe, by exchanging the ſymbols, the different ſeries will become $= \frac{S}{6abc} \times \overline{\frac{abc'}{r} + \frac{ms \cdot c' + c''}{r^2} + \&c.} -$
$\frac{S}{6abc} \times \overline{\frac{bs \cdot c'}{r} + \frac{nt \cdot c' + c''}{r^2} + \&c.} + \frac{S}{3abc} \times \overline{\frac{amc'}{r} + \frac{ns \cdot c' + c''}{r^2} + \&c.} +$
$\frac{2S}{3abc} \times \overline{\frac{ms \cdot c'}{r} + \frac{nt \cdot c' + c''}{r^2} + \&c.} + \frac{S}{c} \times \overline{\frac{c'}{r} + \frac{c''}{r^2} + \frac{c'''}{r^3} + \&c.} -$

$\frac{S}{bc} \times \overline{\frac{mc'}{r} + \frac{n \cdot c + c'}{r^2} + \&c.} - \frac{S}{2ac} \times \overline{\frac{ac'}{r} + \frac{sc''}{r^2} + \frac{tc'''}{r^3} + \&c.} -$

$\frac{S}{2ac} \times \overline{\frac{sc'}{r} + \frac{t \cdot c''}{r^2} + \frac{uc'''}{r^3} + \&c.} - \frac{2S}{3abcr} \times \overline{\frac{ms \cdot c'}{r} + \frac{nt \cdot c + c''}{r^2} + \&c.} -$

$\frac{S}{3abr} \times \overline{\frac{mt \cdot c'}{r} + \frac{nu \cdot c + c''}{r^2} + \&c.} + \frac{S}{6abcr} \times \overline{\frac{2sc'}{r} + \frac{2t \cdot c + c''}{r^2} + \&c.} -$

$\frac{S}{6abcr} \times \overline{\frac{mt \cdot c'}{r} + \frac{nu \cdot c + c''}{r^2} + \&c.} + \frac{S}{bcr} \times \overline{\frac{mc'}{r} + \frac{n \cdot c + c''}{r^2} + \&c.} = S$ into

$\frac{2 \cdot r-1 \cdot AB-ABC}{3r} + \frac{\beta}{6b} \times \overline{\frac{\alpha \cdot HF-HFC}{a} - AF-AFC} + \frac{\alpha \cdot HB+HBC}{3a}$

$- \frac{s}{3ar} \times \overline{NB-NBC} + \frac{m \cdot PN-PNC}{2b} + \frac{m \cdot AP-APC}{6br} + \frac{r+1 \cdot V+C}{r} -$

$\frac{r-1 \cdot B-BC}{r} - \frac{r-1 \cdot A-AC}{2r} - \frac{\alpha \cdot H-HC}{2a} + \frac{s \cdot N-NC}{2ar}$. In order to get the same general rule as that given above in the case of B or C's being the oldest of the three lives, it is to be observed, that the first five expressions, by the second part of the solution of Prob. 1. appear to be $= \frac{r-1 \cdot AB \quad ABC}{r} - M + E$. And (supposing Γ to represent the value of S on the contingency of A's dying after C) that the last three expressions appear to be $= -\Gamma - \frac{r-1 \cdot A-AC}{r}$. But E is $= R + \frac{r-1 \cdot B-AB}{r}$, and W must be $= \frac{r-1 \cdot V-A-C+AC}{r} - \Gamma$; therefore, the sum of the above expressions may be easily found $= \frac{S \cdot r-1 \cdot BC-ABC}{r} + R + W - M$.

If the three lives be of equal age, the value of the reversion will be $= \frac{2S \cdot r-1}{3r} \times \overline{V-L}$. This expression may be derived either from the foregoing general rule, or immediately from

from the different series, and is known to be accurately true from self-evident principles. Q. E. D.

I have now given general rules for determining the values of reversions depending upon three lives in every case which, as far as I can discover, will admit of an *exact* solution. The remaining cases, which are nearly equal in number to those I have investigated, involve a contingency for which it appears very difficult to find such a general expression as shall not render the rules much too complicated and laborious. The contingency to which I refer is that of *one life's failing after another in any given time*. The fractions expressing this probability are every year increasing, so that the value of the reversion must be represented by as many series at least as are equal to the difference between the age of one of the lives, and that of the oldest life in the table of observations. I have indeed so far succeeded in the method of approximation as that the reversion may be generally ascertained within about $\frac{1}{10}$th part of its exact value; but I shall not trouble the Royal Society at present with these investigations.

The 34th, 35th, and 36th problems in Mr. SIMPSON's Select Exercises involve this contingency, and, by the assistance of M. DE MOIVRE's hypothesis, admit of an easy solution. But such is the fallacy of this hypothesis, that it renders Mr. SIMPSON's conclusions obviously wrong, though his reasoning is perfectly correct: for it cannot surely be an *equal* chance *in all cases* that one life shall die after another. In the short term of a single year the chances are indeed so nearly equal, that it would be wrong to perplex the solution by attempting

greater accuracy. But when the number of years, and the difference between the ages of the two lives, are considerable, those chances must vary in proportion; and, therefore, unless the contingency is blended with another which shall very much diminish the probability of the event, the solution, by thus indiscriminately supposing the chances to be equal, must be rendered extremely inaccurate. In Mr. SIMPSON's 36th problem the solution by this means appears to be absurd: for, in the particular case in which C is the oldest of the three lives, the value of the reversionary annuity becomes $= \frac{C-AC}{2}$; that is, the value of an annuity in this case during the life of C after B and A, provided A dies first, is the same whatever be the age of B; for no mention is made of his life in the foregoing expression. It should be observed, however, that the rule itself is strictly true, and that the error arises from Mr. SIMPSON's having been misled by the hypothesis in determining the probability of B's dying after A in his investigation of the 34th problem, which is applied to the solution of this problem *.

I have declined giving specimens of the different values of the reversions as deduced from the foregoing rules and those which have been hitherto published, not only from an apprehension of becoming tedious, but also from the conviction that at present they are unnecessary; those which I have formerly given being, I think, sufficient to prove the inaccuracy of M. DE MOIVRE's hypothesis. In those instances in which I have compared some of the foregoing rules with the approxi-

* It is proper to observe, that I have followed Mr. SIMPSON's method of determining this contingency in the 23d, 27th, 28th, and 29th Problems in my Treatise on Annuities.

mations now in uſe, I have invariably found the latter to be erroneous; nay, in ſome caſes, the values were almoſt twice as great as they ought to have been. This is particularly true when one of the lives is very young, and both or either of the other lives are very old. In reverſions of this kind I believe that this is generally the caſe, and that it ſeldom happens that the ages of the three lives are nearly equal. The approximations therefore can hardly ever be uſed with ſafety, and it will certainly be moſt prudent not to have recourſe to them when the correct values can be obtained. Should the difficulties attending the ſolution of the remaining problems which involve three lives be ſurmounted (and the taſk may not perhaps be impoſſible), the hypotheſis of an equal decrement of life, as far as it relates to any uſeful purpoſe in the doctrine of annuities, may then be totally abandoned. Or ſhould it even be found impracticable to deduce ſolutions of thoſe problems which are ſtrictly and accurately true; yet, I am ſatisfied from my own experience that ſuch near approximations may be procured as to render this hypotheſis equally unneceſſary.

XVIII. *Abstract of a Register of the Barometer, Thermometer, and Rain, at* Lyndon *in* Rutland; *by* Thomas Barker, *Esq.*; *with the Rain in* Surrey *and* Hampshire; *for the Year* 1790. *Communicated by* Thomas White, *Esq.* F.R.S.

Read June 2, 1791.

		Barometer.			Thermometer.						Rain.			
					In the House.			Abroad.			Lyndon	Surrey. S. Lambeth	Hampshire. Selbourn.	Hampshire. Fyfield.
		Highest.	Lowest.	Mean.	High.	Low.	Mean	High.	Low.	Mean				
		Inches.	Inches.	Inches.	°	°	°	°	°	°	Inch.	Inch.	Inch.	Inch.
Jan.	Morn.	29,97	28,48	29,55	49	36	41	50½	26½	37	1,871	1,49	1,99	1,72
	Aftern.				49	36	41½	52½	34½	41				
Feb.	Morn.	30,05	29,20	29,71	48	40	43½	48	30	38½	0,236	0,20	0,49	0,43
	Aftern.				49½	41	44½	57	40½	47				
Mar.	Morn.	30,13	29,31	29,77	50½	42½	46	48½	31	39½	0,259	0,24	0,45	0,38
	Aftern.				51	44	47	60	42½	50				
Apr.	Morn.	29,85	29,01	29,42	51	40	44	49½	32	38½	0,676	2,54	3,64	1,27
	Aftern.				52½	41	45	61	38	49				
May	Morn.	29,80	28,93	29,45	58	50½	54	57	43½	49½	2,911	3,70	4,38	3,66
	Aftern.				59½	51	55	72	52½	61				
June	Morn.	29,86	28,98	29,56	71	54½	59	65½	48½	55½	2,385	0,64	0,13	0,55
	Aftern.				75½	55½	60½	85	56	67½				
July	Morn.	29,73	29,90	29,35	64	57	60	62	49½	57	2,246	2,42	3,24	1,71
	Aftern.				66	58	61	78½	62	69				
Aug.	Morn.	29,68	29,17	29,49	66	56	60½	62½	49	56	1,735	2,26	2,30	1,97
	Aftern.				67½	57	62	80	57	69				
Sept.	Morn.	29,95	28,88	29,53	62	52	55½	59½	42	49½	1,566	0,52	0,66	0,62
	Aftern.				61	53	57	72½	55½	61				
Oct.	Morn.	29,81	28,89	29,43	57	45½	52	55	36	45½	0,991	1,72	2,10	1,25
	Aftern.				59	45½	53	65	46	55				
Nov.	Morn.	29,88	28,49	29,35	49½	37½	44	48	30	40	3,145	3,40	6,95	5,11
	Aftern.				49½	38	45	51½	32	44½				
Dec.	Morn.	29,87	28,32	29,33	46	36	40	48	25½	38½	3,608	3,18	5,94	3,38
	Aftern.				47	36	40½	49	31	43				
										Inches	21,629	22,31	32,27	22,05

The

The thermometer abroad being broken in December, that month is defective in that article.

THE winter proved a remarkably open one; no froſt of two whole days together, nor any ſnow; ſo that there were young artichokes in the garden all winter, where they were ſhort and among the leaves; and the ſecond crop of figs, which are generally killed in winter, eſcaped this year, and ripened in the ſummer, but did not grow large. It was ſaid there were young hares in winter, and birds' eggs laid, ſome of which were perhaps hatched. Several people ſaid they ſaw a covey of young partridges near Oakham in February; but I cannot affirm it, for others ſuſpected ſome miſtake. January was ſhowery and mild, but often windy; as the days grew longer, it became dryer, finer, and forward. A ſwarm of bees roſe and ſettled at Tixover, March 2.; but it was rather a ſhifting of their hive for want, than a regular ſwarm, which could not have been bred ſo early. The owner, having a deſerted hive with combs and honey in it, put them into that, where they lived, and grew ſtrong in the ſummer. The forwardneſs of the ſeaſon continued till the N.E. winds of March made it colder and more froſty; the growth of all things was then ſtopped, and the firſt fortnight in April was ſeverer than any part of the winter, with the only ſnow this ſeaſon.

The ſeed time was dry, and in general good; but, for want of froſt in the winter, the ground ploughed ſtiff, and required a great deal of working; and the ſeaſon continued dry and backward till after the middle of May, when rain and warmer weather made every thing grow as faſt as they were ſlow before. The ſummer was chiefly cloudy and windy, and after

the middle of May very ſhowery and cool, except ſome few violent hot days; and a very growing ſeaſon, ſo that grounds laid at the end of May had good crops of hay on them, which in general was well got, for the rains in this country were not great, though frequent; and ſo much land was laid late, and ſo much over-eaten graſs cut, that there was hardly ever more good hay made than there was this year. The harveſt alſo was ſhowery, yet not ſo much but that thoſe who were not too haſty got it well, and it was plentiful, though very rank, and pretty much laid, yielding well to the acre, though not to the flail.

The autumn was dryer, fine, and pleaſant, ſo that the ground began to burn. The wheat ſeed time was remarkably good; for it was dry at firſt for ſowing the clay land; then rain came in good time to bring it up, and that made the ground moiſt for ſowing the dry fields. There was a good deal of rain at times in November, yet ſo much fine weather between, that it was pleaſant and not dirty; but moſt part of December and January was uncommonly ſtormy and uncertain weather, changing ſometimes two or three times a day; ſevere ſtorms, much rain, great floods, and remarkable thunder, on ſeveral different days, and in many different places. Near a fortnight's froſt at Chriſtmas, with clear weather at firſt, but afterward one great rain, and ſeveral leſſer, which yet did not take away the froſt till January the 6th.

Chalk

Chalk found in a new place.

There is a great deal of chalky ground in the ſouthern part of England; I think it begins at the ſea in Devonſhire, and one vein of it runs all along the ſouthern counties to Dover. Another vein parts off from that about Reading in Berkſhire, goes by Dunſtable, Baldock, and Gogmagog Hills, and ſo on to the ſea in Norfolk; the whole croſſing the kingdom in a Y. Along theſe two diſtricts it is almoſt all chalk to a great depth in the ground; but out of them chalk is ſeldom found. I believe it may be met with in many places in the countries between theſe two diſtricts, and ſometimes deep in the ground, where it does not come up to the ſurface; but beyond the northern limits of them, which are at Wantage in Berkſhire, and over the river from Shillingford in Oxfordſhire, and at Maddingley by Cambridge, chalk is hardly any where to be found; no where in any conſiderable quantity, unleſs it be much farther north, in the wolds of Yorkſhire, beyond Pocklington toward Scarborough.

I did not know till lately that we had any chalk nearer us than Maddingley; but ſeveral years ago, the people of Ridlington in Rutland, digging for ſtone to mend the roads, met with a bed of chalk; at which they were much ſurprized, and did not know what it was, having never ſeen a chalk pit before. After I had heard of it, I went to examine the place, and found it a regular chalk pit, with rows of flints lying in it as is uſual in the ſouth of England. The chalk is not ſoft like that they write with, but very much like that they dig about Baldock; nor are the flints ſo black as thoſe in the ſouth of England, but veined, of a light-coloured flint, and white,

ſome parts much mixed with chalk; and are broken, not whole ones. They may have dug the pit ſix yards long and two deep; but how far the chalk reaches I do not know. The ground about it has plainly been formerly dug, perhaps thirty yards ſquare, but completely turfed over again, with the ſame ſtrong turf as the reſt of the cloſe, which is rich paſture land, and feeds oxen for Smithfield market, not like the ſhort graſs on the chalky downs.

Riding laſt autumn along the turnpike road near Stukeley in Huntingdonſhire, I ſaw a little patch of chalk, a few yards long, in a bank which had been dug away by the road ſide; ſo that though we did not know there was any chalk at all in this country, and there certainly is very little, yet here are now two places where it has been met with.

XIX. *Deſcription of a ſimple Micrometer for meaſuring ſmall Angles with the Teleſcope.* By *Mr.* Tiberius Cavallo, *F. R. S.*

Read June 2, 1791.

THE various teleſcopical micrometers, or machines which have been conſtructed for the meaſurement of ſmall angles, may be divided into two claſſes; namely, thoſe which have not, and thoſe which have, ſome movement amongſt their parts. The micrometers of the former ſort conſiſt moſtly of fine wires, or hairs, variouſly diſpoſed, and ſituated within the teleſcope, juſt where the image of the object is formed. In order to determine an angle with thoſe micrometers, a good deal of calculation is generally required. The micrometers of the other ſort, of which there is a great variety; ſome being made with moveable parallel wires, others with priſms, others again with a combination of lenſes, and ſo on; are more or leſs ſubject to ſeveral inconveniences, the principal of which are the following. 1ſt, Their motions generally depend upon the action of a ſcrew, and of courſe the imperfections of its threads, and the greater or leſs quantity of loſt motion, which is obſervable in moving a ſcrew, eſpecially when ſmall, occaſion a conſiderable error in the menſuration of angles. 2dly, Their complication and bulk renders them difficultly applicable to a variety of teleſcopes, eſpecially to the

pocket ones. 3dly, They do not measure the angle without some loss of time, which is necessary to turn the screw, or to move some other mechanism. 4thly, and lastly, They are considerably expensive, so that some of them cost even more than a tolerably good telescope.

After having had long in view the construction of a micrometer, which might be in part at least, if not intirely, free from all those objections; and, after various attempts, I at last succeeded with a simple contrivance, which, after repeated trials, has been found to answer the desired end, not only from my own experience, but from that also of several friends, to whom it has been communicated.

This micrometer, in short, consists of a thin and narrow slip of mother of pearl finely divided, and situated in the focus of the eye-glass of a telescope, just where the image of the object is formed. It is immaterial whether the telescope be a refractor or a reflector, provided the eye-glass be a convex lens, and not a concave one, as in the Galilean construction.

The simplest way of fixing it is to stick it upon the diaphragm, which generally stands within the tube, and in the focus of the eye-glass. When thus fixed, if you look through the eye-glass, the divisions of the micrometrical scale will appear very distinct, unless the diaphragm is not exactly in the focus; in which case the micrometrical scale must be placed exactly in the focus of the eye-glass, either by pushing the diaphragm backwards or forwards, when that is practicable; or else the scale may be easily removed from one or the other surface of the diaphragm by the interposition of a circular piece of paper or card, or by a bit of wax. This construction is fully sufficient when the telescope is always to be used by the same person; but when different persons are to use it, then

the diaphragm, which ſupports the micrometer, muſt be conſtructed ſo as to be eaſily moved backwards or forwards, though that motion needs not be greater than about a tenth or an eighth of an inch. This is neceſſary, becauſe the diſtance of the focus of the ſame lens appears different to the eyes of different perſons, and therefore, whoever is going to uſe the teleſcope for the menſuration of any angle, muſt firſt of all unſcrew the tube, which contains the eye-glaſs and micrometer, from the reſt of the teleſcope, and, looking through the eye-glaſs, muſt place the micrometer where the diviſions of it may appear quite diſtinct to his eye.

In caſe that any perſon ſhould not like to ſee always the micrometer in the field of the teleſcope, then the micrometrical ſcale, inſtead of being fixed to the diaphragm, may be fitted to a circular perforated plate of braſs, wood, or even paper, which may be occaſionally placed upon the ſaid diaphragm.

I have made ſeveral experiments to determine the moſt uſeful ſubſtance for this micrometer. Glaſs, which I had ſucceſsfully applied for a ſimilar purpoſe to the compound microſcope, ſeemed at firſt to be the moſt promiſing; but it was at laſt rejected after ſeveral trials: for the diviſions upon it generally are either too fine to be perceived, or too rough; and though with proper care and attention the diviſions may be proportioned to the ſight, yet the thickneſs of the glaſs itſelf obſtructs in ſome meaſure the diſtinct view of the object. Ivory, horn, and wood, were found uſeleſs for the conſtruction of this micrometer, on account of their bending, ſwelling, and contracting very eaſily; whereas mother of pearl is a very ſteady ſubſtance, the diviſions upon it may be marked very

very eaſily, and, when it is made as thin as common writing paper, it has a very uſeful degree of tranſparency.

Tab. VIII. fig. 1. exhibits this micrometer ſcale, but ſhews it four times larger than the real ſize of one, which I have adapted to a three-feet achromatic teleſcope, that magnifies about 84 times. It is ſomething leſs than the 24th part of an inch broad; its thickneſs is equal to that of common writing paper; and the length of it is determined by the aperture of the diaphragm, which limits the field of the teleſcope. The diviſions upon it are the 200dths of an inch, which reach from one edge of the ſcale to about the middle of it, excepting every fifth and tenth diviſion, which are longer. The divided edge of it paſſes through the center of the field of view, though this is not a neceſſary precaution in the conſtruction of this micrometer. Two diviſions of the above deſcribed ſcale in my teleſcope are very nearly equal to one minute; and as a quarter of one of thoſe diviſions may be very well diſtinguiſhed by eſtimation, therefore an angle of one-eighth part of a minute, or of $7''\frac{1}{2}$, may be meaſured with it.

When a teleſcope magnifies more, the diviſions of the micrometer muſt be more minute; and I find, that when the focus of the eye-glaſs of the teleſcope is ſhorter than half an inch, the micrometer may be divided with the 500dths of an inch; by means of which, and the teleſcope magnifying about 200 times, one may eaſily and accurately meaſure an angle ſmaller than half a ſecond.

On the other hand, when the teleſcope does not magnify above 30 times, the diviſions need not be ſo minute: for inſtance, in one of DOLLOND's pocket teleſcopes, which when drawn out for uſe, is about 14 inches long, a micrometer with the

the hundredths of an inch is quite ſufficient, and one of its diviſions is equal to little leſs than three minutes, ſo that an angle of a minute may be meaſured by it.

In looking through a teleſcope, furniſhed with ſuch a micrometer, the field of view appears divided by the micrometer ſcale, the breadth of which occupies about one-ſeventh part of the aperture, and as the ſcale is ſemitranſparent, that part of the object, which happens to be behind it, may be diſcerned ſufficiently well to aſcertain the diviſion, and even the quarter of a diviſion, with which its borders coincide. Fig. 2. ſhews the appearance of the field of my teleſcope with the micrometer, when directed to the title page of the Philoſophical Tranſactions, wherein one may obſerve that the thickneſs of the letter C is equal to three-fourths of a diviſion, the diameter of the O is equal to three diviſions, and ſo on.

At firſt view one is apt to imagine that it is difficult to count the diviſions which may happen to cover or to meaſure an object; but upon trial it will be found, that this is readily performed; and even people, who have never been uſed to obſerve with the teleſcope, ſoon learn to meaſure very quickly and accurately with this micrometer; for, ſince every fifth and tenth diviſion is longer than the reſt, one ſoon acquires the habit of ſaying five, ten, fifteen, and then, by adding the other diviſions leſs than five, completes the reckoning. Even with a teleſcope, which has no ſtand, if the object end of it be reſted againſt a ſteady place, and the other end be held by the hand near the eye of the obſerver, an object may be meaſured with accuracy ſufficient for ſeveral purpoſes, as for the eſtimation of ſmall diſtances, for determining the height of a houſe, &c.

After

After having conſtructed and adapted this micrometer to the teleſcope, it is then neceſſary to aſcertain the value of the diviſions. It is hardly neceſſary to mention in this place, that though thoſe diviſions meaſure the chords of the angles, and not the angles or arches themſelves, and the chords are not as the arches, yet it has been ſhewn by all the trigonometrical writers, that in ſmall angles the chords, arches, ſines, and tangents, follow the ſame proportion ſo very nearly, that the very minute difference may be ſafely neglected: ſo that if one diviſion of this micrometer is equal to one minute, we may ſafely conclude, that two diviſions are equal to two minutes, three diviſions to three minutes, and ſo on. There are various methods of aſcertaining the value of the diviſions of ſuch a micrometer, they being the very ſame that are uſed for aſcertaining the value of the diviſions in other micrometers. Such are the paſſage of an equatorial ſtar over a certain number of diviſions in a certain time, the meaſuring of the diameter of the ſun, by computation from the focal diſtance of the object, and other lenſes of the teleſcope, the laſt of which, however, is ſubject to ſeveral inaccuracies; but as they are well known to aſtronomical perſons, and have been deſcribed in many books, need not be farther noticed in this Paper. However, for the ſake of workmen and other perſons not converſant in aſtronomy, I ſhall deſcribe an eaſy and accurate method of aſcertaining the value of the diviſions of the micrometer.

Mark upon a wall, or other place, the length of ſix inches, which may be done by making two dots or lines ſix inches aſunder, or by fixing a ſix-inch ruler upon a ſtand; then place the teleſcope before it ſo that the ruler or ſix-inch length may be at right angles with the direction of the teleſcope, and juſt 57 feet 3¼ inches diſtant from the object-glaſs of the teleſcope: this done, look through the teleſcope at the ruler or other

extenſion

extenſion of ſix inches, and obſerve how many diviſions of the micrometer are equal to it, and that ſame number of diviſions is equal to half a degree, or 30′; and this is all that needs be done for the required determination; the reaſon of which is, becauſe an extenſion of ſix inches ſubtends an angle of 30′ at the diſtance of 57 feet 3½ inches, as may be eaſily calculated by the rules of plane trigonometry.

In one of DOLLOND's 14-inch pocket teleſcopes, if the diviſions of the micrometer be the hundredths of an inch, 11½ of thoſe diviſions will be found equal to 30′, or 23 to a degree.

When this value has been once aſcertained, any other angle meaſured by any other number of diviſions is determined by the rule of three. Thus, ſuppoſe that the diameter of the ſun, ſeen through the ſame teleſcope, be found equal to 12 diviſions, ſay as 11½ diviſions are to 30 minutes, ſo are 12 diviſions to $\left(\frac{12' \times 30'}{11,5}\right)$ 31′,3, which is the required diameter of the ſun.

Notwithſtanding the facility of this calculation, a ſcale may be made anſwering to the diviſions of a micrometer, which will ſhew the angle correſponding to any number of diviſions to mere inſpection. Thus, for the above-mentioned ſmall teleſcope the ſcale is repreſented in fig. 3. AB is a line drawn at pleaſure; it is then divided into 23 equal parts, and thoſe diviſions, which repreſent the diviſions of the micrometer that are equal to one degree, are marked on one ſide of it. The line then is divided again into 60 equal parts, which are marked on the other ſide of it; and theſe diviſions repreſent the minutes which correſpond to the diviſions of the micrometer: thus the figure ſhews, that ſix diviſions of the micrometer are equal to 15½ minutes, 11½ diviſions are nearly equal to 29

minutes, &c. What has been ſaid of minutes may be ſaid of ſeconds alſo, when the ſcale is to be applied to a large teleſcope.

Thus far this micrometer, and its general uſe, have been ſufficiently deſcribed, and mathematical perſons may eaſily apply it to the various purpoſes to which micrometers have been found ſubſervient. But as the ſimplicity, cheapneſs, and at the ſame time the accuracy of this contrivance, may render the uſe of it much more general than that of any other micrometer; and I may venture to ſay, that it will be found very uſeful in the army, and amongſt ſea-faring people, for the determination of diſtances, heights, &c.; I ſhall therefore ſubjoin ſome practical rules to render this micrometer uſeful to perſons unacquainted with trigonometry and the uſe of logarithms.

Problem I. The angle, not exceeding one degree, which is ſubtended by an extenſion of one foot being given, to find its diſtance from the place of obſervation.

N. B. This extenſion of one foot, or any other which may be mentioned hereafter, muſt be perpendicular to the direction of the teleſcope through which it is obſerved. The diſtances are reckoned from the object-glaſs of the teleſcope, and the anſwers obtained by the rules of this problem, though not exactly true, are however ſo little different from the truth, that the difference ſeldom amounts to more than two or three inches, which may be ſafely neglected.

Rule 1. If the angle be expreſſed in minutes, ſay, as the given angle is to 60, ſo is 687,55 to a fourth proportional, which gives the anſwer in inches.

2. If the angle be expreſſed in ſeconds, ſay, as the given angle is to 3600, ſo is 687,55 to a fourth proportional, which expreſſes the anſwer in inches.

3. If the angle be expreſſed in minutes and ſeconds, turn it all into ſeconds, and proceed as above.

Example. At what diſtance is a globe of one foot in diameter when it ſubtends an angle of two ſeconds?

$2 : 3600 :: 687.55 : \frac{3600 \times 687.55}{2} = 1237590$ inches, or $103132\frac{1}{2}$ feet, which is the anſwer required.

This calculation may be ſhortened; for ſince two of the three proportionals are fixed, their product in the firſt caſe is 41253, and in the other two caſes is 2475180; ſo that in the firſt caſe, *viz.* when the angle is expreſſed in minutes, you need only divide 41253 by the given angle; and in the other two caſes, *viz.* when the angle is expreſſed in ſeconds, divide 2475180 by the given angle, and the quotient in either caſe is the anſwer in inches.

Problem II. The angle, not exceeding one degree, which is ſubtended by any known extenſion, being given, to find its diſtance from the place of obſervation.

Rule. Proceed as if the extenſion were of one foot by Problem I. and call the anſwer B; then, if the extenſion in queſtion be expreſſed in inches, ſay, as 12 inches are to that extenſion, ſo is B to a fourth proportional, which is the anſwer in inches; but if the extenſion in queſtion be expreſſed in feet, then you need only multiply it by B, and the product is the anſwer in inches.

Example. At what diſtance is a man, ſix feet high, when he appears to ſubtend an angle of 30″.

By Problem I. if the man were one foot high, the diſtance would be 82506 inches; but as he is ſix feet high, therefore multiply 82506 by 6, and the product gives the required diſtance, which is 495036 inches, or 41253 feet.

For greater conveniency, eſpecially in travelling, or in ſuch circumſtances in which one has not the opportunity of making even the eaſy calculations required in thoſe problems, I have calculated the following two tables; the firſt of which ſhews the diſtance anſwering to any angle from one minute to one degree, which is ſubtended by an extenſion of one foot; and the ſecond table ſhews the diſtance anſwering to any angle from one minute to one degree, which is ſubtended by a man, the height of which has been called an extenſion of ſix feet; becauſe, at a mean, ſuch is the height of a man when dreſſed with hat and ſhoes on. Theſe tables may be tranſcribed on a card, and may be had always ready with a pocket teleſcope furniſhed with a micrometer. Their uſe is evidently to aſcertain diſtances without any calculation; and they are calculated only to minutes, becauſe with a pocket teleſcope and micrometer it is not poſſible to meaſure an angle more accurately than to a minute.

Thus, if one wants to meaſure the extenſion of a ſtreet, let a foot ruler be placed at the end of the ſtreet; meaſure the angular appearance of it, which ſuppoſe to be 36′, and in the table you will have the required diſtance againſt 36′, which is 95½ feet. Thus alſo a man, who appears to be 49′ high, is at the diſtance of 421 feet.

T. CAVALLO.

Wells-ſtreet,
May 26, 1791.

Angles

Angles ſubtended by an extenſion of one foot at different diſtances.

Angles.	Diſtances in feet.	Angles.	Diſtances in feet.
Min. 1	3437,7	Min. 31	110,9
2	1718,9	32	107,4
3	1145,9	33	104,2
4	859,4	34	101,1
5	687,5	35	98,2
6	572,9	36	95,5
7	491,1	37	92,9
8	429,7	38	90,4
9	382,0	39	88,1
10	343,7	40	85,9
11	312,5	41	83,8
12	286,5	42	81,8
13	264,4	43	79,9
14	245,5	44	78,1
15	229,2	45	76,4
16	214,8	46	74,7
17	202,2	47	73,1
18	191,0	48	71,6
19	180,9	49	70,1
20	171,8	50	68,7
21	163,7	51	67,4
22	156,2	52	66,1
23	149,4	53	64,8
24	143,2	54	63,6
25	137,5	55	62,5
26	132,2	56	61,4
27	127,3	57	60,3
28	122,7	58	59,2
29	118,5	59	58,2
30	114,6	60	57,3

163,7

92 Div. = 6 in. 257,3

Angles

Angles subtended by an extenſion of ſix feet at different diſtances.

Angles.	Diſtances in feet.	Angles.	Diſtances in feet.
Min. 1	20626,8	Min. 31	665,4
2	10313.	32	644,5
3	6875,4	33	625.
4	5156,5	34	606,6
5	4125,2	35	589,3
6	3437,7	36	572,9
7	2946,6	37	557,5
8	2578,2	38	542,8
9	2291,8	39	528,9
10	2062,6	40	515,6
11	1875,2	41	503,1
12	1718,8	42	491,1
13	1586,7	43	479,7
14	1473,3	44	468,8
15	1375.	45	458,4
16	1289,1	46	448,4
17	1213,3	47	438,9
18	1145,9	48	429,7
19	1085,6	49	421.
20	1031,4	50	412,5
21	982,2	51	404,4
22	937,6	52	396,7
23	896,8	53	389,2
24	859,4	54	381,9
25	825.	55	375.
26	793,3	56	368,3
27	763,9	57	361,9
28	736,6	58	355,6
29	711,3	59	349,6
30	687,5	60	343,7

Fig. 2.

Fig. 1.

Fig. 3.

Cavallo del.

Basire sc.

XX. *A new Method of inveſtigating the Sums of Infinite Series.* By the *Rev.* Samuel Vince, *A. M. F. R. S.*

Read June 2, 1791.

THE ſummation of infinite ſeries is a ſubject, not only of curious ſpeculation, but alſo of the greateſt importance in the various branches of mathematics and philoſophy; in conſequence of which it has always claimed a very conſiderable ſhare of attention from the moſt celebrated mathematicians. I ſhall therefore make no apology for offering to the public the following new and very expeditious method, by which we may obtain the ſums of a great variety of ſeries, moſt of which have never before been treated of. As the ſummation depends on the ſums of the reciprocals of the powers of the natural numbers, tables of ſuch ſums are given as far as the 40th power to twelve places of decimals, by which the ſums of the ſeries will be found true to ten or eleven places; and if greater accuracy were required (which is a caſe that can very rarely happen) it might eaſily be obtained by continuing the tables. The firſt and third columns ſhew the ſums, and the ſecond and fourth the powers correſponding.

TABLE

TABLE I.

Sum of $\frac{1}{2^n}+\frac{1}{3^n}+\frac{1}{4^n}+$ &c. *ad infinitum.*

Sum	n =	Sum	n =
A =,644934066848	2	W =,000000238450	22
B =,202056903159	3	X =,000000119219	23
C =,082323233711	4	Y =,000000059608	24
D =,036927755107	5	Z =,000000029803	25
E =,017343061984	6	A′ =,000000014901	26
F =,008349277387	7	B′ =,000000007450	27
G =,004077356198	8	C′ =,000000003725	28
H =,002008392826	9	D′ =,000000001863	29
I =,000994575128	10	E′ =,000000000931	30
K =,000494188604	11	F′ =,000000000465	31
L =,000246086553	12	G′ =,000000000233	32
M =,000122713347	13	H′ =,000000000116	33
N =,000061248135	14	I′ =,000000000058	34
O =,000030588236	15	K′ =,000000000029	35
P =,000015282259	16	L′ =,000000000015	36
Q =,000007637196	17	M′ =,000000000007	37
R =,000003817292	18	N′ =,000000000004	38
S =,000001908212	19	O′ =,000000000002	39
T =,000000953961	20	P′ =,000000000001	40
V =,000000476932	21		

TABLE

TABLE II.

Sum of $\frac{1}{2^n} - \frac{1}{3^n} + \frac{1}{4^n} - \frac{1}{5^n} +$ &c. *ad infinitum.*

Sum	$\cong$	Sum	$\cong$
a =,177532966576	2	w =,000000238386	22
b =,098457322630	3	x =,000000119199	23
c =,052967170503	4	y =,000000059602	24
d =,027880229587	5	z =,000000029801	25
e =,014448908703	6	a' =,000000014901	26
f =,007406180072	7	b' =,000000007450	27
g =,003766998147	8	c' =,000000003725	28
h =,001905702459	9	d' =,000000001863	29
i =,000960492403	10	e' =,000000000931	30
k =,000482856502	11	f' =,000000000465	31
l =,000242314856	12	g' =,000000000233	32
m =,000121457237	13	h' =,000000000116	33
n =,000060829654	14	i' =,000000000058	34
o =,000030448787	15	k' =,000000000029	35
p =,000015235790	16	l' =,000000000015	36
q =,000007621708	17	m' =,000000000007	37
r =,000003812130	18	n' =,000000000004	38
s =,000001906491	19	o' =,000000000002	39
t =,000000953389	20	p' =,000000000001	40
v =,000000476742	21		

TABLE III.

Sum of $\frac{1}{2^n}+\frac{1}{4^n}+\frac{1}{6^n}+$&c. *ad infinitum.*

Sum	$\frac{n}{\|\|}$	Sum	$\frac{n}{\|\|}$
A″ =,411233516712	2	W″ =,000000238419	22
B″ =,102571128951	3	X″ =,000000119209	23
C″ =,067645202107	4	Y″ =,000000059605	24
D″ =,032403992347	5	Z″ =,000000029802	25
E″ =,015895985344	6	A‴ =,000000014901	26
F″ =,007877728730	7	B‴ =,000000007450	27
G″ =,003922177173	8	C‴ =,000000003725	28
H″ =,001957047643	9	D‴ =,000000001863	29
I″ =,000977533765	10	E‴ =,000000000931	30
K″ =,000488522553	11	F‴ =,000000000465	31
L″ =,000244200705	12	G‴ =,000000000233	32
M″ =,000122085292	13	H‴ =,000000000116	33
N″ =,000061038895	14	I‴ =,000000000058	34
O″ =,000030518512	15	K‴ =,000000000029	35
P″ =,000015259024	16	L‴ =,000000000015	36
Q″ =,000007629452	17	M‴ =,000000000007	37
R″ =,000003814712	18	N‴ =,000000000004	38
S″ =,000001907352	19	O‴ =,000000000002	39
T″ =,000000953675	20	P‴ =,000000000001	40
V″ =,000000476837	21		

TABLE

TABLE IV.

Sum of $\frac{1}{3^n}+\frac{1}{5^n}+\frac{1}{7^n}+$ &c. *ad infinitum.*

Sum	n	Sum	n
a'' =,233700550136	2	n'' =,000000209240	14
b'' =,051799790264	3	o'' =,000000069724	15
c'' =,014678031604	4	p'' =,000000023234	16
d'' =,004523762760	5	q'' =,000000007744	17
e'' =,001447076640	6	r'' =,000000002581	18
f'' =,000471548657	7	s'' =,000000000864	19
g'' =,000155179025	8	t'' =,000000000286	20
h'' =,000051345183	9	v'' =,000000000095	21
i'' =,000017041362	10	w'' =,000000000032	22
k'' =,000005666051	11	x'' =,000000000011	23
l'' =,000001885848	12	y'' =,000000000004	24
m'' =,000000628055	13	z'' =,000000000001	25

PROP. I.

To find the sum of the sums of the reciprocal squares, cubes, &c. &c. *ad infinitum.*

By division $\frac{1}{x-1\times x}=\frac{1}{x^2}+\frac{1}{x^3}+\frac{1}{x^4}+$&c. *ad inf.*; hence if we make each of these terms the general term of a series, and write 2, 3, 4, &c. *ad inf.* for x, we have $\frac{1}{1\cdot 2}+\frac{1}{2\cdot 3}+\frac{1}{3\cdot 4}+$ &c. = (Table 1.) A+B+C+D+&c.; but $\frac{1}{1\cdot 2}+\frac{1}{2\cdot 3}+\frac{1}{3\cdot 4}+$ &c. *ad inf.*=1; hence A+B+C+D+&c. *ad inf.* = 1.

As $\frac{1}{x \times x+1} = \frac{1}{x^2} - \frac{1}{x^3} + \frac{1}{x^4} - \frac{1}{x^5} +$ &c. *ad inf.*; we have, by the ſame method of proceeding, $A - B + C - D +$ &c. *ad inf.* $= \frac{1}{2}$; conſequently $A + C + E +$ &c. $= \frac{1}{2}$, and $B + D + F +$ &c. $= \frac{1}{4}$.

Becauſe $\frac{1}{x-1 \times x} = \frac{1}{x^2} + \frac{1}{x^3} + \frac{1}{x^4} +$ &c. *ad inf.*; if for x we write 2, 4, 6, &c., then will $\frac{1}{1 \cdot 2} + \frac{1}{3 \cdot 4} + \frac{1}{5 \cdot 6} +$ &c. = (Tab. 3) $A'' + B'' + C'' + D'' +$ &c.; but $\frac{1}{1 \cdot 2} + \frac{1}{3 \cdot 4} + \frac{1}{5 \cdot 6} +$ &c. = hyp. log. 2; hence $A'' + B'' + C'' + D'' +$ &c. = hyp. log. 2.

If in the ſame expreſſion we write 3, 5, 7, &c. for x, then $\frac{1}{2 \cdot 3} + \frac{1}{4 \cdot 5} + \frac{1}{6 \cdot 7} +$ &c. = (Tab. 4.) $a'' + b'' + c'' +$ &c.; but $\frac{1}{2 \cdot 3} + \frac{1}{4 \cdot 5} + \frac{1}{6 \cdot 7} +$ &c. = 1 − hyp. log. 2; hence $a'' + b'' + c'' +$ &c. = 1 − hyp. log. 2.—Hence from either of theſe two laſt caſes, we have a very expeditious method of finding the hyp. log. 2.

PROP. II.

To find the ſum of the infinite ſeries whoſe general term is $\frac{1}{mx^r \pm n}$.

By diviſion $\frac{1}{mx^r \pm n} = \frac{1}{mx^r} \mp \frac{n}{m^2x^{2r}} + \frac{n^2}{m^3x^{3r}} \mp \frac{n^3}{m^4x^{4r}} +$ &c. *ad inf.*; hence, if $\frac{1}{mx^r \pm n}$ be made the general term of a ſeries, and for x we write 2, 3, 4, &c., its ſum will be equal to the ſums of another ſet of ſerieſes, whoſe terms are the powers of the reciprocals of the natural numbers reſpectively multiplied into

into $\frac{1}{m}$, $\frac{n}{m}$, $\frac{n^2}{m^3}$, &c.; hence the ſum of each of theſe ſeries being known from the tables, the ſum of the given ſeries will be found.

Ex. 1. Let $\frac{1}{x^2+1}$ be the general term; now $\frac{1}{x^2+1} = \frac{1}{x^2} - \frac{1}{x^4} + \frac{1}{x^6} - \frac{1}{x^8} +$ &c.; hence if for x we write 2, 3, 4, &c. we have $\frac{1}{5} + \frac{1}{10} + \frac{1}{17} + \frac{1}{26} +$ &c. $=$ A $-$ C $+$ E $-$ G $+$ &c. $=$ (by Tab. 1.) ,576674037469.

Ex. 2. Let $\frac{1}{x^2-1}$ be the general term; then, by the ſame method of proceeding, $\frac{1}{3} + \frac{1}{8} + \frac{1}{15} + \frac{1}{24} +$ &c. $=$ A $+$ C $+$ E $+$ &c. $=$ (by Prop. 1.) $\frac{3}{4}$.

Cor. Becauſe $\frac{1}{8} + \frac{1}{24} + \frac{1}{48} +$ &c. $= \frac{1}{8} \times \overline{1 + \frac{1}{3} + \frac{1}{6} +}$ &c. $=$ (as $1 + \frac{1}{3} + \frac{1}{6} +$ &c. is the reciprocal of the figurative numbers of the ſecond order) $\frac{1}{8} \times 2 = \frac{1}{4}$; therefore $\frac{1}{3} + \frac{1}{15} + \frac{1}{35} +$ &c. $= \frac{1}{2}$. Alſo, as $\frac{1}{x^2-1} = \frac{1}{x^2} + \frac{1}{x^4} + \frac{1}{x^6} +$ &c.; if we write 2, 4, 6, &c. for x, we have $\frac{1}{3} + \frac{1}{15} + \frac{1}{35} +$ &c. $=$ (by Tab. 3.) A″ $+$ C″ $+$ E″ $+$ &c. $= \frac{1}{2}$; but, by Prop. 1. A″ $+$ B″ $+$ C″ $+$ D″ $+$ &c. $=$ hyp. log. 2; hence B″ $+$ D″ $+$ F″ $+$ &c. $= -\frac{1}{2} +$ hyp. log. 2.

Ex. 3. Let the general term be $\frac{1}{x^3-1} = \frac{1}{x^3} + \frac{1}{x^6} + \frac{1}{x^9} +$ &c., and, by writing 2, 3, 4, &c. for x, we have $\frac{1}{7} + \frac{1}{26} + \frac{1}{63} +$ &c. $=$ B $+$ E $+$ H $+$ &c. $=$,221689395104.

Ex.

Ex. 4. Let the general term be $\frac{1}{3x^4-2} = \frac{1}{3x^4} + \frac{2}{9x^8} + \frac{4}{27x^{12}} +$ &c., and, by writing 2, 3, 4, &c. for x, &c. we have $\frac{1}{46} + \frac{1}{241} + \frac{1}{766} +$ &c. $\frac{1}{3}C + \frac{2}{9}G + \frac{4}{27}L +$ &c. = ,028385252052.

Ex. 5. To find the ſum of the ſeries $\frac{1}{9} - \frac{1}{20} + \frac{1}{65} - \frac{1}{124} +$ &c. If we write 2, -3, 4, -5, &c. for x, the general term will be $\frac{1}{x^3+1} = \frac{1}{x^3} - \frac{1}{x^6} + \frac{1}{x^9} - \frac{1}{x^{12}} +$ &c. Now, by writing 2, -3, 4, -5, &c. for x, the ſerieſes of which $\frac{1}{x^3}$, $\frac{1}{x^9}$, &c. are the general terms, will be alternately $+$ and $-$, and therefore their ſums will be found in Tab. 2. and the ſerieſes of which $\frac{1}{x^6}$, $\frac{1}{x^{12}}$, &c. are the general terms will have their terms all $+$, and therefore their ſums will be found in Tab. 1. Hence the ſum required $= b + b + o +$ &c. $-E-L-R-$ &c. = ,082800993.803.

PROP. III.

To find the ſum of the ſums of the reciprocals of the odd *powers in* Tab. 2.

By diviſion $\frac{1}{x-1 \times x} = \frac{1}{x^2} + \frac{1}{x^3} + \frac{1}{x^4} + \frac{1}{x^5} +$ &c.; hence by writing 2, -3, 4, -5, &c. for x, the ſums of the ſerieſes of which $\frac{1}{x^3}$, $\frac{1}{x^5}$, &c. are the general terms, may be found by Tab. 2. and the other ſums by Tab. 1.; hence $\frac{1}{1.2} + \frac{1}{3.4} + \frac{1}{5.6} +$ &c. $= A + C + E +$ &c. $+ b + d + f +$ &c.; but $\frac{1}{1.2} + \frac{1}{3.4} + \frac{1}{5.6} +$ &c.

&c. $= -\frac{1}{2} + 2$ hyp. log. 2; and by Prop. 1. $A+C+E+$&c. $= \frac{3}{4}$; hence $b+d+f+$&c. $= -\frac{5}{4} + 2$ hyp. log. 2.

PROP. IV.

To find the sum of the infinite series whose general term is $\frac{x^s}{mx^r \pm n}$.

By division $\frac{x^s}{mx^r \pm n} = \frac{1}{mx^{r-s}} \mp \frac{n}{m^2 x^{2r-s}} + \frac{n^2}{m^3 x^{3r-s}} \mp$ &c. *ad inf.*; hence the sum of the series of which $\frac{x^s}{mx^r \pm n}$ is the general term, is found as in Prop. 2. Here r must be greater than s at least by 2, otherwise the sum will be infinite.

Ex. 1. Let the general term be $\frac{x^2}{x^5+1} = \frac{1}{x^3} - \frac{1}{x^6} + \frac{1}{x^{10}} -$ &c.; hence if for x we write 2, 3, 4, &c. we have $\frac{4}{17} + \frac{9}{82} + \frac{16}{257} +$ &c. $=$ A $-$ E $+$ I $-$ N $+$ &c. $=$,538527924723.—If for x we write 2, 4, 6, &c. we get $\frac{4}{17} + \frac{16}{257} + \frac{36}{1296} +$ &c. $=$ A″ $-$ E″ $+$ I″ $-$ N″ $+$ &c. $=$,396257616555.

Ex. 2. Let the general term be $\frac{x}{3x^3-1} = \frac{1}{3x^2} + \frac{1}{9x^5} + \frac{1}{27x^8} +$ &c.; hence if we write 2, 3, 4, &c. for x, we have $\frac{2}{23} + \frac{3}{80} + \frac{4}{191} +$ &c. $= \frac{1}{3}$A $+ \frac{1}{9}$D $+ \frac{1}{27}$G $+$ &c. $=$,219238483448.

By this proposition we may find the sum of any series whose general term is $\frac{ax^s + bx^{s-1} + cx^{s-2} +}{mx^r \pm n}$ &c.; for this resolves itself into $\frac{ax^s}{mx^r \pm n}$, $\frac{bx^{s-1}}{mx^r \pm n}$, &c. &c., the sum of each of which series is

found by this propoſition. Now the $\overline{s+1}$th differences of the numerators of this general term are $=0$, and therefore it comprehends all ſeries under ſuch circumſtances. For example, let the given ſeries be $\frac{4}{17}+\frac{13}{82}+\frac{26}{257}+\frac{43}{626}$. Here the third differences of the numerators $=0$; to find therefore the general expreſſion for the numerator, aſſume ax^2+bx+c for it; and, by writing 2, 3, 4, for x, we have $4a+2b+c=4$, $9a+3b+c=13$, $16a+4b+c=26$; hence $a=2$, $b=-1$, $c=-2$; and as the denominator is manifeſtly x^4+1, the general term will be $\frac{2x^2-x-2}{x^4+1}=\frac{2x^2}{x^4+1}-\frac{x}{x^4+1}-\frac{2}{x^4+1}$, each of which being made the general term of a ſeries, their ſum will be found to be reſpectively 1,077055849446, 0,194173022145, and 0,156955159332; hence the ſum of the given ſeries is 0,725927667969.

If s be negative, the general term becomes $\frac{1}{x^s\times mx^r\pm n}=\frac{1}{mx^{r+s}\mp\&c.}=\frac{1}{m^2x^{2r+s}}+\frac{1}{m^3x^{3r+s}}+\&c.$

Ex. 1. To find the ſum of $\frac{1}{1\cdot2\cdot3}-\frac{1}{2\cdot3\cdot4}+\frac{1}{3\cdot4\cdot5}-\&c.$ *ad inf.* Here the general term is $\frac{1}{x-1\times x\times x+1}=\frac{1}{x\times x^2-1}=\frac{1}{x^3}+\frac{1}{x^5}+\frac{1}{x^7}+\&c.$; hence, by writing 2, -3, 4, -5, &c. for x, we have the ſum $=b+d+f+\&c.=$ (by Prop. 3.) $-\frac{5}{4}+2$ hyp. log. 2.

If $\frac{1}{x-1\times x^3\times x+1}$ be the general term it reſolves itſelf into $\frac{1}{x^5}+\frac{1}{x^7}+\frac{1}{x^9}+\&c.$; conſequently the ſum of $\frac{1}{1\cdot2^3\cdot3}-\frac{1}{2\cdot3^3\cdot4}$

+

$+\frac{1}{3\cdot 4^3\cdot 5}$ – &c. $= -b-\frac{5}{4}+2$ hyp. log. 2. In like manner the sum of $\frac{1}{1\cdot 2^2\cdot 3}-\frac{1}{2\cdot 3^2\cdot 4}+\frac{1}{3\cdot 4^2\cdot 5}$ – &c. $= -b-d-\frac{5}{4}+$ 2 hyp. log. 2. Thus we may proceed as far as we please by adding two powers to the middle term; and hence this remarkable property of the series, that the difference of the sums of the series where the middle term is x, x^3, x^5, &c. is b, d, f, &c. respectively.

Ex. 2. In like manner if the general term be $\frac{1}{x-1\times x^3\times x+1}$, and we write 2, 3, 4, &c. for x, we have $\frac{1}{1\cdot 2^3\cdot 3}+\frac{1}{2\cdot 3^3\cdot 3}$ $+\frac{1}{3\cdot 4^3\cdot 5}+$ &c. $=$ D+F+H+&c. $=$ (by Prop. 1.) $\frac{1}{4}$ – B. Hence also $\frac{1}{1\cdot 2^3\cdot 3}+\frac{1}{2\cdot 3^3\cdot 4}+$ &c. $=\frac{1}{4}-$B – D; and so on as before.

If the general term be under the form $\frac{1}{x^n\cdot x+m}$, it will be most convenient to resolve it thus: by division $\frac{1}{x+m}=\frac{1}{x}-\frac{m}{x^2}+\frac{m^2}{x^3}-$ &c. $\pm\frac{m^n}{x^n\cdot x+m}$; hence $\pm\frac{1}{x^n\cdot x+m}=$ $\frac{1}{x+m}-\frac{1}{x}+\frac{m}{x^2}-\frac{m^2}{x^3}+$ &c. $\times\frac{1}{m^n}=-\frac{m}{x\cdot x+m}+\frac{m}{x^2}-\frac{m^2}{x^3}+$&c. $\times\frac{1}{m^n}$, where the sign on the left hand will be + or – according as n is even or odd, and the number of terms on the right is $=n$. Now the sum of the series whose general term is $\frac{m}{x\cdot x+m}$ is well known, and the sums of the other are found from the tables.

Ex. 1. To find the sum of $\frac{1}{2^2\cdot 3}+\frac{1}{3^2\cdot 4}+\frac{1}{4^2\cdot 5}+$ &c. *ad inf.*

Here the general term is $\frac{1}{x^2 \times x+1} = -\frac{1}{x \cdot x+1} + \frac{1}{x^2}$, and by writing 2, 3. 4, &c. for x, we have the sum $= -\frac{1}{2 \cdot 3} - \frac{1}{3 \cdot 4}$ - &c. $+ A = -\frac{1}{2} + A$. In like manner $\frac{1}{2^3 \cdot 3} + \frac{1}{4^3 \cdot 5} + \frac{1}{6 \cdot 7}$ + &c. $= -1 +$ hyp.log. $2 + A''$. Also $\frac{1}{2^3 \cdot 5} + \frac{1}{3^3 \cdot 6} + \frac{1}{4^3 \cdot 7}$ + &c. $= \frac{13}{12} - 3A + 9B \times \frac{1}{27}$.

If m be negative, then $\frac{1}{x^n \cdot x-m} = \overline{\frac{m}{x \cdot x-m} - \frac{m}{x^2} - \frac{m^2}{x^3}} -$ &c. $\times \frac{1}{m^n}$. Hence $\frac{1}{2^4 \cdot 1} + \frac{1}{3^4 \cdot 2} + \frac{1}{4^4 \cdot 3}$ + &c. $= 1 - A - B - C$; and so on for others of the same kind.

If the general term be under this form $\frac{1}{x^{rn} \cdot ax^n + m}$, then, in like manner, we have $\pm \frac{1}{x^{rn} \cdot ax^n + m} = \overline{\frac{1}{ax^n + m} - \frac{1}{ax^n} + \frac{m}{a^2 x^{2n}}} -$ &c. $\times \frac{a^r}{m^r}$, where the sign on the left hand will be + or −, according as r is even or odd, and the number of terms on the right is $= r + 1$.

Ex. 1. To find the sum of $\frac{1}{2^4 \cdot 5} + \frac{1}{3^4 \cdot 10} + \frac{1}{4^4 \cdot 17}$ + &c. Here $m = 1$, $n = 2$, $r = 2$, $a = 1$, and the general term $\frac{1}{x^4 \times x^2+1} = \frac{1}{x^2+1} - \frac{1}{x^2} + \frac{1}{x^4}$; now the sum of the series whose general term is $\frac{1}{x^2+1}$ is $= ,5766740374 69$. by Prop. 2.; consequently the sum required $= ,5766740374 69 - A + C = ,0140632043 32$.

Ex. 2. If the given series be $\frac{1}{4 \cdot 5} + \frac{1}{9 \cdot 10} + \frac{1}{16 \cdot 17}$ + &c. the general

general term will be $\frac{1}{x^2 \cdot x+1} = -\frac{1}{x^2+1} + \frac{1}{x^2}$; hence, by writing 2, 3, 4, &c. for x, we have the sum $= -,576674037469 + A = ,068260029 38$.

If m be negative, then $\frac{1}{x^{rn} \cdot ax^n - m} = \frac{1}{ax^n - m} - \frac{1}{ax^n} - \frac{m}{a^2 x^{2n}}$ - &c. $\times \frac{a^r}{m^r}$.

Ex. 1. To find the sum of $\frac{1}{1 \cdot 2^2 \cdot 3} + \frac{1}{2 \cdot 3^2 \cdot 4} + \frac{1}{3 \cdot 4^2 \cdot 5}$ + &c. Here the general term is $\frac{1}{x-1 \times x^2 \times x+1} = \frac{1}{x^2 \times x^2 - 1} = \frac{1}{x^2-1} - \frac{1}{x^2}$; now, by writing 2, 3, 4, &c. for x, the sum of the series whose general term is $\frac{1}{x^2-1}$ is $= \frac{3}{4}$, by Prop. 2.; hence the sum required $= \frac{3}{4} - A$.

Ex. 2. Let the given series be $\frac{1}{1 \cdot 2^2 \cdot 3} + \frac{1}{3 \cdot 4^2 \cdot 5} + \frac{1}{5 \cdot 6^2 \cdot 7} +$ &c. Here the general term is the same as before, writing 2, 4, 6, &c. for x; and, by Prop. 2. the sum of the series whose general term is $\frac{1}{x^2-1}$ is $= \frac{1}{2}$; hence the sum $= \frac{1}{2} - A''$.

Ex. 3. In like manner the sum of the series $\frac{1}{1 \cdot 2^2 \cdot 3} + \frac{1}{2 \cdot 3^2 \cdot 4} + \frac{1}{3 \cdot 4^2 \cdot 5}$ + &c. $= ,221689395104 - B$.

Ex. 4. To find the sum of $\frac{1}{3 \cdot 4^2 \cdot 5} + \frac{1}{8 \cdot 9^2 \cdot 10} + \frac{1}{15 \cdot 16^2 \cdot 17}$ + &c. Here the general term is $\frac{1}{x^2-1 \times x^4 \times x^2+1} = \frac{1}{x^4 \times x^4-1}$ $= \frac{1}{x^4-1} - \frac{1}{x^4}$; but the sum of the series whose general term is $\frac{1}{x^4-1}$ is $= ,086662976264$; hence the sum required $= ,086662976264 - C$.

PROP. V.

To find the sum of the infinite series $\frac{1}{15}+\frac{1}{40}+\frac{1}{85}+\frac{1}{156}+\frac{1}{259}+$ &c.

In this series the fourth differences of the denominators $=0$; therefore the general term must be represented by $\frac{1}{ax^3+bx^2+cx+d}$; write therefore 2, 3, 4, &c. for x, and we have $8a+4b+2c+d=15$, $27a+9b+3c+d=40$, $64a+16b+4c+d=85$, $125a+25b+5c+d=156$; hence $a=1$, $b=1$, $c=1$, $d=1$, and the general term is $\frac{1}{x^3+x^2+x+1}=\frac{1}{x^3}-\frac{1}{x^4}+\frac{1}{x^7}-\frac{1}{x^8}+$ &c.; hence the sum $=B-C+F-G+K-L+$ &c. $=$,1242700165.

PROP. VI.

To find the sum of $\frac{2}{5^2}+\frac{3}{10^2}+\frac{4}{17^2}+$ &c. *ad inf.*

The general term $=\frac{x}{x^3+1^2}=\frac{1}{x^3}-\frac{2}{x^5}-\frac{3}{x^7}+$ &c.; hence, by writing 2, 3, 4, &c. for x, we have the sum $=B-2D+3F-$ &c. $=$,147115771469.

In like manner $\frac{2}{3^2}+\frac{3}{8^2}+\frac{4}{15^2}+$ &c. $=B+2D+3F+$ &c. $=$,312498999865.

PROP.

PROP. VII.

To find the sum of $\frac{1}{3^2 \cdot 5^2} + \frac{1}{8^2 \cdot 10^2} + \frac{1}{15^2 \cdot 17^2}$ + &c. *ad inf.*

The general term is $\frac{1}{x^2-1^2 \times x^2+1^2} = \frac{1}{x^4} + \frac{2}{x^{12}} + \frac{3}{x^{16}}$ + &c.; hence, by writing 2, 3, 4, &c. for x, we have the sum = G + 2L + 3P + &c. =,009447690684.

PROP. VIII.

To find the sum of $\frac{1}{1^3 \cdot 2^3 \cdot 3^3} + \frac{1}{2^3 \cdot 3^3 \cdot 4^3} + \frac{1}{3^3 \cdot 4^3 \cdot 5^3}$ + &c. *ad inf.*

Here the general term is $\frac{1}{x-1^3 \cdot x^3 \cdot x+1^3} = \frac{1}{x^9} + \frac{3}{x^7} + \frac{6}{x^5}$ + &c. and hence the sum = H + 3K + 6M + &c. =,004707148337.

PROP. IX.

To find the sum of the infinite series $1 - \frac{1}{3} + \frac{1}{6} - \frac{1}{10}$ + &c. *being a series of the reciprocal of the figurative numbers of the* 3rd *order, having the signs alternately* + *and* −.

This series, by resolving two terms into one, becomes $\frac{4}{1 \cdot 2 \cdot 3} + \frac{4}{3 \cdot 4 \cdot 5} + \frac{4}{5 \cdot 6 \cdot 7}$ + &c. whose general term, by writing 2, 4, 6, &c. for x, is $\frac{4}{x-1 \times x \times x+1} = \frac{4}{x^3} + \frac{4}{x^5} + \frac{4}{x^7}$ + &c. consequently the sum = 4B″ + 4D″ + 4F″ + &c. = (by Cor. Ex. 2. Prop. 2.) − 2 + 4 hyp. log. 2.

Cor.

Cor. Hence, as $1+\frac{1}{3}+\frac{1}{6}+\frac{1}{10}+\&c.=2$, we have $1+\frac{1}{6}+\frac{1}{15}+\&c.=2$ hyp. log. 2, and $\frac{1}{3}+\frac{1}{10}+\frac{1}{24}+\&c.=2-2$ hyp. log. 2.

PROP. X.

To find the sum of the infinite series $1-\frac{1}{4}+\frac{1}{10}-\frac{1}{20}+\&c.$ *being the reciprocals of the figurative numbers of the 4th order, having the signs alternately* + *and* −.

If we write 2, −3, 4, −5, &c. for x, the general term will be $\frac{6}{x^3-x}=\frac{6}{x^3}+\frac{6}{x^5}+\frac{6}{x^7}+\&c.$; hence the sum required = $6b+6d+6f+\&c.=$ (by Prop. 3.) $-7\frac{1}{2}+12$ hyp. log. 2.

Cor. Becauſe the sum of $1+\frac{1}{4}+\frac{1}{10}+\frac{1}{20}+\&c.=\frac{3}{2}$; therefore $1+\frac{1}{10}+\frac{1}{35}+\&c.=-3+6$ hyp. log. 2; and $\frac{1}{4}+\frac{1}{10}+\frac{1}{56}+\&c.=4\frac{1}{2}-6$ hyp. log. 2.

PROP. XI.

To find the sum of $\frac{2^3}{1^2\cdot 3^2}+\frac{3^3}{2^2\cdot 4^2}+\frac{4^3}{3^2\cdot 5^2}+\&c.$ *ad infinitum.*

The general term, by writing 2, 3, 4, &c. for x, is $\frac{x^3}{x-1\times x+1}=\frac{1}{x^2}+\frac{2}{x^4}+\frac{3}{x^6}+\&c.$; hence the sum $=A+2C+3E+\&c.=,884966993407$.

PROP.

PROP. XII.

To find the sum of $\frac{1}{1\cdot 2^2\cdot 3^2}+\frac{1}{2\cdot 3^2\cdot 4^2}+\frac{1}{3\cdot 4^2\cdot 5^2}$ &c. *ad infinitum.*

Here the general term, by writing 2, 3, 4, &c. for x, is $\frac{1}{x-1\cdot x^2\cdot x+1}=\frac{1}{x^6}-\frac{2}{x^7}+\frac{4}{x^8}+\frac{6}{x^9}-\frac{9}{x^{10}}-\frac{12}{x^{11}}+\frac{16}{x^{12}}-$ &c.; consequently the sum $= E-2F+4G-6H+9I-12K$ + &c. = ,010370898482.

PROP. XIII.

To find the sum of $\frac{1}{2}A-\frac{1}{4}B+\frac{1}{6}C-$ &c. *ad infinitum.*

The hyp. log. $2=1-\frac{1}{2}+\frac{1}{3}-\frac{1}{4}+\frac{1}{5}-$ &c. $=1+\frac{1}{3}+\frac{1}{5}+$ &c. $-\frac{1}{2}-\frac{1}{4}-\frac{1}{6}-$ &c.; hence $2\times$ hyp. log. 2, or hyp. log. 4, $=\frac{2}{1}+\frac{2}{3}+\frac{2}{5}+$ &c. $-1-\frac{1}{2}-\frac{1}{3}-$ &c. Now, by division, $\frac{2}{2x+1}=\frac{1}{x}-\frac{1}{2x^2}+\frac{1}{4x^3}-\frac{1}{8x^4}+$ &c.; hence, by writing 2, 3, 4, &c. for x, we have (after transposition) $\frac{2}{5}+\frac{2}{7}+$ &c. $-\frac{1}{2}-\frac{1}{3}-\frac{1}{4}$ $-$ &c. $=-\frac{1}{2}A+\frac{1}{4}B-\frac{1}{6}C+$ &c.; hence, by adding equal quantities to each side, we have $\frac{2}{1}+\frac{2}{3}+\frac{2}{5}+$ &c. $-\frac{1}{2}-\frac{1}{3}-\frac{1}{4}-$ &c. $=\frac{5}{3}-\frac{1}{2}A+\frac{1}{4}B-\frac{1}{6}C+$ &c., consequently $\frac{1}{2}A-\frac{1}{4}B+\frac{1}{6}C-$ &c. $=\frac{5}{3}-\frac{2}{1}-\frac{2}{3}-\frac{2}{5}-$ &c. $+1+\frac{1}{2}+\frac{1}{3}+$ &c. $=\frac{5}{3}-$ hyp. log. 4.

PROP.

PROP. XIV.

To find the sum of the infinite series $\frac{1}{2\cdot 5}+\frac{1}{3\cdot 7}+\frac{1}{4\cdot 9}+$&c.

The general term, by writing 2, 3, 4, &c. for x, is $\frac{1}{x\cdot 2x+1}=\frac{1}{2x^2}-\frac{1}{4x^3}+\frac{1}{8x^4}-$&c.; hence the sum$=\frac{1}{2}A-\frac{1}{4}B+\frac{1}{8}$C $-$&c. $=$ (by Prop. 13.) $\frac{1}{2}-$hyp. log. 4.

PROP. XV.

To find the sum of $1+\frac{1}{2}+\frac{1}{3}+\ldots\ldots$ *to* $\frac{1}{x}$.

The hyp. log. $\frac{x}{x-1}=\frac{1}{x}+\frac{1}{2x^2}+\frac{1}{3x^3}+\frac{1}{4x^4}+$&c.; consequently hyp. log. $\frac{x}{x-1}-\frac{1}{2x^2}-\frac{1}{3x^3}-\frac{1}{4x^4}-$&c. $=\frac{1}{x}$; hence, if we write 2, 3, 4, &c. for x, we have hyp. log. $\frac{2}{1}+$hyp. log. $\frac{3}{2}+$ &c. .... hyp. log. $\frac{x}{x-1}-\left\{\begin{array}{l}\frac{1}{2}\times\frac{1}{2^2}+\frac{1}{3^2}+\text{&c.}\ldots\ldots\frac{1}{x^2}\\ -\frac{1}{3}\times\frac{1}{2^3}+\frac{1}{3^3}+\text{&c.}\ldots\ldots\frac{1}{x^3}\\ -\frac{1}{4}\times\frac{1}{2^4}+\frac{1}{3^4}+\text{&c.}\ldots\ldots\frac{1}{x^4}\\ -\quad\text{&c.}\quad\text{&c.}\qquad\text{&c.}\end{array}\right\}=$ $\frac{1}{2}+\frac{1}{3}+\frac{1}{4}+$&c. .... $\frac{1}{x}$; but hyp. log. $\frac{2}{1}+$hyp. log. $\frac{3}{2}+$ hyp. log. $\frac{4}{3}+$&c. .... hyp. log. $\frac{x}{x-1}=$hyp. log. $\frac{2}{1}\times\frac{3}{2}\times\frac{4}{3}\times$ &c. .... $\frac{x}{x-1}=$hyp. log. x; also $\frac{1}{2^2}+\frac{1}{3^2}+$&c. .... $\frac{1}{x^2}=$ the

ſum of the ſame ſeries *ad infinitum*, *minus* the ſum of all the terms from $\frac{1}{x^2} = (\text{if } x+1=n)\ A - \frac{1}{n} - \frac{1}{2n^2} - \frac{1}{6n^3} + \frac{1}{30n^5} - \frac{1}{42n^7} + \&c.$; in the ſame manner $\frac{1}{2^3} + \frac{1}{3^3} + \&c. \ldots \frac{1}{x^3} = B - \frac{1}{2n^2} - \frac{1}{2n^3} - \frac{1}{4n^4} + \frac{1}{12n^6} - \frac{1}{12n^8} + \&c.$; and ſo on for the other ſerieſes; hence, by ſubſtitution, and adding unity to each ſide, we have hyp. log. $x+1-\frac{1}{2}A-\frac{1}{2}B-\frac{1}{2}C-\&c. + \frac{1}{2n} + \frac{5}{12n^2} + \frac{1}{3n^3} + \frac{31}{120n^4} + \frac{1}{5n^5} + \frac{41}{252n^6} + \frac{1}{7n^7} + \frac{31}{240n^8} + \&c. = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \&c. \ldots \frac{1}{x}$; but $1-\frac{1}{2}A-\frac{1}{2}B-\frac{1}{2}C-\&c. =$,577215664901; hence $1 + \frac{1}{2} + \frac{1}{3} + \&c. \ldots \frac{1}{x} =$ hyp. log. x + ,577215664901 $+ \frac{1}{2n} + \frac{5}{12n^2} + \frac{1}{3n^3} + \frac{31}{120n^4} + \frac{1}{5n^5} + \frac{41}{252n^6} + \frac{1}{7n^7} + \frac{31}{240n^8} + \&c.$

Ex. 1. Let $x = 10000$; then

hyp. log. 10000	= 9,210340371976
conſt. quant.	= ,577215664901
$\frac{1}{2n}$	= ,000049995000
$\frac{5}{12n^2}$	= ,000000004166
therefore the ſum required	= 9,787606036043

Ex. 2. Let $x = 10000000$; then

hyp. log. 10000000	= 16,118095650958
conſt. quant.	= ,577215664901
$\frac{1}{2n}$	= ,000000049999
therefore the ſum required	= 16,695311365858

PROP. XVI.

To find the value of $\alpha \times \beta \times \gamma \times \delta \times$ &c. *ad infinitum, supposing the general term to be a rational function of* x.

Let π be the general term, then resolve $\frac{\dot{\pi}}{\pi}$ into an infinite series, and take the fluent on both sides; then write 2, 3, 4, &c. for x, and one side will become the hyp. log. of the given series, and the value of the other side may be found from the tables.

Ex. 1. To find the value of $\frac{4}{3} \times \frac{9}{8} \times \frac{16}{15} \times$ &c. *ad infinitum.*

Here the general term is $\frac{x^2}{x^2-1}$; hence $\frac{\dot{\pi}}{\pi} = -\frac{2\dot{x}}{x^3-x} = -\frac{2\dot{x}}{x^3} - \frac{2\dot{x}}{x^5} - \frac{2\dot{x}}{x^7} -$ &c.; hence the hyp. log. $\frac{x^2}{x^2-1} = \frac{1}{x^2} + \frac{1}{2x^4} + \frac{1}{3x^6} +$ &c. Write 2, 3, 4, &c. for x, and we have the hyp. log. $\frac{4}{3}$ + hyp. log. $\frac{9}{8}$ + hyp. log. $\frac{16}{15}$ + &c. = A + $\frac{1}{4}$C + $\frac{1}{3}$E + &c. = ,693147180574, which is the hyp. log. 2; but hyp. log. $\frac{4}{3}$ + hyp. log. $\frac{9}{8}$ + hyp. log. $\frac{16}{15}$ + &c. = hyp. log. $\frac{4}{3} \times \frac{9}{8} \times \frac{16}{15} \times$ &c. consequently $\frac{4}{3} \times \frac{9}{8} \times \frac{16}{15} \times$ &c. = 2.

Ex. 2. To find the value of $\frac{8}{7} \times \frac{27}{26} \times \frac{64}{63} \times$ &c. *ad infinitum.*

Here the general term is $\frac{x^3}{x^3-1}$; hence $\frac{\dot{\pi}}{\pi} = -\frac{3\dot{x}}{x^4-x} = -\frac{3\dot{x}}{x^4} - \frac{3\dot{x}}{x^7} - \frac{3\dot{x}}{x^{10}} -$ &c.; hence the hyp. log. $\frac{x^3}{x^3-1} = \frac{1}{x^3} + \frac{1}{2x^6} + \frac{1}{3x^9} +$ &c. Write 2, 3, 4, &c. for x, and we have hyp. log. $\frac{8}{7}$ + hyp. log. $\frac{27}{26}$

$\frac{27}{26}$+hyp. log. $\frac{64}{63}$+&c.=B+$\frac{1}{2}$E+$\frac{1}{3}$H+&c.=,211466250444; or hyp. log. $\frac{8}{7}\times\frac{27}{26}\times\frac{64}{63}\times$ &c. =,211466250444; hence $\frac{8}{7}\times\frac{27}{26}\times\frac{64}{63}\times$ &c.=1,627295, &c.

Hence we may find the value of such a quantity, supposing the number of factors to be finite.

Ex. To find the value of $\frac{2}{1}\times\frac{4}{3}\times\frac{6}{5}\times$ &c. . . . . to $\frac{2x}{2x-1}$.

Here the general term being $\frac{2x}{2x-1}$, we have $\frac{\dot{w}}{w}=-\frac{\dot{x}}{2x^2\cdot x}=-\frac{\dot{x}}{2x^2}-\frac{\dot{x}}{4x^3}-\frac{\dot{x}}{8x^4}-\frac{\dot{x}}{16x^5}-$ &c.; hence hyp. log. $\frac{2x}{2x-1}=\frac{1}{1\cdot 2x}+\frac{1}{2\cdot 4x^2}+\frac{1}{3\cdot 8x^3}+\frac{1}{4\cdot 16x^4}+$ &c.. Now write 2, 3, 4, &c. for x, and we have the hyp. log. $\frac{4}{3}\times\frac{6}{5}\times\frac{8}{7}\times$ &c. . . . .

$$\left.\begin{aligned}\frac{2x}{2x-1}=&\frac{1}{2}\times\overline{\frac{1}{2}+\frac{1}{3}+\frac{1}{4}+\ldots\ldots\frac{1}{x}}\\&+\frac{1}{2\cdot 4}\times\overline{\frac{1}{2^2}+\frac{1}{3^2}+\frac{1}{4^2}+\ldots\ldots\frac{1}{x^2}}\\&+\frac{1}{3\cdot 8}\times\overline{\frac{1}{2^3}+\frac{1}{3^3}+\frac{1}{4^3}+\ldots\ldots\frac{1}{x^3}}\\&+\text{ &c.}\qquad\text{ &c.}\qquad\overline{\text{ &c.}}\end{aligned}\right\}$$

But, by Prop. 15. $\frac{1}{2}\times\overline{\frac{1}{2}+\frac{1}{3}+\frac{1}{4}+\ldots\ldots\frac{1}{x}}=\frac{1}{2}$hyp. log. $x-,211392167549+\frac{1}{4x}+\frac{5}{24x^2}+\frac{1}{6x^3}+\frac{31}{240x^4}+\frac{1}{10x^5}+\frac{41}{504x^6}+$ &c.; also $\frac{1}{2\cdot 4}\times\overline{\frac{1}{2^2}+\frac{1}{3^2}+\ldots\ldots\frac{1}{x^2}}=\frac{1}{2\cdot 4}\times\overline{A-\frac{1}{x}-\frac{1}{2x^2}-\frac{1}{6x^3}+\frac{1}{30x^5}-}$ &c. and $\frac{1}{3\cdot 8}\times\overline{\frac{1}{2^3}+\frac{1}{3^3}+\ldots\ldots\frac{1}{x^3}}=\frac{1}{3\cdot 8}\times\overline{B-\frac{1}{2x^2}-\frac{1}{2x^3}-\frac{1}{4x^4}+\frac{1}{12x^6}-}$ &c., and so on for the other series: hence, by substitu-

tion, hyp. log. $\frac{4}{3} \times \frac{6}{5} \times \frac{8}{7} \times$ &c. . . . . $\frac{2x}{2x-1} = \frac{1}{2}$ hyp. log. $x -$,1207822376 4 $+ \frac{1}{8x} + \frac{1}{8x^2} + \frac{23}{192x^3} + \frac{7}{64x^4} + \frac{61}{640x^5}$ + &c.; consequently $\frac{4}{3} \times \frac{6}{5} \times \frac{8}{7} \times$ &c. . . . . $\frac{2x}{2x-1}$ = the natural number corresponding to the right hand side of the equation; hence $\frac{2}{1} \times \frac{4}{3} \times \frac{6}{5} \times$ . . . . $\frac{2x}{2x-1}$ = twice that natural number.

Ex. Let $x = 10000$; then

$\frac{1}{2}$ hyp. log. $x = 4,6051701859 88$

const. quant. = ,1207822376 40

4,4843879483 48

$\frac{1}{8x} =$,0000124987 50

$\frac{1}{8x^2} =$,0000000012 49

4,4844004483 47 the natural number

corresponding to which hyp. log. is 88,6238, &c., consequently $\frac{2}{1} \times \frac{4}{3} \times \frac{6}{5} \times$ . . . . . $\frac{20000}{19999} = 177,2476$.

If x be a very large number, it may be sufficiently exact in most cases to take twice the natural number corresponding to the hyp. log. of $\frac{1}{2}$ hyp. log. $x -$,1207822376 40.

XXI. *Experiments and Obſervations to inveſtigate the Compoſition of* James's *Powder*. *By* George Pearſon, *M. D. F. R. S.*; *communicated by Sir* Joſeph Banks, *Bart. P. R. S.*

Read June 23, 1791.

THE medicine upon which many phyſicians principally depend in the cure of continued fevers is JAMES's *Powder*; but, although it has been very extenſively uſed above thirty years, the public have not, I believe, been informed of the particular nature of this ſubſtance. This *powder* was originally a patent medicine; but it is well known that it cannot be prepared by following the directions of the ſpecification in the Court of Chancery. Preſuming that I have made ſome experiments and obſervations which may explain the nature and manner of preparing this medicine, and, perhaps, may extend the hiſtory of antimony; I beg leave to have the honour of preſenting an account of them to the Royal Society.

Senſible properties of JAMES's *Powder*.

Some parcels of this preparation are white, but in general it has a yellowiſh caſt; and this ſhade is more evident in ſome ſpecimens than in others. It is ſaid, that this powder cannot, in general, be made at different times of preciſely the ſame ſhade of yellow or degree of whiteneſs. Sometimes with the aid of a lens a few very ſmall ſhining *ſpicula* are ſeen mixed with powder. When preſſed between the fingers it feels ſmooth,

ſmooth, with ſome rather rough particles, and it is gritty in the mouth. Moſt parcels at firſt are taſteleſs; but in about a minute there is a ſlight braſſy taſte. It is perfectly inodorous.

Specific gravity.

This powder feels much heavier than any of the common earths and ſtones in a pulverized ſtate. One of the phials, nearly full, in which it is ſold, reckoned to hold a quantity equal to twelve packets, or 480 grains, contained 470 grains troy weight of JAMES's powder. This phial, filled with diſtilled water to the ſame height that it had been by the powder, was found to contain nearly four drachm-meaſures, or about 240 grains, of this liquid *.

Effects of fire.

(*a*) The exterior part of the flame of a candle applied, by means of the blow-pipe, to about one, two, and three grains of JAMES's Powder on charcoal, and alſo in the ſpoon, only made it yellowiſh while hot, but, on cooling, this colour diſappeared. The interior and hotteſt part of the blue flame turned this powder yellow, and when continued ſo as to ignite it, a white inodorous fume or vapour aroſe, which ſoon ceaſed; and though the heat was continued, the powder neither appeared to diminiſh nor melt; but, on cooling, a ſlightly co-

* After this Paper had been read, an experiment, in a different manner, was made to aſcertain the ſpecific gravity of this powder. The quantity which nearly filled a phial weighed 437 grains; and filling the ſame phial, to the ſame height, with diſtilled water, the temperature of which was 65°, the water weighed 250,2 grains. The reaſon of the variation in theſe reſults, in making uſe of different parcels of this medicine, will be obvious, from the following account of its preparation, and the great difficulty of determining, with accuracy, the ſpecific gravity of powders.

hering

hering white ſolid, about three-fourths of the original weight of the powder, was left. If the flame was ſuddenly withdrawn, as ſoon as the white fumes appeared, they aſcended with a kind of revolving motion.

(*b*) Two grains of this powder, mixed with about three of pulverized tartar, being expoſed on charcoal to flame applied by the blow-pipe, the mixture turned black, boiled, and ſwelled; and by continuing to apply the flame, the coaly matter of the tartar diſappeared, a part of the mixture fuſed, and in that ſtate ſeveral ſmall, ſilvery, apparently metallic grains were perceived. On cooling, they were ſeen with the naked eye, or with a lens, adhering to an irregularly figured, partially melted, whitiſh maſs. On a ſecond application of the flame, theſe metallic globules diſappeared.

(*c*) JAMES's Powder, with glaſs of phoſphoric acid, melted into an opaque yellowiſh globule while hot, which on cooling grew whitiſh.

(*d*) This Powder, with ſeveral times its weight of melted borax, afforded a colourleſs tranſparent glaſs while fluid; but, on adding a larger proportion of powder, the globule turned opaque, and when cold became of a milky whiteneſs. As the JAMES's Powder mixed, or melted, with the fuſed ſalt, ſlight exploſions were ſeveral times heard.

(*e*) With foſſil alkali, in the ſpoon, this powder apparently fuſed, and afforded a colourleſs tranſparent fluid in a ſtate of rotatory motion; but on cooling it grew opaque, and had a horny appearance.

(*f*) 100 grains of this powder in a two-inch Engliſh crucible, the cover of which was luted on it as cloſely as poſſible, were expoſed above two hours to a fierce fire in a melting furnace. On breaking this crucible, when cold, the powder was

found changed into an entire very hard white ſolid, receiving its figure from the veſſel, and weighed 95 grains. On breaking this hard ſolid, the lower part of it ſeemed to be vitrified, or in a ſtate of enamel; and being powdered, it afforded a much whiter powder, and of greater ſpecific gravity, than before. The degree of fire denoted by WEDGWOOD's pyrometer was 166°.

Theſe experiments indicated the preſence of a metallic calx, a part at leaſt of which was that of antimony, mixed with earthy matter.

Experiments with different menſtrua applied to JAMES's *Powder.*

I. *With water.*

300 grains of this powder were digeſted for ſeveral hours in twenty-ſeven ounces of diſtilled water, and then boiled for one hour. While boiling the water appeared milky; but in half a minute's time, after withdrawing the lamp over which it boiled, the liquid became nearly clear, and the ſediment depoſited was apparently the powder undiminiſhed in quantity, and in other reſpects unaltered. While hot the liquid was decanted upon a filter of ſeveral folds of paper previouſly weighed, through which twenty ounces of quite clear liquid, like water, readily paſſed. Very little powder could be perceived on the filter; but when it was well dried, it weighed fourteen grains more than before the experiment. The filtered liquor was taſteleſs. In about three quarters of an hour it grew ſlightly *turbid*, and in ten minutes after became *milky*. On ſtanding eight days longer, the milkineſs diminiſhed, and a ſmall quantity, perhaps four grains, of cloſe white ſediment, firmly adhering to the ſides of the veſſel, were depoſited. A little of this ſlightly milky fluid being made hot, it grew quite clear, and on cooling

turned milky as before, or depoſited a ſediment; therefore, this milkineſs depended on cold water diſſolving a ſmaller proportion of JAMES's Powder than hot water. The whole of the filtered milky liquor was poured upon a filter as before, through which it paſſed without any diminution of its turbid appearance; and, on drying, the filter was found to have increaſed only a quarter of a grain in weight. Some of the vapour that aroſe during the ebullition being condenſed was found to be pure water.

On a repetition of this experiment the phænomena above related were always obſerved; but the ſemi-vitrified JAMES's Powder above-mentioned (*f*) afforded a much leſs milky fluid and ſediment than the powder uſed in the preceding experiment.

In order to determine the kind of ſubſtances in water after boiling in it JAMES's Powder, the following re-agents were added to the above filtered liquor.

1. Acid of ſugar ſometimes occaſioned immediately more turbidneſs, and at other times tranſparency was inſtantly produced; but in all caſes, on ſtanding, more ſediment fell than from the filtered liquor alone.

2. Muriated barytes in about an hour rendered this liquid evidently more turbid, and on ſtanding more ſediment was depoſited than from the filtered liquid to which nothing had been added.

3. Lime-water occaſioned immediately a curdy appearance.

4. Infuſion of turnſole was ſometimes turned to a ſlight red; but in general it was not altered in colour.

5. Nitre of ſilver produced in a few hours a ſlight ſediment.

6. Pruſſianated alkali of tartar occaſioned no alteration immediately, nor for four days after adding acetous acid to this mixture.

7. Mild alkali of tartar, and likewiſe mild foſſil alkali, though boiled in this liquor, did not occaſion any additional precipitation.

The ſediment that fell, on merely ſtanding, from the above filtered liquid, was next examined.

1. It did not diſſolve in 100 times its weight of boiling hot concentrated acetous acid; nor,

2. Was it apparently acted upon by boiling with mild alkali of tartar; for, after decanting the clear liquid of this mixture, the ſediment from it was not diſſolved by a large quantity of acetous acid, but readily by marine acid.

3. This ſediment immediately diſappeared on pouring upon it a much ſmaller quantity of marine acid, and alſo of acid of nitre, than had been applied of acetous acid, without any ſolution enſuing.

4. To one portion of this ſolution in marine acid juſt mentioned, was added gradually lixivium of alkali of tartar; after the effervescence had ceaſed to be produced, the firſt drop occaſioned a turbid appearance, and the liquid did not grow clear again on adding a large proportion of vinegar to make it ſour.

5. Another portion of this ſolution in marine acid, being boiled to carry off the ſuperabundant acid, was poured into a large proportion of lime-water, by which it was rendered cloudy, and did not become clear again on adding concentrated acetous acid to make it ſour.

6. To a third portion of this ſolution in marine acid, from which the redundant acid had been carried off, twenty drops of Pruſſian alkaline lixivium were added, which immediately turned it of a bluiſh caſt without diſturbing its tranſparency; and, after ſtanding four days, a ſmaller quantity of pale blue ſediment

ſediment was depoſited than had fallen from a quantity of diſtilled water equal to this muriatic ſolution, to which had been added twenty drops of this Pruſſian alkaline lixivium, and one-fourth part of a drop of muriated antimony. The ſame Pruſſian alkali mixed with vinegar, on ſtanding, turned bluiſh, but depoſited nothing.

7. Though the ſediment that fell in the filtered fluid, p. 320. on merely ſtanding, diſſolved in the nitrous acid as above mentioned; yet, when this acid was added in a ſmaller proportion, but to render the mixture ſour, a partial ſolution only took place. On adding, however, a very ſmall quantity of marine acid, the ſolution was total, and with leſs ſuperabundance of this menſtruum than of the nitrous acid in which a part remained undiſſolved. This ſolution

(*a*) With water depoſited Algaroth powder:

(*b*) With Pruſſian alkali turned bluiſh, and, on ſtanding, a ſmall quantity of ſediment took place.

(*c*) A bright plate of copper, immerſed in this ſolution, did not appear to be at all whitened, or rendered paler.

(*d*) Muriated barytes rendered this ſolution very turbid inſtantly.

(*e*) Acid of ſugar produced no change, except in two or three days a very minute portion of ſediment; but the ſaccharated ſoda immediately occaſioned a ſlight precipitation.

(*f*) Alkali of tartar, fully aërated, rendered this ſolution ſlightly turbid; but cauſtic mineral alkali induced no change.

(*g*) Lime-water, in a ſmall proportion, did not affect the tranſparency; but in a larger produced copious clouds.

The JAMES's Powder which afforded the ſolution in water, on which the experiments above related were made, was boiled a ſecond time in eighteen ounces of diſtilled water for

two hours. The decanted and filtered liquid on cooling grew leſs turbid than before, p. 320. and depoſited leſs ſediment. The filter, on drying, was found to have gained ten grains, though very little powder could be ſeen upon it. The experiments above related, were made on this filtered liquor, and in a ſlighter degree the ſame appearances were obſerved.

The JAMES's Powder remaining after theſe experiments, being well dried, weighed 260 grains, and therefore was found to have loſt 40 grains, partly by ſolution in water, but ſtill more by its adhering to the filters.

I wiſhed to know the proportion in which JAMES's Powder diſſolved in water; and therefore evaporated, in a very thin light glaſs pan, previouſly weighed, twenty-four ounces of the filtered liquids, p. 320. and p. 324. Very little precipitated matter appeared till nearly the whole of the liquor was evaporated; and, when the whole was carried off, a taſteleſs whitiſh, leafy, or mica-like ſediment, but in ſome parts black, was left, that weighed ſix grains. This ſediment required above 100 times its weight of hot water to diſſolve it. It was boiled in 500 times its weight of diſtilled water, and it paſſed through ſeveral folds of filtering paper rather turbid even while hot; nor could it be rendered clear by repeated filtration through paper. This filtered liquid,

(*a*) With infuſion of turnſole and turmeric, betrayed no alkaline ſubſtance, nor deciſively any acidity.

(*b*) Lime-water rendered it curdy; and on adding vinegar, it grew milky.

(*c*) With acid of ſugar it grew clear; but, on ſtanding, was more turbid than before.

(*d*) Salited

(*d*) Salited barytes made it inſtantly turbid.

(*e*) Alkalies, mild and cauſtic, induced no change.

(*f*) Pruſſian alkali produced only a clear greeniſh colour, after the addition of vinegar, and long ſtanding.

(*g*) A very ſmall quantity of marine acid rendered it quite clear; but it required much more nitrous acid to produce this effect; and this mixture did not whiten copper.

(*b*) With nitre of ſilver the filtered liquor turned of a ſomewhat bluiſh hue, and afterwards curdy.

The undiſſolved matter that remained on the filter, p. 324. l. 22. above mentioned, appeared, on examination, to be the ſame kind of ſubſtance, with a larger proportion of iron, as that which was diſſolved by water, the experiments on which have been juſt mentioned. In particular, it afforded Algaroth powder, but did not whiten, in the ſmalleſt degree, a copper plate.

The following concluſions may, perhaps, be juſtifiably drawn from theſe experiments on JAMES's Powder with water.

1. That the whole, or a part, is ſoluble, or at leaſt may be ſuſpended, in about 2000 times its weight of pure water cold; and in about half this quantity of boiling water.

2. That this ſolution contains calcareous earth united to an acid, or ſome other ſubſtance, from which it cannot be diſunited by cauſtic or mild fixed alkalies; therefore, the precipitation by muriated barytes cannot be referred to vitriolated lime.

3. That this ſolution contains a metallic calx, a part of which at leaſt is that of antimony uncombined, or at leaſt not united to any acid with which it forms a compound ſoluble in water.

4. That

4. That the ſubſtance in the nitrous ſolution of the part of JAMES's Powder that had been diſſolved in water, which precipitates lime from lime-water, and which precipitate is not ſoluble in a large quantity of vinegar, is, probably, phoſphoric acid from phoſphorated lime decompoſed by nitrous acid.

The precipitation by muriated barytes and nitrated ſilver could not be from vitriolic and marine acids conſiſtently with the preceding experiments; and I could not have conjectured what was the ingredient in JAMES's Powder which occaſions it, if I had not found, that muriated barytes is not only a teſt of vitriolic but of phoſphoric acid united to lime and alkalies; and the acid of phoſphorus will alſo produce a turbid appearance with nitrated ſilver. The calx of iron, in the above experiments, is in perhaps too ſmall a quantity to be conſidered in any other light than as an accidental ſubſtance.

II. *With acetous acid.*

The 260 grains of JAMES's Powder, remaining after boiling 300 grains of it in diſtilled water, and after the unavoidable waſte of it in the above experiments with water, were put into a tubulated retort that would contain four ounce-meaſures, on which were poured three ounces of concentrated acetous acid, the ſpecific gravity of which was as 106 to 100, the neck of the retort being immerſed in quickſilver, and the tubulated part being immediately cloſely ſtopped. No effervescence was perceived; nor did any elaſtic fluid riſe during twenty-four hours into an inverted veſſel of quickſilver; and when heat was applied to boil the acid, nothing but the common air of the retort and a little water and acetous acid came over.

This mixture of acetous acid and powder being poured, while hot, on a filter of two folds of paper, a clear and colourleſs liquid paſſed through, that remained ſo, when cold, without any ſediment. The filter, with the powder upon it, being well dried, weighed ten grains more than the ſum of their weight ſeparately before the experiment; but the powder being carefully ſcraped off weighed only a little more than 240 grains, and appeared to have ſuffered no change in its properties.

This filtered liquor was ſubjected to diſtillation; it remained clear till about half of it had come over: but then it became rather turbid, and grew more ſo to the end of the diſtillation. There remained in the retort apparently four or five grains of brown ſediment, that adhered very cloſely to the bottom and ſides of it, nearly half as high as the liquid reached.

1ſt, This acetous acid, in which JAMES's Powder had been boiled, and afterwards diſtilled, was found to contain no earth, ſalt, or metallic matter: nor did the acid itſelf appear to be altered in its chemical qualities.

2dly, The reſiduum in the retort had no taſte. It did not appear to diſſolve by trituration in one ounce of diſtilled water, nor in lixivium of alkali of tartar, mild or cauſtic. After boiling this reſiduum in one and a half ounce of water, part of it ſeemed to be diſſolved; and this ſolution being filtered was examined with the following re-agents.

(*a*) Acid of ſugar rendered it turbid, and tranſparency did not enſue on adding a further quantity of this acid.

(*b*) Muriated barytes produced a conſiderable turbidneſs, which was not removed by adding concentrated acetous acid.

(*c*) Nitrated ſilver induced a ſlight turbid appearance.

(*d*) Mild alkalies induced no viſible change.

(*e*) Pruſſianated

(*e*) Pruſſianated foſſil alkali occaſioned a very ſlight opal coloured appearance, and after ſtanding ſix days a very ſmall quantity of whitiſh ſediment was thrown down, which diſſolved on concentrated acetous acid being added, and the liquid turned greeniſh; but, after ſtanding, a little greeniſh ſediment was depoſited, not, however, apparently more than was produced by a mixture of this Pruſſian alkali, acetous acid, and water.

(*f*) Phoſphorated foſſil alkali produced no turbidneſs or precipitation on ſtanding a week; but on the addition of $\frac{1}{100}$ gr. of nitrated mercury (which had been made by fully ſaturating the nitrous acid with mercury) the mixture grew inſtantly thick, and depoſited a copious white ſediment.

3dly, The remainder of the reſiduum, in the retort above mentioned to have been left after diſtilling to dryneſs acetous acid in which JAMES's Powder had been boiled, did not totally diſſolve in a large proportion of nitrous acid diluted; but was wholly taken up on adding a little marine acid. A great part of the ſuperabundant acid of this ſolution being carried off by evaporation, it was examined with the following ſubſtances.

(*a*) Adding a little of it to a large proportion of water, milkineſs enſued.

(*b*) The ſame appearance took place with a large proportion of lime-water.

(*c*) A turbid bluiſh colour was produced on adding Pruſſian mineral alkali, and on ſtanding a bluiſh ſediment took place.

(*d*) A poliſhed copper plate was not at all whitened by immerſion in this ſolution; but, on adding to it $\frac{1}{400}$ of a grain of nitre of mercury, the copper was rendered paler coloured.

The deficiency of twenty grains of powder in theſe experiments with acetous acid muſt be aſcribed partly to the

ſolution in this menſtruum, and ſtill more to the waſte in the paper filter, and to its adhering to the retort when firſt boiled.

Theſe experiments indicated the ſame kind of ſubſtances as the experiments with water, namely, calcareous earth in a combined ſtate; phoſphoric acid; calx of antimony and of iron. It appears alſo, that JAMES's Powder is either wholly or partially ſoluble in about 300 times its weight of concentrated acetous acid.

III. *With nitrous acid.*

The 240 grains of powder remaining after the experiments with acetous acid, p. 327. were digeſted in the cold twelve hours, in three ounce meaſures and a half of purified and concentrated nitrous acid, diluted with four ounces of pure water, and then diſtilled with a gentle heat till there remained about two ounce meaſures. After ſtanding twenty-four hours, about one ounce and a half of clear liquid, which was very corroſive and acid, was drawn off by means of a ſyphon. The turbid liquid and cloſe white ſediment remaining, being mixed with one ounce of diſtilled water, were poured upon a filter of paper, and hot diſtilled water was repeatedly poured upon this filter till it paſſed through almoſt taſteleſs. The firſt portions of the filtered liquid, in quantity ten ounces, being mixed together, were ſet to evaporate in a glaſs pan. As ſoon as the liquor grew hot, the turbidneſs diſappeared; and as the evaporation went on, firſt clouds, and afterwards portions of ſediment, appeared in a clear liquid. The evaporation being continued nearly to dryneſs, a white, porous, or cellular cake was left, that weighed 129 grains. The liquid obtained

by diſtillation from the ſolution which left this maſs was found to be merely diluted nitrous acid.

The reſiduum left on the filter which had reſiſted ſolution in nitrous acid, being well dried, weighed a little more than 142 grains. This reſiduum was digeſted, and boiled as before in nitrous acid; and this menſtruum, diſtilled from the reſiduum, being evaporated to dryneſs, afforded 6,5 grains of a whitiſh maſs. The reſiduum left on the filter after this ſecond application of nitrous acid, being well dried, weighed 132 grains.

The maſs of 129 grains, left on evaporation of the above ſolution of JAMES's Powder in nitrous acid, in a few hours began to deliqueſce, eſpecially at the edges. Some of the deliqueſcing part of the maſs was diſſolved in one ounce and a half of water, forming an opal-coloured ſolution, with a white ſediment. This opal-coloured ſolution being filtered was examined.

(*a*) It rendered lime-water milky; and the milkineſs did not diſappear on adding concentrated acetous acid, but readily on pouring into the mixture a little acid of nitre.

(*b*) It turned thick and white, and ſoon depoſited a copious ſediment of white matter, with a few drops of nitrous ſolution of mercury; and became turbid alſo with nitrated ſilver.

(*c*) With muriated barytes it became very turbid, and remained ſo after adding acetous acid; but grew clear again on adding nitrous acid.

(*d*) Acid of ſugar produced a turbid appearance.

(*e*) Cauſtic volatile alkali produced very little precipitation; but a copious one took place with mild alkali of tartar; which precipitated matter, after decanting the clear ſolution, was nearly all diſſolved by acetous acid, and the remainder was readily taken up by the marine as well as by the nitrous acid.

(*f*) Pruſſian

(*f*) Pruſſian alkali occaſioned a light blue colour, but no turbid appearance till the mixture had ſtood ſeveral days, which was then ſlight.

(*g*) No change of colour was produced on a copper plate.

A little of the ſoft and deliqueſcing part, juſt mentioned to have taken place at the edges of the ſolid maſs, taſted bitter and ſour. It melted under the blow-pipe into a horny kind of globule; but the dry part of this maſs could not be fuſed by this means.

This ounce and a half of ſolution being conſumed in theſe trials, the ſame quantity of boiling diſtilled water was poured on the precipitate or part not diſſolved by this quantity of water on the firſt affuſion. After ſtanding and boiling, the precipitate appeared to be but little diminiſhed. The clear liquid did not as before render lime-water turbid; but a precipitate enſued with nitrated mercury, which on compariſon was found to be a more delicate teſt of phoſphoric acid than lime-water. This ſolution alſo, by this ſecond affuſion of water, did not as before grow thick with mild alkali of tartar, nor blue with Pruſſian alkali. The ſediment, undiſſolved by theſe two applications of water, did not totally diſſolve in a ſuperabundant quantity of nitrous acid; but completely and immediately in a ſmaller quantity of marine acid; and this ſolution in marine acid, with a large proportion of water, produced milkineſs; with Pruſſian alkali, it turned of a deep blue colour; it did not whiten copper; ſaccharine acid and ſalited barytes only ſlightly diſturbed its tranſparency.

By theſe experiments I found the ſolution of JAMES's Powder in nitrous acid contained, probably, a pretty conſiderable proportion of calcareous earth united to both nitrous

and phoſphoric acid; a little phoſphoric acid in a free ſtate; and a ſmall proportion of calx of antimony and of iron.

Ninety grains of the dry part of the above maſs of 129 grains, p. 329. were repeatedly triturated and digeſted in alcohol till almoſt nothing was taken up by it. This ſolution, being filtered, was evaporated to dryneſs, and afforded 20 grains and a half of a fine white ſalt, very bitter, which, on expoſure to the air, ſoon became liquid, but very turbid.

The powder that had thus ceaſed to yield any thing further to alcohol was repeatedly triturated and boiled in pure water, till the liquid paſſed taſteleſs through the filter; and the filtered liquors, being evaporated, left eight grains more of a leſs bitter and leſs deliqueſcent ſubſtance than that from alcohol. Part only of theſe eight grains was ſoluble readily in water; and they appeared to be a mixture of the ſaline matter diſſolved in alcohol, and of the inſoluble reſiduum in that menſtruum.

The powder remaining on the filter after theſe ſolutions in alcohol and water being dried, weighed nearly 59 grains. It was white and taſteleſs.

I next examined theſe products more particularly; and *firſt* the twenty grains and a half which had been diſſolved in alcohol.

(*a*) With a large proportion of water it produced a rather turbid appearance, and, after ſtanding, a ſediment of calcareous earth was depoſited.

(*b*) This laſt ſolution (*a*) being filtered, with mild alkalies grew very thick, and depoſited a ſediment that was readily taken up by acetous acid.

(*c*) With cauſtic volatile alkali its tranſparency was ſcarcely diſturbed.

(*d*) With acid of ſugar it became thick and white; and

(*e*) White

(*e*) White as cream of milk with phoſphorated mineral alkali; the ſediment from which mixture did not diſſolve in a quantity of boiling water that would have diſſolved vitriolated lime, nor in vinegar, but was readily taken up by nitrous acid.

(*f*) The tranſparency of nitrated and muriated barytes was ſcarcely diſturbed.

(*g*) It turned infuſion of turnſole red.

(*h*) A little of the deliqueſcent ſalt above mentioned, that had been diſſolved in alcohol, being made nearly dry, on adding to it a mixture of alcohol and acid of vitriol, vapours of nitrous ether were detached with ebullition

(*i*) With lime-water it produced a ſlight ſediment.

(*k*) With Pruſſian alkali at firſt a pale green colour, and afterwards a blue colour was produced; but without any precipitation on ſtanding.

(*l*) This ſubſtance, which had been diſſolved in alcohol, was infuſible under the blow-pipe; and after being heated red-hot on charcoal it was no longer ſoluble in water. Being further examined, it was found to be merely calcareous earth.

This ſoluble part then in alcohol appeared to be nothing but nitrated lime, with ſome traces of calcined iron.

Secondly, The 59 grains of powder, not ſolubie in alcohol, were examined.

(*a*) A mixture of vitriolic acid and alcohol detached from this powder no nitrous ether, nor any vapour that formed white clouds with volatile alkali.

(*b*) It did not effervеſce, and required above 200 times its weight of concentrated acetous acid to diſſolve it.

(*c*) Under the blow-pipe it emitted no ſmell or fume, and with great difficulty melted imperfectly, affording an irregular figured, horn-like, opaque maſs.

(*d*) It

(*d*) It was not diminiſhed or altered in its properties by boiling in lixivium of alkali of tartar.

(*e*) Nitrous acid formed with it, without effervescence, a very ſlightly turbid ſolution; which ſolution produced the ſame appearances as thoſe related, p. 330, 331.

(*f*) 50 grains of this powder were diſſolved in nitrous acid, and a great part of the redundant acid being carried off by evaporation, to one half of this ſolution was added lime-water till it ceaſed to produce any milkineſs, and the mixture taſted of lime-water. After ſtanding excluded from the air, the ſediment depoſited from a clear liquid was collected, and being dried it weighed 26,3 grains. This precipitate had the properties of phoſphorated lime, with that proportion of lime and phoſphoric acid which forms a compound ſcarcely fuſible. The liquid in which this precipitate fell ſeemed to contain a little phoſphorated lime, but principally calcareous earth.

(*g*) To the other half of this ſolution in nitrous acid (*f*) was added vitriolic acid, drop by drop, till it no longer diſturbed its tranſparency. After ſtanding, the clear liquid was decanted from the precipitated matter which had taken place, and the precipitate with a ſmall quantity of water was thrown upon a filter. The filtered and decanted liquids mixed together were boiled till the ſmell of nitrous acid ceaſed, and there remained about half an ounce of acid liquor, which being filtered to ſeparate the vitriolated lime precipitated during evaporation, foſſil alkali was added to perfectly ſaturate it. During this union there was an effervescence and a ſeparation of more vitriolated lime, which being removed, the ſaturated liquor, by cryſtallization, afforded nearly 26 grains of cryſtals of phoſphorated foſſil alkali, beſides

ſides a little cubic nitre, vitriolated foſſil alkali and iron, with ſome veſtiges of calx of antimony, and phoſphorated lime.

The precipitate thrown down on adding vitriolic acid, and left upon the filter, weighed when dried 26,1 grains, and was vitriolated lime, with a minute portion of calcined antimony and iron.

The 6,5 grains, p. 330. left on evaporation to dryneſs of the ſecond ſolution in nitrous acid, conſiſted of nearly three grains of calcined antimony, and the reſt phoſphorated lime, with a little iron.

1. It appears from the above experiments with nitrous acid, that this menſtruum, by two affuſions, in a large proportion, aided by trituration, digeſtion, and heat, diſſolved $\frac{108}{240}$ of JAMES's Powder that had been expoſed to the action of water and acetous acid; but from the ſmallneſs of the quantity contained in the nitrous acid the ſecond time it was applied, and from its being principally calcined antimony, not more than two of the ſix grains afforded by this ſolution, perhaps, ſhould be conſidered to be *diſſolved*, for the reſt may be ſuppoſed to be merely *ſuſpended*.

The firſt ſolution alſo in this menſtruum was not filtered, and the acid was conſiderably redundant, and there was found in it ſeveral grains of calcined antimony. The real quantity *diſſolved* might therefore probably be eight grains leſs than the above 108 ſtated. According to this mode of calculation, the proportion of the ſoluble part of JAMES's Powder in nitrous acid is $\frac{100}{240}$, or about $\frac{41}{100}$.

2. The whole of this ſoluble part, except a little calx of antimony, is, deciſively, phoſphoric acid and calcareous earth: which

which two ſubſtances may reaſonably be ſuppoſed, from theſe experiments, to have been united together, and to have been in the ſtate of phoſphorated lime in this powder. Conſequently, the proportion of this phoſphorated lime, conſidered as the ſoluble part of JAMES's powder in theſe experiments with nitrous acid, appears to be 40 *per cent.* making a deduction of 1 *per cent.* for the antimonial calx contained in the nitrous acid in the above experiments. It is however already obvious to ſuſpect, that the powder which reſiſted ſolution in this menſtruum may contain more phoſphorated lime; and this conſideration prevents me aſſigning at preſent the above 40 *per cent.* as the whole quantity of it in JAMES's Powder. It cannot however, I think, be a ſmaller proportion.

I do not reckon the calx of iron in theſe calculations, becauſe it is in too ſmall a quantity, and is apparently only to to be looked upon as an accidental extraneous ſubſtance. I ſuppoſe too, that the water and acetous acid applied to the JAMES's Powder uſed in theſe experiments, carried off a proportion of its ingredients equal to that in the remaining powder.

IV. *With marine acid.*

The 132 grains of heavy, white, taſteleſs powder, the reſiduum after boiling 240 grains of JAMES's Powder in nitrous acid, till it had diſſolved that part for which it has any conſiderable affinity, p. 330. were digeſted for twenty-four hours in eight ounce-meaſures of marine acid, the ſpecific gravity of which was 1,170, and diluted with half its bulk of diſtilled water. This mixture was diſtilled in a

gentle heat till there remained about two ounce-meaſures of a very turbid liquor. After ſtanding in a jar two days, it depoſited a cloſe white ſediment, obviouſly much leſs in bulk than the powder added to this menſtruum; and nearly one ounce meaſure and three quarters of clear yellow liquid were drawn off, by means of a ſiphon, which was marked N° 1.

The diſtilled liquid, which was merely diluted marine acid, was poured back on the ſediment and remaining liquid; and after digeſtion twenty-four hours, this mixture was diſtilled as before, till there remained about one and a half ounce-meaſure; but after ſtanding in a jar ſeveral days, the quantity of ſediment depoſited was apparently as great as before the ſecond application of this menſtruum. The clear liquor was drawn off as before, and marked N° 2. The diſtilled liquor being found to be merely diluted marine acid, was poured on the remaining liquid and ſediment a third time; and, after digeſtion, the diſtillation was repeated as before. The remaining liquid having ſtood upon the ſediment ſome time, one ounce-meaſure of clear liquor was drawn off, and marked N° 3. The ſediment did not appear diminiſhed by this third diſtillation; but, as the decanted liquid, N° 3. was found to contain a ſmall quantity of ſome ſubſtance diſſolved or ſuſpended in it, the marine acid diſtilled in this experiment was poured a fourth time on this reſiduum, and after digeſtion boiled. Having ſtood ſeveral days, the clear liquor was decanted, and marked N° 4. To the reſiduum, after theſe four affuſions of marine acid, one ounce of boiling diſtilled water was added, and this mixture was poured on a filter. The powder upon the filter being well dried, was found to weigh 60,1 grains.

In the ounce of boiling water that had been filtered from this reſiduum, I could find nothing but ſome minute particles of that ſubſtance and veſtiges of iron. Two or three drops of the liquor, N° 1. added to three ounces of diſtilled water, produced a pretty conſiderable milkineſs; and, on ſtanding, a cloſe white ſediment was depoſited. Two or three drops of the liquor, N° 2. produced leſs milkineſs and ſediment in the quantity of water juſt mentioned than N° 1. N° 3. ſcarcely diminiſhed the tranſparency of diſtilled water; and N° 4. did not affect it all. The liquids, N° 3. and 4. were diſtilled till there remained about two drachm-meaſures of clear brown liquid, with a cloudy ſediment. N° 3. being poured into two ounces of diſtilled water ſcarcely made it milky; and N° 4. did not diminiſh the tranſparency of water at all. On evaporating to dryneſs theſe mixtures of the liquors, N° 3. and 4. with diſtilled water; that with N° 3. left a yellowiſh dry ſediment which weighed 3,1 grains; and that of N° 4. left 2,2 grains of ſediment. Nothing but calx of antimony could be diſcovered in theſe ſediments. They were reducible readily with tartar; ſcarcely fuſible in the ſpoon; but with phoſphoric acid eaſily melted into an opaque yellowiſh globule.

The liquor, marked N° 1. being poured into twenty-four ounces of diſtilled water produced a very milky appearance; and the ſame appearance, but in a much leſs degree, enſued on pouring the liquor N° 2. into this quantity of pure water. After ſtanding ſeveral days, a white ſediment being depoſited from a clear watery fluid, the clear liquid was drawn off by means of a ſiphon, and the ſediments being dried, that of N° 1. weighed 51 grains; that of N° 2. weighed 6,15 grains; and both were found to be purely Algaroth powder. The water, *viz.* 48 ounces, in which theſe precipitates fell, being evaporated

evaporated to dryneſs, left a little more than four grains of an infuſible kind of ſediment, which was calx of antimony, like that of N° 3. and 4. with a minute portion of Algaroth powder and iron.

The 60,1 grains, p. 337. which reſiſted ſolution in marine acid, were a lighter powder than JAMES's Powder itſelf, rather gritty, quite taſteleſs. This powder was digeſted in one-ounce meaſure of concentrated nitrous acid mixed with two of acid of ſalt for a week; and then this mixture was diluted with four ounces of diſtilled water, and gently boiled till it was reduced to about two ounce-meaſures. On cooling and ſtanding ſeveral days, a ſediment of the ſame kind apparently, and in the ſame quantity as before evaporation, was depoſited, and, after ſtanding, a clear yellow liquid was drawn off. The ſediment, well dried, weighed 55 grains. The decanted liquor being reduced by evaporation from one and a half ounce to about two drachm-meaſures, a ſediment, while hot, appeared, which was calx of antimony, with a minute portion of earthy matter that had ſome properties of phoſphorated lime and calx of iron. A little of the clear liquor on this ſediment being dropped into water produced no milkineſs, nor ſediment, after ſtanding; but the whole of this clear liquor, with ſediment, being evaporated to dryneſs, afforded four grains of the ſame kind of infuſible calx of antimony, mentioned to have been obtained from the acid liquors, N° 3. and 4. above mentioned.

It appears then, that by repeatedly digeſting and boiling in marine acid, and in *aqua regia*, that part of JAMES's Powder which reſiſted ſolution in nitrous acid, which was $\frac{133}{144}$, p. 335. 77 grains were carried off by theſe menſtrua; but conſidering the ſmall proportion contained in theſe acids after the two firſt

affusions, which afforded 57,15 grains, p. 338., and supposing the calx to be neither increased nor diminished in weight by the acids, the real quantity of soluble and fusible calx of antimony may be stated to be that of Algaroth powder; for the other kind of antimonial calx obtained by subsequent affusions was probably only *suspended*. Consequently 240 grains of JAMES's Powder afforded, by the above experiments with marine acid, 57,15 grains of Algaroth powder, and 19,85 grains of a less soluble and more difficultly fusible calx of antimony, with a small proportion of phosphorated lime. The residuum, amounting to 55 grains, was of course next examined.

Experiments upon the part of JAMES's *Powder which was insoluble in the above menstrua.*

(*a*) A few grains of this insoluble substance could neither be melted nor carried off in vapour by means of the flame of a candle applied to it upon charcoal, and also in the spoon, with the blow-pipe; but,

(*b*) Mixed with an equal weight of tartar, it melted on charcoal; and, while in fusion, small, apparently, metallic grains were distinctly perceived; and on cooling they could be seen, even without a lens, adhering to an irregularly figured, opaque, whitish mass. Sometimes slight explosions were heard while the flame was applied. The metallic grains appeared more distinctly when this powder was mixed with one-third of its weight of powdered calcined bone, than in JAMES's Powder.

(*c*) Ten grains of this powder were melted as above mentioned, by repeatedly applying flame with the blow-pipe to two

two or three grains of it at a time mixed with tartar. The opaque whitiſh maſſes, with metallic grains in them, thus obtained, being pulverized, were digeſted and gently boiled in diluted nitrous acid. The filtered ſolution afforded nitre, and nitrous acid in a free ſtate, the greateſt part of which ſuperabundant acid being carried off, the lixivium did not whiten copper, or throw down any calx but iron with Pruſſian alkali. The reſiduum that reſiſted ſolution in nitrous acid was digeſted and gently boiled in *aqua regia*. On ſtanding it was decanted, and this decanted liquid being heated, to carry off ſuperabundant acid and water, it afforded on mixture with water 1,2 grain of Algaroth powder, and no metallic matter could be detected in the water excepting a little iron. A ſmall part only being diſſolved by the *aqua regia*, the reſiduum was expoſed to the flame of a candle with tartar as before; and, by the aid of a lens, I could juſt perceive two or three metallic grains in the fuſed maſs. To this maſs the *aqua regia* was again applied, and 0,15 grain of Algaroth powder was obtained, and no other metallic calx was found but iron. A third affuſion of *aqua regia* indicated an exceedingly minute portion of Algaroth powder; but I could afterwards perceive no more metallic grains in the reſiduum expoſed to flame with tartar, nor obtain more Algaroth powder from the ſolution of the fuſed maſs in *aqua regia*. The reſiduum that reſiſted ſolution melted readily with a little phoſphoric acid into an opaque, ſomewhat yellowiſh, white globule, not unlike calcined bone fuſed with phoſphoric acid, and a minute portion of flowers of antimony. The quantity, however, of this reſiduum was ſo ſmall, that I deſpaired of determining its nature further by more experiments.

Having found that this insoluble powder would not melt with sulphur when heated *red-hot*, I made the following experiment.

(*d*) Twenty grains of it being mixed with three times its weight of sulphur, were put into one of Mr. WEDGWOOD's crucibles that would contain one ounce-measure, which, with a cover well luted on, was put into a three-inch English crucible, and calcined bone in powder filled the space between the two crucibles. After exposing this charge to a red heat half an hour, and in a white heat ten minutes, the crucibles were cooled; and being opened, the pyrometer piece of WEDGWOOD in the bone ashes was found to indicate 65°, and the mixture in the inner crucible had apparently been melted into a resin-like mass that adhered firmly to the sides of the vessel. Twenty-eight grains were scraped off, which, after digestion and boiling in marine acid, afforded six grains of Algaroth powder. A great deal of hepatic air was discharged during this solution, and very little sulphur was left on the filter with the part not dissolved by the marine acid. This undissolved part, which weighed six grains, was blackish, tasteless, not heavy. It was infusible with the blow-pipe, both alone, and mixed with sulphur and tartar; but with phosphoric acid it melted into a blackish scoria-like mass. I could only conjecture, that this last part was antimonial calx, so far vitrified with phosphorated lime as to be neither soluble nor reducible or fusible, except with phosphoric acid.

(*e*) By a similar experiment, but with alkali of tartar twenty grains, sulphur thirty grains, and ten grains of this insoluble part of JAMES's Powder, a fused mass was obtained that partially dissolved in water, and afforded kermes mineral on pouring an acid into this solution; but a great part was insoluble

inſoluble in water and acids, and ſeemed to be of the ſame nature as the ſix grains of reſiduum juſt mentioned (*d*).

I could only conclude from theſe experiments on this inſoluble matter, that it contained calx of antimony; but as to the proportion of it, and the other ſubſtance with which it is joined, I conjecture that it may be about half the quantity of the inſoluble powder; and that the other half is antimonial calx and phoſphorated lime, ſo highly calcined and vitrified together as to reſiſt ſolution in acid menſtrua, decompoſition by charcoal, and fuſion with fixed alkalies, but not by phoſphoric acid.

I ſhould not have been ſatisfied with here terminating this analyſis without enquiring further into the nature of this inſoluble matter; but I diſcontinued this analytic inveſtigation in order to derive light from the ſynthetic experiments which will be related hereafter.

Theſe laſt experiments ſeem to ſhew, that the proportion of antimonial calx is not ſo great as might have been aſſigned from the experiments with nitrous acid, marine acid, and *aqua regia*.

The ſubſtances and proportions of them, obtained from 240 grains of JAMES's Powder, by the above experiments, are as follow:

	Grains.
Phoſphorated lime, with a little antimonial calx,	100,
Algaroth powder,	57,15
Inſoluble antimonial calx, with a little phoſphorated lime,	19,85
The ſame inſoluble calx, with, probably, a little phoſphorated lime,	55,
Waſte,	8,
	240,0

As it may be objected, that concluſions drawn concerning the nature of calces might be erroneous if nitrous acid had been applied previouſly to ſubſtances containing them, I made the following experiment.

Experiment with marine acid applied to JAMES's *Powder, which had not been expoſed to the action of nitrous acid, or any other menſtruum.*

50 grains of JAMES's powder were digeſted, and gently boiled in two ounce-meaſures and a half of concentrated marine acid diluted with one ounce of diſtilled water till there remained only about one ounce-meaſure. A great part of the powder appeared to be evidently diſſolved. On cooling, cryſtals of muriated antimony were formed upon a white ſediment. The clear liquid with the cryſtals being decanted, the ſediment was boiled twice, as before, with marine acid; but the ſecond affuſion of this menſtruum brought off but eight grains of this powder, and the third only four grains. The remaining ſediment, being well dried, weighed 14 grains. Now it has been ſhewn already, that the nitrous and marine acids, ſucceſſively applied, diſſolved $\frac{177}{240}$, or all but about 60 grains; and in the preſent experiment, the marine acid diſſolved $\frac{36}{50}$, which is in the proportion of $\frac{180}{250}$, or nearly $\frac{173}{240}$; ſo that, on account of the trifling difference in theſe proportions, it may, perhaps, be fairly concluded, that the properties of the calx in JAMES's Powder are not altered by nitrous acid to affect its ſolubility in marine acid. And further, this inſoluble powder in the preſent experiment was found to have the ſame properties as that in the former experiments.

To

To know whether JAMES's Powder contained any ſubſtance that could be decompoſed by mild fixed alkalies, the following experiments were made.

Theſe experiments with fixed alkali ſeemed to be eſpecially neceſſary, becauſe phoſphoric acid, lime, and antimonial calx, are ingredients in JAMES's Powder; and it was obvious to ſuſpect, that this acid might be united with calx of antimony as well as with lime; which phoſphorated antimony would be decompoſed by alkalies, and yield phoſphorated alkali.

Experiments with fixed alkalies.

100 grains of JAMES's Powder were boiled in ſix ounces of water, with 50 grains of mild alkali of tartar, for three hours, and then the remaining liquid was filtered, and evaporated to dryneſs; but the matter left after evaporation was nothing but the alkali uſed in the experiment, with a little of the powder itſelf.

The reſult was the ſame on making the experiment with cryſtallized mineral alkali inſtead of alkali of tartar.

Synthetic Experiments.

ALTHOUGH the inability to prepare JAMES's Powder would not prove the above concluſions, with reſpect to its compoſition, to be erroneous; the being able to compoſe a ſubſtance

poſſeſſing all the ſame properties as JAMES's Powder, by uniting or mixing together the ſubſtances ſhewn by the above analyſis to enter into its compoſition, would afford all the proof and demonſtration which can be had in the ſcience of chemiſtry.

The above analyſis ſhewed no eſſential ingredients of JAMES's Powder but antimonial calces, phoſphoric acid, and calcareous earth, which two laſt ſubſtances appeared to be united together; but it would have been vain and unneceſſary labour to have attempted to make this powder by mixtures of any of the commonly known calces of antimony and phoſphorated lime; becauſe none of them, from their well known qualities, could form a powder of the ſame colour and ſpecific gravity as JAMES's Powder, and like it partially ſoluble in acids. From the above experiments, however, the probability was evident, that this ſubſtance might be made by calcining together antimony and bone-aſhes; which operation produces a powder called LILE's and SCHAWANBERG's fever-powder; a preparation deſcribed by SCHRODER and other chemiſts 150 years ago. The receipts for this preparation differed in the proportion of the antimony to the bone aſhes, and in the ſtate of the bone; ſome directing bone ſhavings to be previouſly boiled in water; others ordered them to be burnt to aſhes before calcining them with antimony; and in other preſcriptions the bone ſhavings were directed to be burnt with the antimony. According to the receipt in the poſſeſſion of Mr. BROMFIELD, by which this powder was prepared forty-five years ago, and before any medicine was known by the name of JAMES's Powder, two pounds of hart's horn ſhavings muſt be boiled to diſſolve all the mucilage, and then, being dried, be calcined with one pound of crude antimony, till the ſmell of

ſulphur

ſulphur ceaſes, and a light grey powder is produced. The ſame preſcription was given to Mr. WILLIS, above forty years ago, by Dr. JOHN EATON, of the College of Phyſicians, with the material addition, however, of ordering the calcined mixture to be expoſed to a great heat in a cloſe veſſel to render it *white*. Mr. TURNER made this powder above thirty years ago by calcining together equal weights of burnt hart's horn and antimony in an open veſſel, till all the ſulphur was driven off, and the mixture was of a light grey colour. He likewiſe was acquainted with the fact, that by a ſufficient degree of fire in a cloſe veſſel this cineritious powder turned *white* *. Mr. TURNER alſo prepared this powder with a pound and a half of hart's horn ſhavings and a pound of antimony, as well as with ſmaller proportions of bone. SCHRODER preſcribes equal weights of antimony and calcined hart's horn; and POTERIUS and MICHAELIS, as quoted by FREDERIC HOFFMAN, merely order the calcination of theſe two ſubſtances together (aſſigning no proportion), in a reverberatory fire for ſeveral days. In the London *Pharmacopœia* of 1788, this powder is called *Pulvis antimonialis*; and it is directed to be prepared by calcining together equal weights of hart's horn ſhavings and antimony.

Powders made from various proportions of antimony and bone-aſhes, after ſolution in nitrous acid, left a reſiduum of antimonial calx much leſs or greater in quantity than JAMES's Powder did by the ſame menſtruum, except two of Mr. TUR-

* It is probable, that this powder was made for ſeveral years with merely the heat neceſſary to carry off the ſulphur and calcine the bone, in an open veſſel over a charcoal fire in a common grate, and conſequently it was of a light clay or aſh colour. In this manner, Mr. BROMFIELD told me, he prepared SCHAWANBERG's Powder 46 or 47 years ago. Its property of turning *white* in a greater degree of fire appears to have been a ſubſequent diſcovery.

NER's proportions, *viz.* two parts of antimony and one of calcined bone, and equal weights of bone shavings and antimony. The quantity of this calx was, however, greater in the powder from the former of these two last proportions than the latter of them; which latter corresponded sometimes exactly, and always nearly, with the weight of the calx from a given weight of JAMES's Powder. This calx afforded also the same proportion of Algaroth powder as the calx in JAMES's Powder; and the insoluble part of the calx afforded metallic grains like those from the insoluble part of the calx in that powder.

I found then an exact correspondence between what I consider to be the essential and peculiar properties of JAMES's Powder, and the properties of a powder made by uniting or mixing together the ingredients of JAMES's Powder found by analysis. But, in order to shew the identity or difference of the qualities of these two substances, I made comparative observations on them, and repeated the above analytic experiments on JAMES's Powder with the preparation made by calcining together equal weights of bone shavings and antimony, in an open vessel, to carry off the sulphur, and then in close vessels applying such a degree of fire as to render them *white*, that is, on the same preparation as the *Pulvis antimonialis* of the London *Pharmacopœia*.

First, I compared, more particularly, the sensible qualities of several different specimens of JAMES's Powder with various parcels of the *Pulvis antimonialis* made by different chemists. All of these would be called white powders, but not two of them were so in the same degree. Most of the papers of the *Pulvis antimonialis* were whiter than those of JAMES's Powder; but others were of a very light stone colour, and some had a shade of yellow, so as to resemble very exactly JAMES's Powder; but

but all the parcels of JAMES's Powder had either a ſhade of yellow or of ſtone colour, and none were perfectly white, or ſo white as ſome ſpecimens of the *Pulvis antimonialis*. Some of the parcels of JAMES's Powder and of the *Pulvis antimonialis* taſted braſſy; and other ſpecimens of both powders had no taſte. All of theſe powders were gritty. Moſt of the parcels of the *Pulvis antimonialis* were a little ſpecifically heavier than thoſe of JAMES's Powder. The ſpecific gravity of both powders was increaſed by expoſing them to ſuch a degree of fire as brought them into almoſt a ſemi-vitrified ſtate; and, on the contrary, the ſpecific gravity of the *Pulvis antimonialis* was leſs than it is in its uſual ſtate, when made in ſuch a degree of fire that the mixture preſerves the powdery form.

The experiments with water on the *Pulvis antimonialis* produced the ſame kind of appearances, but more ſlightly than thoſe with JAMES's Powder; for the hot ſolution of the former grew leſs milky on cooling than that of the latter, and on evaporation to dryneſs leſs ſediment was found of the ſolution of *Pulvis antimonialis* than after that of JAMES's Powder *.

The experiments with acetous acid on the *Pulvis antimonialis* ſhewed, that this menſtruum diſſolved ſometimes a greater, and ſometimes a ſmaller proportion of it than of JAMES's powder; and the diſſolved matter was found to be antimonial calx, phoſphorated lime, and calx of iron, and no other ſubſtance.

It has been already ſaid, that the proportion of ſoluble matter in nitrous acid was the ſame, or nearly ſo, of the *Pulvis antimonialis* as that of JAMES's Powder; and this diſſolved matter was phoſphoric acid, calcareous earth, with a little antimonial calx, and a minute portion of calx of iron, as exactly

* The reaſon for this difference is aſſigned in another place.

as

as could be expected from the nature of the ſubſtances and the experiments, in the ſame proportion as thoſe in JAMES's Powder.

The Algaroth powder, obtained by means of ſolution of the *Pulvis antimonialis* in marine acid, was in the ſame proportion as nearly as could reaſonably be expected from the nature of the experiments as that obtained from JAMES's Powder. And the part that reſiſted ſolution in this menſtruum was partially reducible to a metallic form, and had otherwiſe the ſame properties, as far as diſcovered, as the inſoluble part of JAMES's Powder.

Having now formed a powder poſſeſſed of properties ſimilar in *kind* to every one of thoſe aſcertained in JAMES's Powder, with ſcarcely any difference in the *degree* of them, if it be thought that among theſe properties are thoſe which are eſſential and peculiar ones of JAMES's Powder, the concluſion that theſe two are the ſame kind of things muſt be admitted to be juſt. The nature of one of the ingredients of JAMES's Powder, *viz.* the irreducible part of the inſoluble matter, p. 342. is not fully elucidated by the ſynthetic experiments; but in ſo far as they ſhew, that this part equally exiſts in the powder formed by calcining together antimony and bone, which is concluded to be JAMES's Powder, the objection againſt the concluſion with reſpect to the identity of the two ſubſtances, on the ground of this inconſiderable part of JAMES's Powder not being well underſtood, muſt be of little weight.

Several reaſons, more intereſting to myſelf than to the Society, induced me to authenticate by additional teſtimonies thoſe analytic experiments, which may be conſidered to be more deciſive than the reſt for eſtabliſhing the identity of JAMES's Powder, and a powder formed by calcining together antimony and bone-aſhes. I therefore requeſted Mr. CAVALLO and Mr. TURNER to be

preſent when I made thoſe experiments on the *Pulvis antimonialis*, prepared by Mr. GRIFFIN, of Apothecaries' Hall, and JAMES's Powder. Having, in the preſence of theſe two Gentlemen*, broken the ſeal of a phial of JAMES's Powder, bought of F. NEWBERY, and taken out of it the quantity required for the experiments, the bottle was again ſealed by Mr. CAVALLO with his ſeal, as well as the phial from which was taken the *Pulvis antimonialis*. Should any experiments be publiſhed, which eſtabliſh different concluſions from thoſe contained in this Paper, with reſpect to the identity of theſe two powders, I ſhall be happy to endeavour to aſcertain the truth by experiments, on the remaining parcels of the two powders, in the preſence of competent judges.

I ſhall next relate the experiments made with the view of confirming or invalidating the concluſions drawn from the above analyſis, with reſpect to the ingredients and proportions of them in JAMES's Powder; and by which I eſpecially endeavoured to make ſuch antimonial calces as this ſubſtance contains, by proceſſes different from thoſe above related.

EXP. 1. (*a*) Hart's horn ſhavings, of ſix different parcels, well dried, ſeparately calcined in the ſame manner, and apparently to the ſame degree as when calcined with antimony to make LILE's Powder, afforded a light brown coarſe powder, with a few thin light black pieces, and loſt from 43 to 48 *per cent.* of their weight. The mean loſs of weight, of courſe, was 45¼ *per cent.*

(*b*) This calcined bone (*a*), being pulverized, was expoſed to a greater degree of fire, in cloſe veſſels, than that neceſſary to render the calcined mixture of antimony and bone-aſhes *white*. The loſs of weight by this ſecond calcination or expoſure to fire was from two to three *per cent.*; and the aſhes

* Dr. CLARKE alſo was preſent at the beginning of theſe experiments.

were

were as white as ſnow. The total mean loſs of weight, by theſe two calcinations, was then $\frac{4\,8}{1\,0\,0}$.

EXP. II. 2000 grains of coarſely powdered antimony were calcined in an earthen diſh, as in making LILE's Powder, by conſtantly raking them about for above three hours. During a great part of this time the veſſel was red hot at the bottom; and for the laſt hour the ſulphureous fumes had entirely ceaſed. The calx thus produced was of a pale bluiſh colour; it melted, in a low degree of heat, into an opaque, ſcoria-like brittle maſs; it yielded no hepatic air with marine acid; it weighed 1409 grains, or the antimony loſt nearly 29½ *per cent.* The pyrometer in the veſſel with the antimony during its calcination, was contracted to the 6th degree of WEDGWOOD's ſcale.

The ſum therefore of the loſs of antimony and bone by calcination in this manner, ſeparately, was 37½ *per cent.* Theſe two ſubſtances were in the next place calcined together in the ſame manner in an open veſſel, as above mentioned.

EXP. III. 2000 grains of antimony from the ſame parcel as that in the laſt experiment, and an equal weight of hart's horn ſhavings taken from the ſame parcel as thoſe were in Exp. 1. were calcined together in the ſame manner that theſe ſubſtances had been ſeparately. During the firſt quarter of an hour, the mixture ſmoked, was black, ſmelled ſtrongly of ſulphur, and felt ſoft. For half an hour more, the ſmell of ſulphur continued, the mixture turned brown, and the bone was reduced to aſhes. At the end of this time, not only the bottom of the veſſel might be kept red hot without any ſigns of fuſion; but the ſmell of ſulphur, though weakly, continued for half an hour more in a heat to keep a great part of the mixture red hot. At this time the ſulphureous ſmell rather ſuddenly diſappeared, and could not be perceived, though a little

little of the mixture was made quite red hot for a quarter of an hour further; during which no fume was ſeen, or ſmell perceived. After cooling, a light grey or cineritious heavy powder was left; on examining which, argentine *ſpicula* were ſeen in the larger grains of this calcined ſubſtance. It weighed 2200 grains, therefore the loſs of weight was 45 *per cent.* The WEDGWOOD pyrometer pieces indicated 8°. In other ſimilar experiments the loſs, by calcination, was from 37 to 41 *per cent.*; therefore the mean proportion loſt in theſe experiments muſt be ſtated at 41 *per cent.*

It appears, that the calcination of antimony with bone-aſhes is much more ſpeedy than when by itſelf, but the degree of fire was a little greater in the laſt experiment than in that with antimony alone. Conſidering the nature of theſe experiments, perhaps, it may be more reaſonable to impute the 3½ *per cent.* greater loſs in this laſt experiment than the ſum of the loſs in Exp. 1. and 2. to the greater inſenſible ſublimation of the calx from more fire in one caſe than in the other, than to refer it to the larger quantity of air combined with the metal in the former of theſe two laſt experiments.

EXP. IV. The above light clay or aſh-coloured powder, obtained in the laſt experiment by calcining together antimony and bone, being expoſed to various degrees of fire from 20° to 165° of WEDGWOOD's pyrometer, in cloſe crucibles, was not at all increaſed in weight, but generally loſt about 5 *per cent.* when a pretty large quantity, as a pound, was in the veſſel. A part of this loſs muſt be referred to the adheſion or vitrification of the charge with the ſides of the crucible, and part to the deficiency of the bone itſelf, as above ſhewn, by further expoſure to fire. I am ſenſible, that in experiments of this nature all calculation muſt neceſſarily, to a certain degree,

be vague; yet it may be of ſome application to obſerve, that the proportion of antimonial calx, eſtimated to be contained in LILE's Powder or *Pulvis antimonialis*, and JAMES's Powder, p. 343. from the analyſis of them, does not differ more conſiderably from the proportion of this calx than may, perhaps, be reaſonably expected on calculation from theſe four laſt experiments to exiſt in them: for 70½ parts of antimonial calx, p. 352. to 54½ parts of bone-aſhes, p. 351. is as about 56,4 parts of this calx to 43,6 parts of calcined bone; and, on analyſis, JAMES's Powder afforded $\frac{57}{100}$ of antimonial calx, and $\frac{43}{100}$ of phoſphorated lime, or nearly ſo, p. 343. allowing for the waſte.

EXP. V. This experiment ſhews the degree of fire neceſſary to render the antimony calcined with bone of a white colour; and that this whiteneſs does not depend on the air, but on the fire.

(*a*) 1500 grains of the calcined mixture of antimony and bone, Exp. 3. were kept red hot in a cloſe veſſel for half an hour. On cooling, I found the powder changed from a cineritious or clay colour to a whitiſh colour with a ſhade of yellow. The ſides of the crucible were not glazed. The pyrometer in the middle of the powder had contracted to 40°. This powder was much inferior in whiteneſs to JAMES's Powder, being much yellower.

(*b*) Another parcel of the ſame powder, Exp. 3. was expoſed in the ſame manner, but to a greater degree of fire, in which the crucible was almoſt white hot for half an hour. After cooling, the powder was found changed to a looſely cohering, ſnow-white, heavy maſs, and the ſides of the crucible were covered with a yellow glaze. This maſs, which

was

was eaſily detached from the veſſel, was found covered with a yellow vitreous coat over the whole ſurface of it that had been in contact with the crucible. In the white ſolid, on breaking it, many argentine *ſpicula* were ſeen. The pyrometer uſed in all theſe experiments indicated 71°.

(*c*) 1500 grains of the ſame parcel, Exp. 3. were expoſed in an open crucible to the fire of a melting furnace; no fumes aroſe till the crucible began to be almoſt white hot. After inverting another crucible, with a ſmall hole in its bottom, the fumes continued to aſcend at times through the aperture for a quarter of an hour. The crucible was then taken out of the fire, and on cooling a *whitiſh powder* was found, but no glazing, and the pyrometer indicated 28°. On again expoſing this crucible with one inverted over it in the melting furnace, but to a greater degree of fire, ſtill more fumes aroſe; but, on cooling, the charge was ſtill in the ſtate of a powder, though *whiter* than before; and the inſide of the inverted crucible was covered with ſilvery particles, and the hole of it was ſurrounded with argentine *ſpicula*, in a ſtellated form. The pyrometer indicated 39°. On reducing a little of this powder to a greater degree of fineneſs, it was as white as JAMES's Powder, with a yellowiſh caſt like it, but inferior in whiteneſs to a ſpecimen of *Pulvis antimonialis.* This crucible, containing its charge, with a cover cloſely luted on it, was put again into the fire, which was raiſed much higher than before; and, after being expoſed in it twenty minutes, the powder in the crucible became a looſely cohering ſolid, as *white as ſnow*, with a vitreous yellow coat, as before obſerved; the inſide of the crucible was glazed and covered with *ſpicula.* The pyrometer-piece in the middle of the powder was alſo covered with a yellow coat, but not

glazed, and it indicated 81°. This loosely cohering solid being pulverized afforded a *whiter* powder than JAMES's Powder.

(*d*) The crucible, with its charge (*b*), having a cover well luted on it, was again put into the furnace, and the fire raised to almost as great a degree as I was able. This intense heat was kept up above an hour. After cooling, a white hard solid mass was found within the crucible. On breaking the vessel, to detach from it the charge, this solid mass was found as hard as marble, and to have received its figure from the crucible. Its surface was covered with a yellow vitreous coat, and the whole inside of the vessel had a beautiful gold-coloured glaze with many argentine *spicula*. The pyrometer-piece in the middle of the charge was also covered with a fine yellow glaze, and indicated 166°. This solid, hard mass weighed only 21 grains less than before the experiment, though the whole inside of the crucible was glazed, and had shining *spicula* upon it. A piece of this hard mass being pulverized, it afforded a whiter powder than JAMES's powder is in general.

EXP. VI. 2000 grains of coarsely powdered antimony, mixed with 1105 grains of calcined hart's horn in powder, were calcined first in an open vessel, and then exposed to a great degree of fire in a close vessel, as in the above experiments with bone-shavings, Exp. 3. and 4. The calcination of this mixture in the open vessel afforded 2550 * grains, of a less whitish and rather yellowish powder, instead of a light ash-colour, as with bone shavings, Exp. 3. p. 353.; and by the second, and even

* In another experiment of this kind 2400 grains of antimony and 1500 grains of calcined bone afforded 3450 grains of yellowish light-brown powder. In a third trial, 600 grains of antimony and 400 grains of calcined bone gave 850 grains of yellowish brown powder.

repeated

repeated expoſure to fire, it never could be made quite ſo white, but ſeemed more inclined to melt than the powder prepared with unburnt bone. In other reſpects the effects of fire were apparently the ſame, or nearly ſo, as in the experiments with bone ſhavings, Exp. 3, 4.; for though the loſs of weight in this experiment, reckoning that of the antimony at 29½ *per cent.*, and that of the bone aſhes at 2½ *per cent.* ſhould have left 2483 only, inſtead of 2550; yet, in other ſimilar experiments, the product correſponded nearer to this calculation, and the loſs was ſometimes leſs both of the antimony and bone calcined ſeparately. Some of the perſons who prepare the *Pulvis antimonialis* ſay, that the whiteſt colour is obtained by firſt boiling the bone ſhavings to diſſolve their mucilage, and then calcining them with antimony as above ſhewn. Mr. LILE's receipt directs previous decoction of the hart's horn.

It will not be difficult, from theſe experiments, to give a probable reaſon for the JAMES's Powder being generally of a yellowiſh caſt, and for different parcels of it, as well as of the *Pulvis antimonialis*, being generally of different degrees of whiteneſs and ſhades of yellow. The colour of this preparation is, however, a very delicate one. I once directed a perſon to calcine together antimony and bone ſhavings, in the uſual manner, to that ſtate in which the white powder may be produced by a due degree of fire; but, inſtead of a ſnow-white maſs, I could not by any degree of fire obtain any colour but a dirty whitiſh or light ſtone colour; though repeated calcinations were employed. The reaſon of the failure was, that the earthen diſh had been broken during the calcination, and a few very ſmall pieces of it had ſcaled off, and being mixed with the powder occaſioned this diſappointment with

with reſpect to colour. The ſame diſappointment has been alſo occaſioned by uſing a ruſty iron rod in calcining the mixture.

The bone-aſhes procured from the ſal ammoniac and ſpirit of hart's horn manufactories, frequently failed in producing a white powder; and ſo did ſometimes the bone-aſhes, called prepared hart's horn, ſold by the druggiſts. Even after a fine white coloured maſs had been made, if it was pulverized in an iron mortar that had extremely little calx upon its ſurface, or dirt, the powder was not white.

The yellow coat and glaze on the ſides of the crucible and ſurface of the calcined mixture of bone and antimony, in theſe experiments, is to be aſcribed rather to the fuſion of the clay of the crucible with the antimonial calx, than to the greater degree of fire in the part of the crucible in which it takes place; or than to the calx of iron and ſiliceous earth of the veſſel: becauſe the ſame yellow coat and glazing are produced on the WEDGWOOD pyrometer pieces, which are placed in the middle of the charge, and where the degree of heat cannot be ſo great as nearer the ſide of the crucible, and yet a ſnow-white maſs is produced between theſe clay pieces and the ſides of the crucible. This effect of clay, in forming a yellow coat and glaze, is ſhewn by the obſervation of what happens when the calcined mixture is put into a WEDGWOOD's crucible, which is made of much purer clay than other veſſels of this kind, and when it is ſet in a larger Heſſian crucible with the ſpace betwixt the two veſſels filled with the ſame calcined mixture. After expoſure to a ſufficient degree of fire, *viz.* about 120° of WEDGWOOD's ſcale, the inſide and outſide of the inner crucible will be covered with a yellow vitreous coat and glaze as well as the inſide of the outer crucible in contact with the charge, while the reſt of the matter within theſe veſſels is of a ſnowy

a ſnowy whiteneſs. This yellow coat is one reaſon for the powder being of a ſhade of yellow in ſome ſpecimens.

Suppoſing the fuſibility of the antimonial calces to be diminiſhed the more they are calcined; the following experiment ſhews, that the antimonial calx in JAMES's Powder is more calcined than that in Exp. 2.

EXP. VII. 70½ grains of calcined antimony, as prepared in Exp. 2. triturated with 53½ grains of calcined bone, formed a powder of a bluiſh caſt, which being expoſed in a cloſe crucible, for half an hour, in a melting furnace, the degree of fire in which was 120° of WEDGWOOD's ſcale, it was found melted into a vitreous, pale bluiſh maſs; and the inſide of the crucible was glazed yellow, with red ſtreaks, and had argentine *ſpicula* adhering to it.

EXP. VIII. 800 grains of the calcined antimony of Exp. 2. were calcined for eight hours in a diſh, as in making LILE's Powder, by ſtirring it conſtantly, and keeping the bottom of the veſſel red hot during the whole time; the two laſt hours alſo the whole of the powder was kept red hot. On cooling, this calx was an impalpable light-brown powder.

(*a*) 100 grains of this calx, triturated with an equal quantity of calcined hart's horn, formed a powder very unlike JAMES's Powder, for it was of a light-brown colour. On expoſing it to about 120° of fire it melted into a yellow opaque maſs.

(*b*) The remaining 700 grains of the calcined antimony of this experiment were expoſed to fire and air as before for eight hours longer, and kept red hot a great part of the time; but the calx became very little lighter coloured than before.

(*c*) 100 grains of this calx laſt mentioned (*b*), triturated with as much calcined hart's horn, being expoſed to the degree of fire uſually applied in making the *Pulvis Antimonialis*, in a

cloſe veſſel, the mixture melted partially into a greyiſh maſs.

(*d*) 150 grains of the calcined antimony (*b*) of this experiment were mixed with an equal weight of calcined hart's horn. This mixture was raked about in an earthen diſh for an hour, during a great part of which time it was red hot. On cooling, the powder was evidently lighter coloured than before this calcination. It was then expoſed in a cloſe crucible to a white heat for half an hour; and, after cooling, a looſely cohering white ſolid, with a vitreous yellow coat, was found, little inferior in whiteneſs, and otherwiſe reſembling JAMES's Powder.

(*e*) 300 grains of the calcined antimony (*b*) of this experiment were raked about in an earthen diſh for an hour, a great part of which time they were kept red hot. On cooling, the calx was found of the ſame colour as before; and after expoſing it in a cloſe crucible in the melting furnace to almoſt a white heat for half an hour, it was obſerved to have been melted into a yellowiſh maſs.

It ſeems at leaſt very probable, from this experiment, that no degree or duration of fire, applied in open or cloſe veſſels to antimony *alone*, can produce a calx of the ſame kind as that in JAMES's Powder: nor, perhaps, can ſuch a powder be compoſed by fire applied, in cloſe veſſels, to calx of antimony mixed with calcined bone; but if antimony duly calcined be mixed with calcined bone, and expoſed to air, in a due degree of fire, for a ſufficient length of time, and then a ſtill greater degree of fire be applied to it in cloſe veſſels, ſuch a compound may be formed as JAMES's Powder. This experiment alſo proves, that the ſulphur in antimony is no ways neceſſary to the formation of this compound.

The manner in which air and fire act upon the antimonial calx and phoſphorated lime, I ſhall venture to conjecture.

It is probable, that the calx of antimony and phoſphorated lime combine with each other. 1. Becauſe it requires the application of heat and air for a ſhorter ſpace of time to ſeparate the ſulphur from a given quantity of antimony mixed with bone-aſhes than to produce this effect on antimony *by itſelf*: nor can the ſpeedy calcination of antimony with bone-aſhes be explained by ſuppoſing that the antimony can then bear more heat without melting; for the difference in the degree of heat applied in the two caſes is not, apparently, ſufficient to account for the difference of the times required for deſulphurating the antimony. 2. Becauſe it appears, that heat, applied to antimony in a conſiderable variety of degrees, and air for various ſpaces of time, formed a calx very different in colour, fuſibility, and other chemical qualities, from that produced by calcining this metallic ſubſtance with bone-aſhes. The ſtrongeſt confirmation, perhaps, of the opinion that the antimonial calx and phoſphorated lime are chemically united together is, that, however long the calcination of the antimony and bone-aſhes is continued in the open veſſel it will only produce preciſely the ſame ſubſtance, with reſpect to chemical properties, that is produced the moment the ſulphureous fumes ceaſe.

But why is a ſnow-white powder produced by expoſing a mixture of calcined antimony and bone-aſhes to air and fire for a due length of time, and then applying a greater degree of fire in cloſe veſſels, whereas no ſuch white powder is formed by a mixture of any calx of antimony and bone-aſhes, expoſed to any degree of fire in cloſe veſſels, without previous expoſure to fire and air? The reaſon may be, that in order that the calx ſhould unite with the phoſphorated lime, it muſt be calcined to one

certain degree; which is effected by expoſure to air and fire with the bone-aſhes when it can part or combine with air, ſo as to be reduced to that ſtate in which it will be duly calcined for union with that ſubſtance, which could not happen in cloſe veſſels.

If it be objected, that this explanation does not account for the *whiteneſs* of this preparation, which is only produced by a white heat, and to which air is not neceſſary, the difficulty will be removed by conſidering that this whiteneſs may be induced without any chemical alteration effected by the fire: for, after the firſt calcination in the open veſſel, it ſeems to act, principally, in the ſame way that it does in making grey coloured bone-aſhes, or imperfectly burnt bone, of a ſnowy whiteneſs, namely, by totally deſtroying matter extraneous to the phoſphoric ſelenite. Fire alſo, in many inſtances, alters the colour of bodies without occaſioning any change in their compoſition; and, perhaps, the change of the light clay or cineritious powder, formed by the calcination of antimony and bone-aſhes in open veſſels, to a ſnowy-white ſubſtance by further expoſure to fire, depends in part upon its increaſe of ſpecific gravity or other mechanical effects of fire. A ſtriking example of the power of fire to change the colour of bodies, by merely increaſing their ſpecific gravity, is afforded by the operation of quartation, in which proceſs, the ſilver being parted, the gold is left of the colour of copper; but, by expoſure to a due degree of fire, it is changed to its well known yellow colour, without undergoing any alteration except an increaſe of ſpecific gravity.

To elucidate the nature of the inſoluble and infuſible part of JAMES's Powder, I made the following experiments, in which I particularly had in view to determine whether ſeveral antimonial calces be wholly ſoluble in acids.

EXP.

EXP. IX. (*a*) Needle-like cryſtals of Algaroth powder diſſolved readily and totally in about thirty times their weight of marine acid.

(*b*) Part of the ſame parcel of cryſtallized Algaroth powder was calcined for above two hours, during which time it was expoſed to as great a heat as it would bear without melting, and during which time it was conſtantly raked about. Nearly half of this calcined calx readily diſſolved in marine acid, and by boiling the remainder in a proportionally much greater quantity of the ſame acid, great part of it was diſſolved, and the ſmall part which ſtill reſiſted ſolution could not be diſſolved in above 100 times its quantity of hot *aqua regia*. This indiſſoluble part afforded regulus with tartar by means of heat applied with the blow-pipe.

(*c*) White flowers of antimony generally left a reſiduum that was either inſoluble, or diſſolved with great difficulty, and in a ſmall proportion, in marine acid or *aqua regia*; yet this reſiduum was reducible. Some parcels of this calx *totally* diſſolved.

(*d*) A little of the antimony, long calcined in a former experiment, and afterwards melted into a yellow maſs, Exp. 8. (*a*), would only partially diſſolve in marine acid and *aqua regia*; but the copious reſiduum it left was reduced.

(*e*) Equal weights of cryſtals of Algaroth powder and calcined bone mixed together, diſſolved *totally* and readily in marine acid. This ſhews, that diſengaged phoſphoric acid does not precipitate antimonial calx when marine acid is preſent.

(*f*) The *calx antimonii nitrata* of the Edinburgh Diſpenſatory, argentine flowers of antimony, hyacinthine glaſs of antimony, and calx precipitated from antimonial tartar by alkali of tartar, all diſſolved readily and *wholly* in marine acid; but,

(*g*) Diaphoretic antimony left a reſiduum which mixed with tartar formed metallic grains under the flame applied by means of the blow-pipe.

(*b*) Any of the above ſoluble antimonial calces by further calcination with air and fire become more difficultly ſoluble, or partly indiſſoluble.

The next experiments were made principally for the purpoſe of knowing whether antimony calcined with vitriolic ſelenite, calcareous earth, and ſiliceous earth, would afford the ſame ſort of calx as antimony calcined with bone-aſhes.

EXP. X. 1500 grains of well burnt and dry plaſter of Paris, mixed with as much pulverized antimony, were calcined together in the ſame manner as the mixture for making LILE's Powder, Exp. 3. In half an hour the ſulphureous fumes diſappeared; after calcining half an hour longer in a heat that kept the bottom of the diſh red hot, the mixture was of a reddiſh brown or copper colour, and after cooling weighed 2520 grains. Suppoſing, therefore, the whole deficiency of weight in this experiment to be from the ſulphur carried off; and ſuppoſing the quantity of air combined with the metal to be the ſame as in Exp. 2. the loſs of weight *viz.* 32 *per cent.* is more than would have been expected; but as in experiments of this nature it is not perhaps poſſible to repeat them under preciſely the ſame circumſtances, the difference of 2½ *per cent.* deficiency more than would have been calculated, may more reaſonably be aſcribed to the ſublimation of antimony than to other cauſes. By expoſure to 70° of fire in a cloſe crucible, this calcined mixture changed to a pale ſtraw-coloured powder, and the ſides of the veſſel were glazed yellow. The change of colour was the ſame in an open veſſel in 60° of fire.

Though

Though it is probable, from this experiment, that there is an affinity between antimonial calx and vitriolic ſelenite, it is plain that the compound is very different from JAMES's Powder.

The next experiment with chalk and antimony, which Dr. BLAGDEN had the goodneſs to ſuggeſt, would lead to ſeveral concluſions, but I ſhall only take notice of the compoſition produced.

EXP. XI. 1200 grains of antimony were mixed with 800 grains of well waſhed, dried, and pulverized chalk, and calcined as in making LILE's Powder. In leſs than an hour the ſmell of ſulphur diſappeared; after which the mixture was calcined half an hour longer. It afforded a lighter clay-coloured powder than the calcination of antimony with bone-aſhes; and weighed 1800 grains. By expoſure to 100° of fire this powder changed to a dirty white colour. On examination, inſtead of aërated lime or chalk, there was found vitriolic ſelenite, part of which was probably combined with the antimonial calx; for, by means of boiling water repeatedly applied till the lixivium did not become turbid with muriated barytes nor acid of ſugar, I could only obtain 12 *per cent.* of vitriolic ſelenite, mixed with a little antimonial calx; but by means of nitrous acid I ſeparated 45 *per cent.* of this ſelenite, with ſcarcely any antimonial calx in it. The reſiduum, after this ſolution in nitrous acid, was calx of antimony with a little vitriolic ſelenite ſeemingly vitrified. Accordingly the compoſition may be ſtated to conſiſt of 1000 parts of antimonial calx and 950 parts of vitriolic ſelenite which I infer from the quantity of ſelenite diſſolved by the nitrous acid, and eſtimated to remain united to the calx; and from the following calculation of the proportion of theſe two ingredients formed in the experiment.

Antimony.	Sulphur.	Air.		
1200	– 300	+ 100	= 1000	Antimonial calx.
Calcar. earth.	Aerial acid.	Vitriolic acid.		
800	– 300	+ 450.	= 950	Vitriolic ſelenite.
			1950	
Loſs by ſublimation and waſte			150	
			1800	

With regard to the nature of this calx, the greateſt part of it readily diſſolved in marine acid; and part of what then remained was alſo diſſolved, but with great difficulty and very ſparingly; a minute quantity reſiſted ſolution entirely.

EXP. XII. 600 grains of coarſely powdered antimony were mixed with 400 grains of purified white ſand, and calcined as in making LILE's Powder. The ſmell of ſulphur continued for one hour and a half, and the mixture was calcined for half an hour longer. On cooling, a brown powder was obtained which weighed 820 grains, and expoſed to 100° of fire, melted into an irregularly figured, blackiſh maſs, full of cavities.

In this experiment the loſs of weight correſponds nearly to that in experiments above related, *viz.* thoſe in which the deficiency of weight after calcining antimony alone was about 29½ *per cent.* The much longer time required in this experiment for carrying off the ſulphur than in the calcinations with bone-aſhes, gypſum, and chalk, perhaps is owing to there being no affinity between antimonial calx and ſiliceous earth.

I beg leave to mention one more experiment relative to JAMES's Powder.

EXP. XIII. A medicine is ſold by F. NEWBERY, under the title of "JAMES's Powder for Horſes, Horned Cattle, Hounds, &c." It is a light clay-coloured, gritty, taſteleſs ſubſtance, in which

are

are ſeen ſmall *ſpicula*. It appears to me to be nothing more than JAMES's Powder for Fevers, or LILE's Powder above-mentioned, made by calcining antimony and bone-aſhes together in open veſſels; becauſe, 1ſt, by expoſure to a white heat in cloſe veſſels, it turns as white as JAMES's Powder. 2dly, It diſſolves partially in nitrous acid; and the remainder diſſolves partially in marine acid. The nitrous ſolution contains phoſphoric acid and calcareous earth; and the muriatic ſolution affords Algaroth powder.

From the whole of the above *analytical experiments* it appears:

1. That JAMES's Powder conſiſts of phoſphoric acid, lime, and antimonial calx; with a minute quantity of calx of iron, which is conſidered to be an accidental ſubſtance.

2. That either, theſe three eſſential ingredients are united with each other, forming a triple compound, or, phoſphorated lime is combined with the antimonial calx, compoſing a double compound in the proportion of about 57 parts of calx and 43 parts of phoſphorated lime.

3. That this antimonial calx is different from any other known calx of antimony in ſeveral of its chemical qualities. About three-fourths of it are ſoluble in marine acid, and afford Algaroth powder; and the remainder is not ſoluble in this menſtruum, and is apparently vitrified.

From the preceding *ſynthetic experiments* it appears, that by calcining together bone-aſhes, that is, phoſphorated lime, and antimony in a certain proportion, and afterwards expoſing the mixture to a white heat, a compound was formed conſiſting of antimonial calx and phoſphorated lime, in the ſame proportion, and poſſeſſing the ſame kind of chemical properties, as JAMES's Powder.

XXII. *An Account of ſome chemical Experiments on* Tabaſheer. *By* James Louis Macie, *Eſq. F. R. S.*

Read July 7, 1791.

THE Tabaſheer employed in theſe experiments was that which Dr. RUSSELL laid before the Society, as ſpecimens of this ſubſtance, the evening his Paper upon the ſubject was read *.

There were ſeven parcels.

N° 1. conſiſted of Tabaſheer extracted from the bamboo by Dr. RUSSELL himſelf.

N° 2. had been partly taken from the reed in Dr. RUSSELL's preſence, and partly brought to him at different times by a perſon who worked in bamboos.

N° 3. was the Tabaſheer from Hydrabad; the fineſt kind of this ſubſtance to be bought.

N°s 4, 5, and 6. all came from Maſulapatam, where they are ſold at a very low price. Theſe three kinds have been thought to be artificial compoſitions in imitation of the true Tabaſheer, and to be made of calcined bones.

N° 7. had no account affixed to it.

The Tabaſheer from Hydrabad being in the greateſt quantity, and appearing the moſt homogeneous and pure, the experiments were begun, and principally made, with it.

* See Phil. Tranſ. Vol. LXXX. p. 283.

Hydrabad

Hydrabad Tabaſheer. (N° 3.)

§ I. (A) This, in its general appearance, very much reſembled fragments of that variety of calcedony which is known to mineralogiſts by the name of *Cacholong*. Some pieces were quite opaque, and abſolutely white; but others poſſeſſed a ſmall degree of tranſparency, and had a bluiſh caſt. The latter, held before a lighted candle, appeared very pellucid, and of a flame colour.

The pieces were of various ſizes; the largeſt of them did not exceed two or three tenths of an inch cubic. Their ſhape was quite irregular; ſome of them bore impreſſions of the inner part of the bamboo againſt which they were formed.

(B) This Tabaſheer could not be broken by preſſure between the fingers; but by the teeth it was eaſily reduced to powder. On firſt chewing it felt gritty, but ſoon ground to impalpable particles.

(C) Applied to the tongue, it adhered to it by capillary attraction.

(D) It had a diſagreeable earthy taſte, ſomething like that of magneſia.

(E) No light was produced either by cutting it with a knife, or by rubbing two pieces of it together, in the dark; but a bit of this ſubſtance, being laid on a hot iron, ſoon appeared ſurrounded with a feeble luminous *auréole*. By being made red hot, it was deprived of this property of ſhining when gently heated; but recovered it again, on being kept for two months.

(F) Examined with the microſcope, it did not appear different from what it does to the naked eye.

(G) A quantity of this Tabaſheer which weighed 75.7 gr. in air, weighed only 41.1 gr. in diſtilled water whoſe temperature was 52.5 F. which makes its ſpecific gravity to be very nearly = 2.188.

Mr. CAVENDISH, having tried this ſame parcel when become again quite dry, found its ſpecific gravity to be = 2.169.

Treated with water.

§ II. (A) This Tabaſheer, put into water, emitted a number of bubbles of air; the white opaque bits became tranſparent in a ſmall degree only, but the bluiſh ones nearly as much ſo as glaſs. In this ſtate the different colour produced by reflected and by tranſmitted light was very ſenſible.

(B) Four bits of this ſubſtance, weighing together, while dry and opaque, 4.1 gr., were put into diſtilled water, and let become tranſparent; being then taken out, and the unabſorbed water haſtily wiped from their ſurface, they were again weighed, and were found to equal 8.2 gr.

In the experiment § I. (G), 75.7 gr. of this ſubſtance abſorbed 69.5 gr. of diſtilled water.

(C) Four bits of Tabaſheer, weighing together 3.2 gr. were boiled for 30′ in half an ounce of diſtilled water in a Florence flaſk, which had been previouſly rinced with ſome of the ſame fluid. This water, when become cold, did not ſhew any change on the admixture of vitriolic acid, of acid of ſugar, nor of ſolutions of nitre of ſilver, or of cryſtals of ſoda; yet, on its evaporation, it left a white film on the glaſs, which could not be got off by waſhing in cold water, nor by hot

marine

marine acid; but which was diſcharged by warm cauſtic vegetable alkali, and by long ebullition in water.

Upon theſe bits of Tabaſheer, another half ounce of diſtilled water was poured, and again boiled for about half an hour. This water alſo on evaporation left a white film on the glaſs veſſel, ſimilar to the above. The pieces of Tabaſheer having been dried, by expoſure to the air for ſome days in a warm room, were found to have loſt one-tenth of a grain of their weight.

To aſcertain whether the whole of a piece of Tabaſheer could be diſſolved by boiling in water, a little bit of this ſubſtance, weighing three-tenths of a grain, was boiled in 36 ounces of ſoft water for near five hours conſecutively; but being afterwards dried and weighed, it was not diminiſhed in quantity, nor was it deprived of its taſte.

With vegetable colours.

§ III. Some Tabaſheer, reduced to fine powder, was boiled for a conſiderable time in infuſions of turnſole, of logwood, and of dried red cabbage, but produced not the leaſt change in any one of them.

At the fire.

§ IV. (A) A piece of this Tabaſheer, thrown into a red hot crucible, did not burn or grow black. Kept red hot for ſome time, it underwent no viſible change; but when cold, it was harder, and had entirely loſt its taſte. Put into water it

grew tranſparent, juſt as it would have done, had it not been ignited.

(B) 6.4 gr. of this ſubſtance, made red hot in a crucible, were found, upon being weighed as ſoon as cold, to have loſt two-tenths of a grain. This loſs appears to have ariſen merely from the expulſion of interpoſed moiſture; for theſe heated pieces, on being expoſed to the air for ſome days, recovered exactly their former weight.

(C) A bit of this ſubſtance was put into an earthen crucible, ſurrounded with ſand, and kept red hot for ſome time; when cold, it was ſtill white, both exteriorly and interiorly.

(D) Thrown into ſome melted red hot nitre, this ſubſtance did not produce any deflagration, or ſeem to ſuffer any alteration.

(E) A bit expoſed on charcoal to the flame of the blow-pipe did not decrepitate or change colour; when firſt heated it diffuſed a pleaſant ſmell; then contracted very conſiderably in bulk, and became tranſparent; but on continuing the heat it again grew white and opaque, but ſeemed not to ſhew any inclination to melt *per ſe*. Poſſibly, however, it may ſuffer ſuch a ſemi-fuſion, or ſoftening of the whole maſs, as takes place in clay when expoſed to an intenſe heat; for when the bit uſed happened to have cracks, it ſeparated during its contraction, at theſe cracks, and the parts receded from each other without falling aſunder.

If, while the bit of Tabaſheer was expoſed to the flame, any of the aſhes of the coal fell upon it, it inſtantly melted, and ſmall very fluid bubbles were produced. That the opacity which this ſubſtance acquires on continuing to heat it after it is become tranſparent, is not owing to the fuſion of its

ſurface by means of ſome of the aſhes of the charcoal ſettling upon it unobſerved, appeared by its undergoing the ſame change when fixed to the end of a glaſs tube, in the method of M. DE SAUSSURE *.

With acids.

§ V. (A) A piece of Tabaſheer, weighing 1.2 gr. was firſt let ſaturate itſelf with diſtilled water; its ſurface being then wiped dry, it was put into a matraſs with ſome pure white marine acid, whoſe ſpecific gravity was 1.13. No effervescence aroſe on its immerſion into the acid; nor did this menſtruum, even by ebullition, ſeem to have any action upon it, or itſelf receive any colour. The acid being evaporated left only ſome dark coloured ſpots on the glaſs. Theſe ſpots were diſſolved by diſtilled water. No precipitation was produced in this water by vitriolic acid, or by a ſolution of cryſtals of ſoda. The bit of Tabaſheer waſhed with water, and made red hot, had not ſuſtained any loſs of weight.

The pores of the maſs of Tabaſheer were filled with water before it was put into the acid, to expel the common air contained in them, and which would have made it impoſſible to aſcertain with accuracy whether any effervescence was produced on its firſt contact with the menſtruum.

(B) Another portion of Tabaſheer, weighing 10.2 gr. was boiled in ſome of the ſame marine acid. Not the leaſt precipitate was produced on ſaturating this acid with ſolution of mild ſoda. This Tabaſheer alſo, after having been

* Journal de Phyſique, Tom. XXVI. p. 409.

boiled

boiled in water, and dried by expoſure for ſome days to the air, was ſtill of its former weight.

§ VI. This ſubſtance ſeemed in like manner to reſiſt the action of pure white nitrous acid boiled upon it.

§ VII. (A) A bit of Tabaſheer weighing 0.6 gr. was digeſted in ſome ſtrong white vitriolic acid, which had been made perfectly pure by diſtillation. It did not ſeem by this treatment to ſuffer any change, and after having been freed from all adhering vitriolic acid by boiling in water, it had not undergone any alteration either in its weight or properties. The vitriolic acid afforded no precipitate on being ſaturated with ſoda.

(B) Two grains of Tabaſheer reduced to fine powder were made into a paſte with ſome of this ſame vitriolic acid, and this mixture was heated till nearly dry; it was then digeſted in diſtilled water. This water, being filtered, taſted ſlightly acid, did not produce the leaſt turbidneſs with ſolution of ſoda, and ſome of it, evaporated, left only a faint black ſtain on the glaſs, produced doubtleſs by the action of the vitriolic acid on a little vegetable matter, which it had received either from the Tabaſheer, or from the paper. The undiſſolved matter collected, waſhed, and dried, weighed 1.9 gr.

§ VIII. 2 gr. of Tabaſheer, reduced to fine powder, were long digeſted in a conſiderable quantity of liquid acid of ſugar. The taſte of the liquor was not altered; and being ſaturated with a ſolution of cryſtals of ſoda in diſtilled water, it did not afford any precipitate. The Tabaſheer having been freed from all adhering acid, by very careful ablution with diſtilled water, and let dry in the air, was totally unchanged in its appearance, and weighed 1.98 gr. This Tabaſheer being gra-

dually heated till red hot, did not become in the leaſt black, or loſe much of its weight, a proof that no acid of ſugar had fixed in it.

With liquid alkalies.

§ IX. (A) Some liquid cauſtic vegetable alkali being heated in a phial, Tabaſheer was added to it, which diſſolved very readily, and in conſiderable quantity. When the alkali would not take up any more, it was ſet by to cool, but was not found next morning to have cryſtallized, or undergone any change, though it had become very concentrated, during the boiling, by the evaporation of much of the water.

(B) This ſolution had an alkaline taſte, but ſeemingly with little, if any, cauſticity.

(C) A drop of it changed to green a watery tincture of dried red cabbage.

(D) Some of this ſolution was expoſed in a ſhallow glaſs to ſpontaneous evaporation in a warm room. At the end of a day or two it was converted into a firm, milky jelly. After a few days more, this jelly was become whiter, more opaque, and had dried and cracked into ſeveral pieces, and finally it became quite dry, and curled up and ſeparated from the glaſs.

The ſame change took place when the ſolution had been diluted with ſeveral times its bulk of diſtilled water, only the jelly was much thinner, and dried into a white powder.

Some of this ſolution, kept for many weeks in a bottle cloſely ſtopped, did not become a jelly, or undergo any change.

(E) A

(E) A ſmall quantity of this ſolution was let fall into a proportionably large quantity of ſpirit of wine, whoſe ſpecific gravity was .838. The mixture immediately became turbid, and, on ſtanding, a denſe fluid ſettled to the bottom, and which, when the bottle was haſtily inverted, fell through the ſpirit of wine in round drops, like a ponderous oil.

The ſupernatant ſpirit of wine being carefully decanted off, ſome diſtilled water was added to this thick fluid, by which it was wholly diſſolved. This ſolution, expoſed to the air, ſhewed phænomena exactly ſimilar to thoſe of the undiluted ſolution (D).

The decanted ſpirit being alſo left expoſed to the air in a ſhallow glaſs veſſel, did not, after many days, either depoſit a ſenſible quantity of precipitate, or become gelatinous; but having evaporated nearly away, left a few drops of a liquor which made infuſion of red cabbage green; and, on the addition of ſome pure marine acid, effervеſced violently. No precipitate fell during this ſaturation with the acid; nor did the mixture on ſtanding become a jelly; and on the total evaporation of the fluid part, a ſmall quantity of muriate of tartar only remained. The ſpirit of wine ſeems, therefore, to have diſſolved merely a portion of ſuperabundant alkali preſent in the mixture, but none of that united with Tabaſheer.

(F) To different portions of this ſolution were added ſome pure marine acid, ſome pure white vitriolic acid, and ſome diſtilled vinegar, each in exceſs. Theſe acids at firſt produced neither heat, effervеſcence, any precipitate, or the leaſt ſenſible effect, except the vitriolic acid, which threw down a very ſmall quantity of a white matter; but, after ſtanding ſome days, theſe mixtures changed into jellies ſo firm, that

the

the glaſſes containing them were inverted without their falling out.

This change into jelly equally took place whether the mixtures were kept in open or cloſed veſſels, were expoſed to the light, or ſecluded from it; nor did it ſeem to be much promoted by boiling the mixtures.

(G) Some ſolution of mild volatile alkali in diſtilled water, being added to ſome of this ſolution, ſeemed at the firſt inſtant of mixture to have no effect upon it; but in the ſpace of a ſecond or two it occaſioned a copious white precipitate.

(H) The flakes remaining on the glaſſes at (D) and (E) put into marine acid raiſed a ſlight effervescence, but did not diſſolve. Theſe flakes, when taken out of the acid, and well waſhed, were found, like the original Tabaſheer, to be white and opaque when dry; but to become tranſparent when moiſtened, and then to ſhew the blue and flame colour, § II. (A).

(I) The jellies (F), diluted with water, and collected on a filter, appeared to be the Tabaſheer unchanged.

§ X. A bit of Tabaſheer, weighing two-tenths of a grain, was boiled in 127 gr. of ſtrong cauſtic volatile alkali for a conſiderable time; but after being made red hot, it had not ſuſtained the leaſt diminution of weight.

§ XI. (A) 27 gr. of Tabaſheer, reduced to fine powder, were put into an open tin veſſel with 100 gr. of cryſtals of ſoda, and ſome diſtilled water, and this mixture was made boil for three hours. The clear liquor was then poured off, and the Tabaſheer was digeſted in ſome pure marine acid; after ſome time this acid was decanted, and the Tabaſheer waſhed with diſtilled water, which was then added to the acid.

(B) This Tabaſheer was put back into the alkaline ſolution, which ſeemed not impaired by the foregoing proceſs, and again boiled for a conſiderable time. The liquor was then poured from it while hot, and the Tabaſheer edulcorated with ſome cold diſtilled water, which was afterwards mixed with this hot ſolution, in which it inſtantly cauſed a precipitation. On heating the mixture it became clear again; but as it cooled it changed wholly into a thin jelly; but, in the courſe of a few days, it ſeparated into two portions, the jelly ſettling in a denſer ſtate to the bottom of the veſſel, leaving a limpid liquor over it.

(C) The Tabaſheer remaining (B) was boiled in pure marine acid; the acid was then poured off, and the Tabaſheer edulcorated with ſome diſtilled water, which was afterwards mixed with the acid.

(D) The remaining Tabaſheer collected, waſhed, and dried, weighed 24 gr. and ſeemed not to be altered.

(E) The acid liquors (A and C) were mixed together, and ſaturated with ſoda, but afforded no precipitate.

(F) The alkaline mixture (B) was poured upon a filter, the clear liquor came through, leaving the jelly on the paper.

Some of this clear liquor, expoſed to the air in a ſaucer, at the end of ſome days depoſited a ſmall quantity of a gelatinous matter; after ſome days more, the whole fluid part exhaled, and the ſaucer became covered with regular cryſtals of ſoda, which afforded no precipitate during their ſolution in vitriolic acid. What had appeared like a jelly while moiſt aſſumed, on drying, the form of a white powder. This powder was inſoluble in vitriolic acid, and ſeemed ſtill to be Tabaſheer.

Some

Some of this clear liquor, mixed with marine acid, effervesced; did not afford any precipitate; but, on ſtanding ſome days, the mixture became ſlightly gelatinous.

(G) Some of the thick jelly remaining on the filter, being boiled in water and in marine acid, appeared inſoluble in both, and ſeemed to agree entirely with the above powder (F).

With dry alkalies.

§ XII. (A) Tabaſheer melted on the charcoal at the blowpipe with ſoda, with conſiderable effervescence. When the proportion of alkali was large, the Tabaſheer quickly diſſolved, and the whole ſpread on the coal, ſoaked into it, and vaniſhed; but, by adding the alkali to the bit of Tabaſheer in exceedingly ſmall quantities at a time, this ſubſtance was converted into a pearl of clear colourleſs glaſs.

(B) 5 gr. of Tabaſheer, reduced to fine powder, were melted in a platina crucible with 100 gr. of cryſtals of ſoda. The maſs obtained was white and opaque, and weighed 40.2 gr. Put into an ounce of diſtilled water, it wholly diſſolved. An exceſs of marine acid let fall into this ſolution produced an effervescence, and changed it into a jelly. This mixture was ſtirred about, and then thrown upon a filter. The jelly left on the paper did not diſſolve in marine acid by ebullition; collected, waſhed with diſtilled water, and dried, it weighed 4.5 gr. and ſeemed to be the Tabaſheer unaltered.

The liquor which had come through being ſaturated with mineral alkali yielded only a very ſmall quantity of a red precipitate, which was the colouring matter of the pink blotting paper through which it had been paſſed.

(C) 10 gr. of Tabaſheer, reduced to powder, were mixed with an equal weight of ſoda, deprived of its water of cryſtallization by heat. This mixture was put into a platina crucible, and expoſed to a ſtrong fire for 15′. It was then found converted into a tranſparent glaſs of a ſlight yellow colour. This glaſs was broken into pieces, and boiled in marine acid. No effervescence appeared; but the glaſs was diſſolved into a jelly. This jelly, collected on a filter, well waſhed, and dried, weighed 7.7 gr.

The acid liquor which came through, on ſaturation with ſoda, afforded not the leaſt precipitate; but, after ſtanding a day or two, it changed into a thin jelly. This collected on a filter was waſhed with diſtilled water, and then boiled in marine acid, but did not diſſolve. Being again edulcorated, and made red hot, it weighed 1.6 gr. The filtered liquor (B) would in all probability have changed ſimilarly to a jelly, had it been kept. Theſe precipitates were analogous to thoſe § IX. (I).

(D) An equal weight of vegetable alkali and Tabaſheer were melted together in the platina crucible. The glaſs produced was tranſparent; but it had a fiery taſte, and ſoon attracted the moiſture of the air, and diſſolved into a thick liquor. But two parts of vegetable alkali, with three of Tabaſheer, yielded a tranſparent glaſs, which was permanent.

Treated with other fluxes.

§ XIII. (A) A fragment of Tabaſheer put into glaſs of borax, and urged at the blow-pipe, contracted very conſiderably in ſize, the ſame as when heated *per ſe*; after which it continued turning about in the flux, diſſolving with great difficulty

and very ſlowly. When the ſolution was effected, the ſaline pearl remained perfectly clear and colourleſs.

(B) With phoſphoric ammoniac (made by ſaturating the acid obtained by the ſlow combuſtion of phoſphorus with cauſtic volatile alkali) the Tabaſheer very readily melted on the charcoal at the blow-pipe, with efferveſcence, into a white frothy bead.

(C) Fuſed, by the ſame means, on a plate of platina, with the vitriols of tartar and ſoda, it appeared entirely to reſiſt their action; the little particles employed continuing to revolve in the fluid globules without ſuſtaining any ſenſible diminution of ſize, and the ſaline beads on cooling aſſumed their uſual opacity.

(D) A bit of Tabaſheer was laid on a plate of ſilver, and a little litharge was put over it, and then melted with the blow-pipe. It immediately acted on the Tabaſheer, and covered it with a white glaſſy glazing. By the addition of more litharge the maſs was brought to a round bead, though with conſiderable difficulty. This bead bore melting on the charcoal, without any reduction of the lead, but could not be obtained tranſparent.

(E) The eaſe with which this ſubſtance had melted with vegetable aſhes, led to the trial of it with pure calcareous earth. A fragment of Tabaſheer, fixed to the end of a bit of glaſs, was rubbed over with ſome powdered whiting. As ſoon as expoſed to the flame of the blow-pipe, it melted with conſiderable efferveſcence; but could not, even on the charcoal, and with the addition of more whiting, be brought to a tranſparent ſtate, or reduced into a round bead.

Equal weights of Tabaſheer and pure calcareous ſpar, both reduced to fine powder, were irregularly mixed, and expoſed

in

in the platina crucible to a ſtrong fire in a forge for 20′; but did not even concrete together.

(F) When magneſia was uſed, no fuſion took place at the blow-pipe.

(G) Equal parts of Tabaſheer, whiting, and earth of alum precipitated by mild volatile alkali, were mixed in a ſtate of powder, and ſubmitted in the platina crucible to a ſtrong fire for 20′, but were afterwards found unmelted.

Examination of the other ſpecimens.

N° I.

This parcel contained particles of three kinds; ſome white, of a ſmooth texture, much reſembling the foregoing ſort; others of the ſame appearance, but yellowiſh; and others greatly ſimilar to bits of dried mould.

The white and yellowiſh pieces were ſo ſoft as to be very eaſily rubbed to powder between the fingers. They had a diſagreeable taſte, ſomething like that of rhubarb. Put into water, the white bits ſcarcely grew at all tranſparent; but the yellow ones became ſo to a conſiderable degree.

The brown earth-like pieces were harder than the above, had little taſte, floated upon water, and remained opaque.

Expoſed to the blow-pipe, they all charred and grew black; the laſt variety even burned with a flame. When the vegetable matter was conſumed, the pieces remained white, and then had exactly the appearance, and poſſeſſed all the properties, of the the foregoing Tabaſheer from Hydrabad, and like it melted with ſoda into a tranſparent glaſs.

N° II.

N° II.

Alſo conſiſted of bits of three ſorts.

(*a*) Some white, nearly opaque.

(*b*) A few ſmall very tranſparent particles, ſhewing, in an eminent degree, the blue and yellow colour, by the different direction of light.

(*c*) Coarſe, browniſh pieces of a grained texture.

Theſe all had exactly the ſame taſte, hardneſs, &c. and ſhewed the ſame effects at the blow-pipe, as N° I.

27 gr. of this Tabaſheer thrown into a red-hot crucible, burned with a yellowiſh white flame, loſt 2.9 gr. in weight, and became ſo ſimilar to the Hydrabad kind as not to be diſtinguiſhed from it.

Some of this Tabaſheer put into a crucible, not made very hot, emitted a ſmell ſomething like tobacco aſhes, but not the kind of perfume diſcovered in that from Hydrabad, § IV. (E).

N° IV.

All the pieces of this parcel were of one appearance, and a good deal reſembled, in their texture, the third variety of N° II. Their colour was white; their hardneſs ſuch as very difficultly to be broken by preſſure between the fingers. In the mouth they immediately fell to a pulpy powder, and had no taſte.

A bit expoſed on the charcoal to the blow-pipe became black, melted like ſome vegetable matters, caught flame, and burnt to a botryoid inflated coal, which ſoon entirely conſumed away, and vaniſhed.

A piece put into water fell to a powder. The mixture being boiled, this powder diſſolved, and turned the whole to a jelly.

Theſe

These properties are exactly those of common starch.

N° V.

Agreed entirely with N° IV. in appearance, properties, and nature.

N° VI.

The pieces of this parcel were white, quite opaque, and considerably hard. Their taste and effects at the blow-pipe, were perfectly similar to those of the Hydrabad kind.

N° VII.

Much resembled N° VI. only was rather softer, and seemed to blacken a little when first heated. With fluxes at the blow-pipe it shewed the same effects as all the above.

Conclusion.

1. It appears from these experiments, that all the parcels, except N° IV. and V. consisted of genuine Tabasheer; but that those kinds, immediately taken from the plant, contained a certain portion of a vegetable matter, which was wanting in the specimens procured from the shops, and which had probably been deprived of this admixture by calcination, of which operation a partial blackness, observable on some of the pieces of N° III. and VI. are doubtless the traces. This accounts also for the superior hardness and diminished tastes of these sorts.

2. The nature of this substance is very different from what might have been expected in the product of a vegetable. Its indestructibility by fire; its total resistance to acids; its uniting by

by fuſion with alkalies in certain proportions into a white opaque maſs, in others into a tranſparent permanent glaſs; and its being again ſeparable from theſe compounds, entirely unchanged by acids, &c. ſeem to afford the ſtrongeſt reaſons to conſider it as perfectly identical with common *ſiliceous earth*.

Yet from pure quartz it may be thought to differ in ſome material particulars; ſuch as in its fuſing with calcareous earth, in ſome of its effects with liquid alkalies, in its taſte, and its ſpecific gravity.

But its taſte may ariſe merely from its divided ſtate, for chalk and powdery magneſia both have taſtes, and taſtes which are very ſimilar to that of pure Tabaſheer; but when theſe earths are taken in the denſer ſtate of cryſtals, they are found to be quite inſipid; ſo Tabaſheer, when made more ſolid by expoſure to a pretty ſtrong heat, is no longer perceived, when chewed, to act upon the palate, § IV. (A).

And, on accurate compariſon, its effects with liquid alkalies have not appeared peculiar; for though it was found on trial, that the powder of common flints, when boiled in ſome of the ſame liquid cauſtic alkali employed at § IX. (A), was ſcarcely at all acted upon; and that the very little which was diſſolved was ſoon precipitated again, in the form of minute *flocculi*, on expoſing the ſolution to the air, and was immediately thrown down on the admixture of an acid; yet the precipitate obtained from *liquor ſilicum* by marine acid was diſcovered, even when dry to diſſolve readily in this alkali, but while ſtill moiſt to do ſo very copiouſly, even without the aſſiſtance of heat; and ſome of this ſolution, thus ſaturated with ſiliceous matter by ebullition, being expoſed to the air in a ſhallow glaſs, became a jelly by the next day, and the day after dried, and cracked, &c. exactly like the mixtures § IX.

(D and E). And another portion of this ſolution mixed with marine acid afforded no precipitate, and remained perfectly unaffected for two days; but on the third it was converted into a firm jelly like that § IX. (F).

As gypſum is found to melt *per ſe* at the blow-pipe, though refractory to the ſtrongeſt heat that can be made in a furnace, it was thought that poſſibly ſiliceous and calcareous earths might flux together by this means, though they reſiſt the utmoſt power of common fires; but experiment ſhewed, that in this reſpect quartz did not agree with Tabaſheer. But this difference ſeems much too likely to depend on the admixture of a little foreign matter in the latter body, to admit of its being made the grounds for conſidering it as a new ſubſtance, in oppoſition to ſo many more material points in which it agrees with ſilex.

Nor can much weight be laid on the inferior ſpecific gravity of a body ſo very porous. The infuſibility of the mixture § XIII. (G) depended alſo, probably, either on an inaccuracy in the proportions of the earths to each other, or on a deficiency of heat.

3. Of the three bamboos which were not ſplit before the Royal Society I have opened two. The Tabaſheer found in them agreed entirely in its properties with that of N° I. and II.

It was obſerved, that all the Tabaſheer in the ſame joint was exactly of the ſame appearance. In one joint it was all ſimilar to the yellowiſh ſort N° I. In another joint of the ſame bamboo, it reſembled the variety (*c*) of N° II. Probably, therefore, the parcels from Dr. RUSSELL, containing each ſeveral varieties of this ſubſtance, aroſe from the produce of many joints having been mixed together.

4. The

4. The aſhes, obtained by burning the bamboo, boiled in marine acid, left a very large quantity of a whitiſh inſoluble powder, which, fuſed at the blow-pipe with ſoda, effervesced, and formed a tranſparent glaſs. Only the middle part of the joints was burned, the knots were ſawed off, leſt, being porous, Tabaſheer might be mechanically lodged in them. However, the great quantity of this remaining ſubſtance ſhews it to be an eſſential, conſtituent part of the wood.

The aſhes of common charcoal, digeſted in marine acid, left in the ſame manner an inſoluble reſiduum which fuſed with ſoda with effervescence, and formed glaſs; but the proportion of this matter to the aſhes was greatly leſs than in the foregoing caſe.

5. Since the above experiments were made, a ſingular circumſtance has preſented itſelf. A green bamboo, cut in the hot-houſe of Dr. PITCAIRN, at Iſlington, was judged to contain Tabaſheer in one of its joints, from a rattling noiſe diſcoverable on ſhaking it; but being ſplit by Sir JOSEPH BANKS, it was found to contain, not ordinary Tabaſheer, but a ſolid pebble, about the ſize of half a pea.

Externally this pebble was of an irregular rounded form, of a dark-brown or black colour. Internally it was reddiſh-brown, of a cloſe dull texture, much like ſome martial ſiliceous ſtones. In one corner there were ſhining particles, which appeared to be cryſtals, but too minute to be diſtinguiſhed even with the microſcope.

This ſubſtance was ſo hard as to cut glaſs!

A fragment of it expoſed to the blow-pipe on the charcoal did not grow white, contract in ſize, melt, or undergo any change. Put into borax it did not diſſolve, but loſt its colour,

and tinged the flux green. With ſoda it effervefced, and formed a round bead of opaque black glaſs.

Theſe two beads, digeſted in ſome perfectly pure and white marine acid, only partially diſſolved, and tinged this menſtruum of a greeniſh yellow colour; and from this ſolution Pruſſite of tartar, ſo pure as not, under many hours, to produce a blue colour with the above pure marine acid, inſtantly threw down a very copious Pruſſian blue.

P. S. In aſcertaining the ſpecific gravity of the Hydrabad Tabaſheer, § I. (G), great care was taken in both the experiments that every bit was thoroughly penetrated with the water, and tranſparent to its very center, before its weight in the water was determined.

XXIII. *A Second Paper on Hygrometry.*

By J. A. De Luc, *Esq. F. R. S.*

Read July 14, 1791.

PART II.

IN the first *part* of this Paper * I have treated of the fundamental principles of hygrometry, and of some hygroscopic phænomena; and this will relate to a particular application of those premises.

62. Since the publication of my first hygrometer, many others have been invented, two of which are now principally in use; the *hair* hygrometer of M. DE SAUSSURE, and my *hygrometer* made of a *slip of whalebone*. If the comparative points of those instruments could be determined in the whole extent of their scales, the only inconvenience of their being both used would be, the necessity of reducing to one of them, the observations made with the other; but from 70 to 100 of mine, which space includes the most important period of *moisture*, their correspondent indications are as different from one another, and as variable, as if they were the effects of two very different causes. Therefore it is important to decide which of them should remain our only measure of *moisture*, till, if possible, a better one is found. The following pages, I hope, will lead to that decision.

* See Page 1. of this Volume.

63. The

63. The fundamental proceſs of M. DE SAUSSURE, with the view of diſcovering the effects of *moiſture* on the *hair hygrometer*, was this. He repeatedly cauſed ſucceſſive known quantities of *water* to evaporate into a cloſe glaſs veſſel, previouſly reduced to *extreme dryneſs*, and containing that *hygrometer* and a *manometer*; he obſerved the correſpondent changes of thoſe inſtruments, and, by combining the reſults of his experiments, he reduced to regular *ſeries* the correſpondent motions of the two inſtruments by equal quantities of *evaporated water*. Having confined himſelf to that only claſs of experiments, which, from cauſes that I ſhall explain, could not diſcover to him the difficulties of his attempt, he thought himſelf warranted to draw from them the following concluſions. 1ſt, That the degrees of *moiſture* in the incloſed *medium*, were nearly proportional to the quantities of *water* evaporated in the veſſel; and that, conſequently, the *ratio* obſerved between thoſe quantities and the *march* of his *hygrometer*, could be conſidered as giving immediately the *march* of the inſtrument correſpondent to *moiſture* itſelf; which, according to our common opinion, is a certain quantity of *aqueous vapours* ſpread in the *medium*. 2dly, That when no more *water* could *evaporate* in the veſſel, the incloſed *medium* was arrived at *extreme moiſture*; and that, conſequently, the point indicated at that time on his *hygrometer*, was to be the *limit* of its *ſcale* on that ſide. 3dly, That having, from thoſe experiments, a probable determination of the *expanſions* of the *hair* by ſucceſſive equal quantities of *moiſture*, in beginning from the point were this is null, and ending at its *extreme*, his inſtrument could not differ eſſentially from an abſolute *hygrometer*.

64. Theſe concluſions were very natural in the ſtate of M. DE SAUSSURE's experiments; but before their publication I had

gone

gone over a great field of *hygroſcopic* phænomena, in which the *hair*, and a *cloſe veſſel*, had a ſhare; and thereby, ſeeing the objects in another light than M. DE SAUSSURE, I doubted of his concluſions, and I procured three of his *hygrometers*, in order to examine them on ſome particular points. It was after that immediate verification of my conjectures concerning his inſtrument, that I ſettled the following concluſions, very different from thoſe above. 1ſt, That MOISTURE, or the quantity of *vapour* ſpread in the *medium* itſelf, does not increaſe in an incloſed ſpace in proportion to the quantity of water evaporated in it; becauſe of an increaſing, but undetermined, part of that *water* being depoſited on the ſides of the veſſel; and that, conſequently, Mr. DE SAUSSURE's experiments could not afford the determination of a real *hygroſcopic-ſcale*. 2dly, That the circumſtance conſidered by him as a ſure ſign of *extreme moiſture* exiſting in the incloſed *medium*, namely, the *maximum* of *evaporation* in the ſpace, has only that effect when the *temperature* is very little above 32°; but that, by ſucceſſive increaſes of *heat* from that point, *moiſture* recedes farther and farther from its *extreme*; or from the point where no more *vapour* can be introduced in the *medium* without an immediate *precipitation*; though at the ſame time, there are ſucceſſive increaſes in the quantity of *vapour*, and thereby a conſtant *maximum* of *evaporation* correſpondent with the actual *temperature*. 3dly, That, in approaching to *extreme moiſture*, the *hair* hygrometer becomes *ſtationary*, and afterwards a little *retrograde*, in which *march* the unavoidable irregularities of every *hygroſcopic* ſubſtance produce frequent anomalies; from which cauſe it was very difficult for M. DE SAUSSURE, conſidering the form of his experiments, to diſcover the *hygroſcopic law* expreſſed by the ſecond concluſion; and with the unknown exiſtence of that

law,

law, to ſuſpect the *march* of his *hygrometer*: which accidental complication I ſhall explain hereafter.

65. When I publiſhed thoſe reſults of my experiments and obſervations, M. DE SAUSSURE rejected them; not from having made new experiments that had confirmed his opinions; but becauſe he conjectured inverſely, that my theory reſulted from a fallacious *march* of my *hygrometer*: and the well-earned reputation of that celebrated philoſopher engaged me to undertake every experiment that could help me to detect on which ſide was the error. I have related, in the firſt *part* of this Paper, ſome of thoſe experiments; and now, for their application, as well as for giving an account of ſome others, I ſhall follow more particularly M. DE SAUSSURE's proceſs.

66. In a large glaſs veſſel, containing (as I have mentioned above) a *manometer* and his *hair*-hygrometer, which veſſel he had previouſly reduced to a known ſmall diſtance from *extreme dryneſs*, M. DE SAUSSURE introduced from time to time a piece of *wet* cloth, which he weighed both before he put it into the veſſel, and when he took it out. The ſucceſſive increaſes in the quantity of *vapour* reſulting from that proceſs were indicated; on the *manometer*, by ſucceſſive increaſes in the quantity of the incloſed *elaſtic fluids*, which cauſed the quickſilver to aſcend more and more in that inſtrument; and on the *hygrometer*, by ſucceſſive expanſions of the *hair*. The *maximum* of *evaporation* was clearly indicated by the *manometer*; for, during every laſting *temperature*, the quickſilver, after having aſcended to a certain point, remained fixed at that point, notwithſtanding a longer ſtay of the *wet* cloth; and by repeating that operation at different *temperatures*, M. DE SAUSSURE determined the quantities of *evaporated water* that, in a given ſpace, and by a given *temperature*, produced the *maximum* of *evaporation*.

That

That ſingle determination, from its great conſequences not yet generally attended to, would be ſufficient to fix the celebrity of its author, as I have already expreſſed many times in other works.

67. But with reſpect to a different concluſion from the ſame experiments, no leſs important to natural philoſophy, I have alſo ſaid, that M. DE SAUSSURE's *hygrometer* may have miſled him. We have ſeen, that the *manometer* indicated, by an indubitable ſymptom, the *maximum* of *evaporation*; for here the immediate cauſe is clearly known, namely, the *quantity* of *elaſtic fluids*; and it is evident, that the quickſilver muſt aſcend in the inſtrument in proportion to that *quantity*, and ſtop when it ceaſes to increaſe. But it is not the ſame as to the indications of the *hygrometer* in reſpect of *moiſture:* M. DE SAUSSURE found himſelf that they were far from proportional to the intenſities of their cauſe; and in the laſt ſtage of his experiments, though theſe indications did not vary much by the different *maxima* of *evaporation*, they however varied in the ſpace of 1 or 2 *degrees*. But as thoſe ſmall differences on the point where the *hygrometer* ſtopped in different experiments, did not follow apparently any law conformable to the *temperature*, M. DE SAUSSURE conſidered them as ſmall *anomalies*, unavoidable in hygroſcopic ſubſtances, and of little conſequence on a ſcale of 100 *degrees*; therefore, laying aſide that circumſtance, he could have no doubt, that, in every *temperature*, the *maximum* of *evaporation* in a cloſe ſpace was ſynonymous with the *maximum* of *moiſture* in that ſpace; while, from my experiments, theſe two ſuppoſed identical expreſſions may differ $\frac{1}{5}$, and ſometimes $\frac{1}{4}$, of the real ſcale of *moiſture*, which is the caſe in the *temperature* of only 75° or 80°.

68. Let us now ſuppoſe for a moment, that the above *hygroſcopic law*, and the *march* that I attribute to the *hair* hygrometer, are real. In that caſe, if, during a conſtant *maximum* of *evaporation*, the *temperature* varies from 32° to 80°, *moiſture* will diminiſh $\frac{1}{7}$ and even $\frac{1}{8}$ of the whole; or, in other words, the ſtate of the *medium* will be diſtant by ſo much from that in which a new introduction of *vapour* would be followed by a *precipitation*. But at the ſame time, in the whole of that period of *moiſture*, the *hair* hygrometer is ſuppoſed to move only 1 or 2 *degrees* backwards and forwards, with frequent irregularities. Therefore, in the hypotheſis, ſuch a great change of *moiſture* would be hardly ſuſpected from thoſe ſmall deviations of the *hair* hygrometer, in which at firſt nothing appears to be regular; and thereby it is evident, that, by confining himſelf to thoſe experiments, M. DE SAUSSURE could not diſcover thoſe two important laws of *hygrology* and *hygrometry*, of which I have here only ſuppoſed the exiſtence.

69. Let us ſuppoſe again, that the *hair* hygrometer had not exiſted before a certain number of other experiments; and that M. DE SAUSSURE, in his attempt to produce an inſtrument of that kind, had fixed on any of the *ſlips* made of *fibrous* vegetable or animal ſubſtances cut *acroſs* the *fibres*, of which many *hygroſcopes* had been made before in a coarſer manner; and that, in every other reſpect, he had proceeded as he has done with the *hair*. In that caſe, having placed his *hygrometer* and the *manometer* in the ſame veſſel, with a quantity of *water* ſufficient for producing the *maximum* of *evaporation* in every common *temperature*, and obſerved alſo the points where both inſtruments ſtopped in different laſting *temperatures*, he then would have found; that the *hygrometer* indicated leſs and leſs *moiſture*, at the ſame time that the *manometer*, by aſcending more

more and more as the *evaporation* increaſed by more *heat*, indicated that increaſe in the quantity of *vapour*, diſtinct from the expanſions of the *fluids*; and that each of thoſe inſtruments remained alſo fixed in that ſtate of oppoſite changes, by every ſufficiently durable change of *temperature*. The firſt obſervation of that phænomenon would have ſurprized him, as it did me, and it would have induced him alſo to try ſome other of thoſe *ſlips*; and by finding the ſame phænomenon with every one of them, he would have been convinced, that it was a real *law* of *moiſture*. Laſtly, if he had known Mr. JAMES WATT's obſervations on the great *dryneſs* of the *ſteam* of *boiling water*, as long as it remains in a ſpace as *warm* as the *water* that produces it (a condition always underſtood when the general laws of *evaporation* are the object of inquiry); admitting with me, as it follows from his own theory, that ſuch *ſteam* is no other than the ſame kind of *vapour* thus far mentioned, only rendered capable of a greater *denſity* by more *heat*; he would then have ſeen the importance of that *hygroſcopic law* from its great extent. I have hardly any doubt, that *extreme dryneſs* would reign in a cloſe place ſimilar to PAPIN's *digeſtor*, if there were a ſufficient ſpace above the *water* and a *red heat*; though that ſpace would be filled with *vapour* to the *maximum*. This relates only to the *hygroſcopic law* here in view, on which, in the ſuppoſed caſe, I do not think we could have had any controverſy.

70. Laſtly, let us ſuppoſe, that in order to try the effect of *moiſture* on the ſubſtance he had firſt choſen, or on ſome other of the ſame kind of ſubſtances, but taken *lengthwiſe*, M. DE SAUSSURE had happened to try firſt one of thoſe which, uſed in that manner, have a great *retrogradation*, as *gooſe-quill* and *deal*; and that, after having previouſly obſerved it in the

open air, he had incloſed it in the *moiſt veſſel*, at a time when the place where it ſtood before had the degree of *moiſture* corresponding with the *ſtationary* ſtate of that *hygroſcope*, he would then have obſerved a phænomenon as little expected as the former: for after that inſtrument had been incloſed in the *moiſt veſſel*, it would have moved, by that increaſe of *moiſture*, in the ſame direction as it had done in the open *dry* air when there was on the contrary an increaſe of *dryneſs*. Surpriſed no doubt at that phænomenon, M. DE SAUSSURE would have ſubmitted his new inſtrument to more experiments; he would alſo have tried other *threads*, in which he would have found the ſame ſort of *march*, only at various degrees; and if, in the courſe of thoſe trials, he had ſubmitted the *hair* to the ſame experiments, the ſmallneſs of its motions backwards and forwards, and their irregularities, would not have prevented him from diſcovering in it the ſame ſort of *march* as he had then been uſed to ſee in other *threads*; and thereby, he would have abandoned the whole tribe of *threads* as unfit for the *hygrometer*.

71. The whole of that ſuppoſed courſe of experiments with *ſlips* and *threads*, is that which I have followed from the time I had abandoned the conſtruction of my firſt *hygrometer*; which I did eſpecially with the view of being able to try many ſubſtances. Therefore my theory was formed in conſequence of the two above concluſions, which appear to me immediate, and ſuch as M. DE SAUSSURE could not have drawn differently, if he had followed the ſame neceſſary ſteps: and now I will prove, moreover, that if it had not been for *accidental* circumſtances in his own proceſs, the *hair* alone would have engaged him by degrees to undertake the ſame experiments.

72. M. DE SAUSSURE's firſt *hygrometers*, having their *index* at the top of the frame, could be plunged into *water*; and he tried

tried that method for fixing their point of *extreme moiſture*, as I had done for my firſt *hygrometers*. But in thoſe trials he obſerved, that while his inſtruments ſtood in *water*, their indications remained undetermined within a ſpace of four or five degrees; and attributing that irregularity to a *friction* of the *hair* with *water*, he thought it neceſſary to change my method, to that of placing the *inſtrument* in a *moiſt medium*, which he produced by means of a glaſs jar, *wet* on the inſide, and inverted over *water*. In this method, the ſituation of the *index* became indifferent; and, for ſome particular reaſon, he placed it at the bottom of his new *hygrometers*, which then could not be plunged into *water*. This laſt circumſtance was merely accidental; however, we ſhall ſee how much it has influenced his opinions in reſpect of *extreme moiſture*.

73. In the firſt account of his experiments, M. DE SAUSSURE did not enter into thoſe particulars; but they are in his anſwer to me: and when I there ſaw, that he gave as a reaſon for having *abandoned the immerſion in water*, that a *ſtrong adheſion* of the *hair* to that liquid impeded the freeneſs of its motions, I found it more natural to aſſign to the inſtrument itſelf the unſteadineſs of its *index* which he attributed to that cauſe. Indeed, in thoſe firſt *hygrometers*, one end of the *axis* paſſed through a hole for the purpoſe of carrying the *index* outwards, which was a cauſe of much friction: that *axis* beſides was loaded with the weight of pretty large pincers, holding the *hair*, and that weight was counterpoiſed on the other ſide. Laſtly, the connexion of the *hair* with the *index* was produced by a *ſilver lamina*, which, though very thin, oppoſed ſome reſiſtance in bending round the *axis*. Thoſe are defects that M. DE SAUSSURE corrected afterwards; but they exiſted in the inſtruments which he plunged into *water*, and a weight of

only 3 *grains* was not sufficient to keep their *index* steady, either in *water* or any where; and this he observed himself.

74. Notwithstanding that natural explanation of the unsteadiness of M. DE SAUSSURE's first *hygrometer* when in *water*, I thought it necessary to try, in the same circumstance, some well-constructed *hair hygrometer*; therefore I made two, similar to the last of M. DE SAUSSURE's in every respect, except that of having their *index* at the top; and for the connection of the *hair* with the *axis*, I used a kind of *pincers*, not above half a *grain* in weight, with a hair-like bit of *hemp*, which I know does not alter in any sensible manner the march of the instrument. Now, these *hair* hygrometers, with their *weight* of only 3 *grains*, being put into *water*, follow in it their own laws, arrive and remain *fixed* at their once settled point, as well as any of my other hygrometers. This M. DE SAUSSURE would have also found, if in the improvement of his instrument the *axis* had remained at the top.

75. Before I explain the influence which that *accidental* circumstance of the place of his *index* has had on his opinions with respect to *extreme moisture*, I must mention another of the same kind which has contributed to the same effect. It is evident, both from theory and from M. DE SAUSSURE's own experiments related above (§ 66.), that a sufficient quantity of *water* in any part of a close vessel is the only requisite for producing in it the *maximum* of *evaporation*; but, with a view of accelerating that effect for the common purpose of fixing the point of *extreme moisture* on his *hygrometer*, he prescribes *wetting* the inside of the vessel, besides inverting it over *water*. He did not (and indeed he could not) foresee the consequence of that alteration in his first process; but in fact it was such as to prevent him from discovering, even with time,

time, in those operations, what he would have seen immediately if his last *hygrometers* could have been plunged into *water*.

76. After the improvement of his *hygrometers*, their *index* was no more subject to that unsteadiness observed in the first; consequently they remained *fixed* under the *moist vessel*; but they did not *fix* at the same point every time; and it happened *accidentally*, from causes of irregularity in the *vessel* itself, that in some of the cases, when the *hair* was the *longest*, a *precipitation* of *water* happened (by some partial cooling) on some part of that vessel; and he took that appearance for a sure sign that there was a *superfluous* quantity of *water* in the inclosed air. From that *accidental* connexion of circumstances he concluded, that his *hygrometer* indicated two different states of the *medium* in respect of *extreme moisture*; one, correspondent to about 98 on that instrument, which he considered as real *extreme moisture*, or that state of the *medium* in which no more *vapour* could be introduced into it without a *precipitation*; the other, when such a *precipitation* took place, which he made correspondent to 100, or to the greatest *length* of the *hair*. Prepossessed afterwards with that opinion, when I published my comparative experiments of his *hygrometer* and mine, in which the latter stood sometimes at 80 when the former was at 98, he concluded from that circumstance, that while his *hygrometer* moved only 2 *degrees* by the utmost effect of a *supersaturation* of the *medium*, mine had 20 of those *insignificant degrees*. In this centers the whole of our disagreement, and I am now going to trace its cause in the *accidental* circumstances above described.

77. When I plunge my *hair* hygrometers into *water*, where, as I have said, they come to a *fixed* point, that point does not indicate the *greatest length* of the *hair*; for, on the contrary, that

that *thread* is then *ſhorter* than it is moſt times under the *moiſt veſſel*. This M. DE SAUSSURE would have ſeen, if he had not been prevented, by the ſituation of the *index* in his improved *hygrometer*, from trying again the effect of *water* on the *hair*; and that phænomenon alone would ſurely have given a different courſe to his ideas; eſpecially he would not have ſuppoſed, that the *hair* lengthens 2 *degrees* more, by a *ſuper-ſaturation* of the *medium*, or by the immediate contact of concrete *water*.

78. If, alſo, when he ſettled the manner of determining the point of *extreme moiſture* on his *hygrometers*, M. DE SAUSSURE had retained the ſimplicity of the proceſs he had uſed for his fundamental experiments, in which a piece of *wet* cloth had been ſufficient for producing the *maximum* of *evaporation* in his large veſſel; and, in conſequence, had contented himſelf with inverting his glaſs jar over *water*, without *wetting* it on the inſide, he would have avoided a great cauſe of deception which I am going to explain. In my firſt experiments on the comparative *marches* of our *hygrometers*, in which I followed M. DE SAUSSURE's preſcription for the *moiſt veſſel*, I found ſome *anomalies* which puzzled me. M. DE SAUSSURE himſelf took notice of them in the account I gave of thoſe experiments, and attributed them to my inſtrument. I did not agree with him in that reſpect; but it was long before I could diſcover the real cauſe of thoſe anomalies. The firſt ſtep towards that diſcovery, was the reflecting on the uſeleſſneſs of *wetting* the veſſel on the inſide, for the only purpoſe of producing in it the *maximum* of *evaporation*. That conſideration engaged me to undertake a new courſe of the ſame experiments, with a glaſs jar merely inverted over *water*; and by that means, the greateſt part of the real *anomalies* being removed, I obſerved clearly in the *march* of the *hair*, the combined

combined effects of its own property, and of the *hygroſcopic law* reſpecting *evaporation* which was known to me from other phænomena.

79. As for the cauſe of thoſe *anomalies* which had been removed by the change of the proceſs, the following phænomenon led me to diſcover it. I obſerved frequently, at times when my *hygrometer*, placed under the jar, ſtood at a conſiderable diſtance from its point of *extreme moiſture*, that a very ſmall diminution of heat was ſufficient to cauſe, on the lower part of the veſſel, the formation of a *tarniſhed* rim, extending one or two inches above the ſurface of the *water*, with a thin vaniſhing edge. Having reflected on that phænomenon, from the mechaniſm I aſſign to the operation of *fire* in the very act of *evaporation* I concluded, that in ſuch a *ſtagnant air* every *evaporating* ſurface had an *atmoſphere* of *extreme moiſture*, which extended as indicated by the *tarniſhed rim*; and that it was only beyond that limit that reigned the other *law*, of a decreaſing *moiſture*, correſpondent to the increaſing *maxima* of *evaporation* by an increaſing *heat*. That new *law* of *evaporation* offered evidently an adequate cauſe for explaining the anomalies obſerved in the *wet* veſſel; for *glaſs* retains concrete *water* very imperfectly, and it runs down very ſoon from many of the places which had retained it. Conſequently, under ſuch a partially *wet* veſſel, and differently ſo at different times, the inſtruments muſt be variouſly affected by ſcattered *atmoſpheres* of *extreme moiſture*.

80. However, before I could truſt that explanation, I wanted to ſubmit it to ſome direct experiment; and I ſucceeded by a means which at the ſame time realized what M. DE SAUSSURE thought he had obtained, namely, to produce *extreme moiſture* in a cloſe ſpace, during any common *temperature*,

without any *precipitation* of *water* from the *medium*. This I have produced by means of a wire cage, 4 inches in diameter, covered with cotton cloth, having at the top a *reſervoir*, by which the cloth is kept thoroughly *wet* for a long time; which cage beſides is incloſed in the glaſs jar inverted over water. In that apparatus, though in ſummer-time, every *hygrometer*, either *thread* or *ſlip*, moves and fixes itſelf, not ſo ſpeedily, but elſe exactly as if it was plunged into *water*, without any *ſuperſaturation* of the incloſed *medium*, or *precipitation* of *water* on the *hygroſcopic* ſubſtance.

81. We may ſee now that the idea of two ſorts of *extreme moiſture* is without any foundation. In order to enforce the neceſſity of taking the point of *extreme moiſture* in the *air* brought to that ſtate, and not in *water*, M. DE SAUSSURE ſays, "That the *hygrometer* is not to meaſure the *moiſture* of *water*, "but that of the *air*." This at firſt appears plauſible; however, in reality, *moiſture* is no more to be conſidered in *water* itſelf, than *heat* in the fluid called *fire*. *Water* is the cauſe of *moiſture*, as *fire* is the cauſe of *heat*; but thoſe *effects* are not produced *on* their cauſes; it is *on* other ſubſtances. Therefore, if ſome *hygroſcopic* ſubſtances are placed in a *medium* which has attained *extreme moiſture*, and, in proportion as they take *water* from it, the loſs of that *water* be conſtantly repaired by a new *evaporation*, they will receive by degrees in ſuch a *medium*, without any *precipitation*, as much *water* as if they were plunged in *water* itſelf; for the limit is their *capacity*, which I have explained in § 19. This is the ſame theory that I had expreſſed in my firſt Paper on *Hygrometry*; and it is completely confirmed by the above experiment, with the diſcovery of this new *hygroſcopic law*: "That in a *ſtagnant air*, every "*evaporating* ſurface has an *atmoſphere* of *extreme moiſture*,

"which

"which extends in a ſpace of few inches, diminiſhes rapidly, "and does not interfere beyond that limit with the other *laws* "of *moiſture*."

82. I ſhall now explain, by an example, what is the fundamental *deviation* of an *hair hygrometer*, introducing in it, for a moment, thoſe two ſingular points 98 and 100, which, in M. DE SAUSSURE's experiments, were a very natural cauſe of miſtake. A *hair* hygrometer and mine, being in a cloſe veſſel, at a time when the *temperature*, ſenſibly conſtant, ſhall be but little above 32; if *moiſture* is firſt introduced into that veſſel, ſo as to bring the *hair* hygrometer, by a very ſlow direct motion, to 98, my *hygrometer* will ſtop between 70 and 75; and both inſtruments will be *fixed*, if *moiſture* and *heat* remain the ſame. Let *moiſture* then be made to increaſe very ſlowly, till the *hair* hygrometer has attained its point 100; mine will have arrived at 80; and they again will remain at thoſe points as long as, with the ſame *temperature*, the ſame quantity of *vapour* ſhall remain in the veſſel. Laſtly, let a ſufficient or ſuperfluous quantity of *water* be introduced into the veſſel, the *hair* hygrometer will *retrograde* to 98, and mine proceed to 100, at which points they will ſtop, whatever be the quantity of *water*; and they will remain fixed, as long as the *heat* ſhall not increaſe. This explains the riddle of the ſingular point 98, or of a certain *point*, various in different *hair* hygrometers, various even, at different times, in the ſame individual, at which that inſtrument ſtands with very different degrees of *moiſture*; conſequently, its little motions round that *point* may create great deception, eſpecially from the nature of *organized* ſubſtances, on which now I ſhall add a few words.

83. The above is the fundamental *march* of the *hair* hygrometer, ſuch as it is on the whole, and as it would be conſtantly,

ſtantly, if no other cauſe interfered; but it is ſubject to diſturbing *anomalies*, which become worthy of attention within that ſmall critical ſpace which I have deſcribed. The *texture* of organized *hygroſcopic* ſubſtances occaſions a *friction* between their parts, when, by the changes of *moiſture* and *heat* combined with their *elaſticity*, they undergo changes in their reſpective poſitions; whereby they hardly can completely return to the very ſame arrangement, though with the ſame external circumſtances; even at *extreme moiſture*, when, there being leſs *friction* between their particles, the greateſt part of the diſturbances produced in their former motions are reſtored. To that general cauſe of irregularity is added a particular cauſe, when thoſe ſubſtances are in the ſtate of *hygrometers*; this is the influence of two oppoſite forces acting conſtantly on them; one, the tendency of their component parts to remain united; the other, a *weight* or *ſpring* which tends to ſeparate them. Certain accidental arrangements of their component parts give them more power to reſiſt the action which tends to ſeparate them; and thoſe arrangements are very changeable, by the alternate introduction and expulſion of *moiſture*, by a long ſtay within a ſmall compaſs of variations, and by more or leſs *heat*. This is a large field of facts and ſpeculations, not unintereſting in itſelf, but on which I muſt not dwell: what I have ſaid of thoſe cauſes is ſufficient to account for the *anomalies* to which, more or leſs, every *hygrometer* is ſubject. But whereas in the *ſlips*, thoſe *anomalies* create only ſome irregularities in the obſervations, without any deceiving conſequence in reſpect of the *laws* of *moiſture*, they may deceive when they happen to interfere in the critical part of the march of ſome *threads*; for inſtance, if, by a certain accidental arrangement of the conſtituent parts in a *ſlip* of *whalebone*, there happens to be ſome

tenths of a *degree* difference, from one experiment to another, on its point of *extreme moiſture*, that *anomaly* cannot be of any conſequence on the determination of what muſt be conſidered as that ſtate in the medium; but if it happens to the *hair*, which, in approaching *extreme moiſture*, has but very ſmall motions, it may reverſe thoſe which it had had naturally (as I have obſerved it ſometimes) and become a cauſe of deception.

84. I have now explained, how mere *accidental* circumſtances have been the cauſe of a difference in the ideas that M. DE SAUSSURE and I had formed on what is to be underſtood by *extreme moiſture* in every caſe; and I am going to illuſtrate the whole of that ſubject by a ſingular fact. An *hygrometer* made with a *box thread*, or a thin *faſcicle* of the fibres of that wood, being placed in open air, next to a *hair* hygrometer, or to moſt of the other inſtruments of that ſort, moves in a contrary way from them; but we may lay aſide that circumſtance, by ſuppoſing, that the *numbers*, marked on the *dial* of the firſt, are increaſing in the oppoſite direction from thoſe of the other inſtruments. Let us then ſuppoſe, that ſome experimental philoſopher had choſen the *box thread* for his *hygrometer*; with him I ſhould have fallen into no controverſy on the point of *extreme moiſture*; for, either under the *moiſt veſſel*, or in any other caſe approaching *extreme moiſture*, his *hygrometer* would have moved like mine. But the *box thread*, at approaching *extreme dryneſs*, firſt relents much its pace, then becomes *ſtationary*, and afterwards *retrograde*; by which property, with the concourſe of ſome *accidental* circumſtances as have happened in M. DE SAUSSURE's experiments, the very ſame queſtions that I have examined with ſo much labour in reſpect of *extreme moiſture*, only becauſe of the *hair* hygrometer, would have been tranſported to the point of *extreme*

treme dryneſs; which, however, till now, has created no doubt.

85. That ſingular *thread* illuſtrates alſo another point, cloſely connected with the *retrogradation* of its tribe, but not to be confounded with it; I mean the *recoil*. The common cauſe of both phænomena is, two oppoſite *effects* produced by changes of *moiſture*, on the length of *threads*. The *retrogradation* in the *march*, comparatively to that of *moiſture*, is produced by one of the *effects*, which before was ſurpaſſed by the other, becoming predominant; and the *recoil*, or a returning back a part of a firſt ſtride when *moiſture* changes ſuddenly, is produced, by one of the *effects*, that on the *fibres* themſelves, being performed *ſooner* than that on the ſort of *reticle* formed by the *fibres*. Now, the *box thread* having its *retrogradation* at approaching *extreme dryneſs*, there alſo the *recoil* becomes ſenſible: it appears in the firſt modifications of that *thread* when placed in my *dry veſſel*, by motions backwards and forwards, as it happens to ſome other *threads*, when taken out of *water*, or expoſed to any other ſudden change of *moiſture*. The *box thread* having a ſlow motion, no *recoil* is clearly diſtinguiſhable in its common *march*; whereas the *hair* and the *quill-thread*, which, in appearance, are very quick, have generally a very diſturbed motion when *moiſture* changes ſuddenly. I have ſeen them, when in a free air, and happening to be in their *ſtationary* ſtate, moving quickly one way in a ſpace of 1 or 2 degrees, and then *recoiling* ſlower, ſometimes to the ſame point where they were before, while my *hygrometer* underwent a ſteady change, which was in the firſt direction of theirs. This phænomenon, of a complete *recoil* in the *ſtationary* ſtate of quick *threads*, is ſimilar to the *recoil* of the *index* in thoſe *glaſs frames* that I have deſcribed in § 59. of this Paper, which have a

com-

compenſation for the changes of *heat* by a thin *braſs lamina*: this being ſooner affected than the *glaſs rods*, by ſudden changes of *heat*, the *index* moves firſt one way; then it *recoils* completely, by the change being operated later in the *glaſs rods*.

86. The experiments briefly related in this Paper will, I hope, be ſufficient to anſwer the following queſtion, which has been made to me by ſome obſervers of the two principal *hygrometers* here compared: "Why does the *hair* hygrometer, "when expoſed to the open air in day time, come ſo often "near its point of *extreme moiſture*, while the *whalebone* hygro- "meter almoſt never comes within 30 *degrees* diſtance of that "point in ſummer, and very ſeldom within 20 in winter, even "in *rainy* weather (if preſerved from rain)?" The anſwer, according to the reſults of theſe experiments, is this: "The "general *march* of the *hair* hygrometer is much *decreaſing*, "comparatively to equal increaſes of *moiſture*; that *march* for- "wards, ends in a *ſtationary* ſtate, and is followed by a ſmall "*retrogradation*; while the *whalebone* hygrometer has con- "ſtantly a *march*, if not *proportional*, at leaſt conſtantly *ſimilar* "to that of *moiſture* itſelf."

87. There remains an object of inquiry, which is, a determination of thoſe *ratios* here generally expreſſed. I have explained, in the firſt *part* of this Paper, the difficulties of that object, and what help may be found in comparing the *marches* of *hygrometers* with the *acquiſitions* of *weight* of their ſubſtances, of which proceſs I gave ſome *examples*; and here I ſhall relate ſimilar experiments on *hair*, *whalebone*, *box*, and *aloes-pitta*. But as I have already deſcribed the whole proceſs, and the manner of calculation of that claſs of experiments, I ſhall only give here the reſults of theſe laſt.

88. I

88. I muſt, however, firſt explain another *reduction* that I have added to the former. In the firſt TABLE which I have given of thoſe experiments, I followed the immediate diviſion of my inſtruments, in which 0 correſponds to *extreme dryneſs*, and 100 to *extreme moiſture*. But firſt, under that form, the point called 100 by M. DE SAUSSURE would not appear in its true light, as it means the greateſt *length* of the *hair*; while the point 100 of my ſcale deſigns the ſtate of that *thread* in *water*, where it has a *little retrogradation*. Under that form alſo moſt part of the terms in the obſervation on the *box thread* would be *negative*, ſince it moves very long in a contrary direction to the other hygroſcopes. For theſe reaſons, inſtead of calling 0 the point of *extreme dryneſs*, and that of *extreme moiſture* 100, I have, in the following TABLES, applied the firſt of thoſe denominations to the *ſmalleſt length* of each ſubſtance, and the laſt to their *greateſt length*. That *reduction* produces no difference in the *proportions* between the *terms*, and none in the *terms* themſelves in reſpect to *ſlips*; as in theſe, the *greateſt length* is always obſerved by *extreme moiſture*, and the *ſmalleſt* by *extreme dryneſs*.

II. TABLE

II. TABLE *of comparative changes in the* weight *and in the* length *of the ſame ſubſtances, by the ſame* increaſes of moiſture, *correſpondent to the march of the* ſlip of whalebone *from* 5 *to* 5 *of its degrees.*

	WHALEBONE.		HAIR.		ALOES-PITTA.	
	Increaſes of the weight in ſhavings.	March of the ſlip.	Increaſes of the weight in a maſs of hair.	March of the hair.	Increaſes of the weight in a maſs of pitta.	March of the thread of pitta.
Extr. dryneſs	0.0	0	0.0	0.0	0.0	0.0
	6.0	5	4.8	15.7	6.0	20.6
	11.8	10	8.8	29.0	11.8	35.1
	17.3	15	12.5	40.0	17.3	51.6
	22.2	20	15.9	50.4	22.2	57.6
	26.8	25	19.1	59.7	26.8	75.6
	31.2	30	22.2	67.5	31.2	71.9
	35.2	35	26.6	74.4	35.2	76.3
	39.7	40	29.0	79.3	39.7	83.0
	44.0	45	32.0	83.3	44.0	86.6
	48.1	50	35.0	88.0	48.1	93.6
	52.1	55	38.2	90.0	52.1	96.5
	57.1	60	43.3	92.8	57.1	94.7
	61.7	65	49.8	94.1	61.7	98.2
	66.3	70	55.3	95.4	66.3	100.0
	71.9	75	61.9	97.0	71.9	99.2
	77.6	80	68.7	100.0	77.6	98.2
	* 83.2	85	* 76.0	99.5	* 83.2	96.8
	* 88 8	90	* 84.0	99.2	* 88.8	94.1
	* 94.4	95	* 92.0	98.6	* 94.4	91.9
In *water*	*100.0	100	*100.0	97.7	*100.0	88.3

89. In the above, and in the following experiments, the operation of ſucceſſively introducing *moiſture* into the veſſel was ſtopped, when the *ſlip* of *whalebone* was at 80, as beyond that term the ſmalleſt difference in the *temperature* between the parts of the apparatus creates great anomalies; therefore the fol-

lowing *terms* in the three columns of *weights*, which are marked with an *, have only been added (as I have explained for the former *Table)* with the view of having a common *modulum* between the changes of *weight* and the *marches* of the other inſtruments. But the obſerved *terms* remain in their original *proportions*, and from theſe we may ſee, that the *march* of the *ſlip* of *whalebone* does not differ much from the ſucceſſive increaſe of *weight* in its own ſubſtance; and that when taken out of the apparatus, and immediately plunged into *water*, it proceeds in the ſame direction as before, till it has attained its fixed point, while the *hair* takes, comparatively with the increaſe of *weight* of its own ſubſtance, great ſtrides in the beginning of its *march*, and very ſmall ſteps in the latter part of it, before 100; and then *retrogrades* a little, when taken out of the apparatus, and plunged into *water*. We ſee beſides in that TABLE, that the *thread* of *aloes-pitta*, which at firſt takes ſtill greater ſtrides than the *hair*, has, after a longer *ſtationary* or undetermined ſtate, a determined beginning of *retrogradation* at the ſame time that its own ſubſtance continues to acquire *weight* in the apparatus, and continues that *retrograde* march when, being taken out of the veſſel, it is plunged into *water*.

90. The following TABLE will farther illuſtrate theſe characteriſtic differences of *ſlips* and *threads*, the aſcertaining of which was ſo eſſential to *hygrometry*.

III. TABLE

III. TABLE *of experiments on the comparative changes in the* weight *and the* length *of the ſame ſubſtance by* increaſe *of* moiſture.

BOX.

	Slip of whalebone.	March of the ſlip.	Increaſes of the weight in ſhavings.	March of the thread.
Extr. dryneſs	0	0.0	0.0	72.8
	5	4.5	7.3	87.2
	10	9.5	12.8	93.2
	15	14.5	17.8	97.8
	20	20.0	22.6	100.0
	25	25.7	27.3	95.9
	30	31.5	31.8	92.7
	35	38.0	38.5	88.6
	40	45.5	44.5	79.9
	45	51.5	49.7	70.3
	50	56.5	54.8	63.9
	55	61.2	59.1	57 3
	60	65.7	63.1	51.0
	65	69.7	66.4	47.5
	70	73.7	69.6	40.9
	75	77.7	76.6	31.4
	80	81.5	80.0	21.7
	85	85.9	* 85.0	16.0
	90	90.5	* 90.0	10.4
	95	95.5	* 95.0	5.1
In *Water*	100	100.0	*100.0	0.0

We ſee in this TABLE the *ſlip* of *box* following, in its increaſes of *length*, the increaſe of *weight* in the *ſhavings* of the ſame *wood*, nearly in the ſame manner as the *ſlips* of *whalebone*, *quill*, and *deal*, follow thoſe of their own ſhavings; while the *thread* of *box*, after having gained ſome length by

decreasing steps, begins soon to *shorten*, at the same time that its substance continues to imbibe *water*; being thus the *shortest*, when it cannot receive any more *water* in its pores. That excess of the *hygroscopic* phænomenon of *threads* cannot but throw a full light on the nature of those *hygroscopes*.

91. I am now going to assemble, in two TABLES, the comparative *marches* of all the *threads*, and of all the *slips*, which I have hitherto submitted to that regular course of experiments, laying aside many more of each class, the *marches* of which I only know from common observations. The next TABLE shall contain the experiments on *threads*; in the number of which are two thin natural bodies, which in that respect are similar to the *hair*; one, an animal substance, is a very thin *porcupine quill*; the other, a vegetable, is a thin stem of *gramen*.

TABLE

TABLE *of the correſpondent* marches, *by the ſame* increaſes of moiſture, *of different* THREADS *or vegetable and animal ſubſtances taken* lengthwiſe.

	Porcupine quill.	Whale-bone.	Hair.	Gut.	Aloes-pitte.	Goose-quill.	Deal.	Gramen.	Box.	Slip of wh. bone.
Ext. dryneſs	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	72.8	0
	18.0	12.0	15.6	9.7	20.6	37.0	33.2	26.8	87.4	5
	34.0	29.9	29.4	19.2	35.1	66.6	54.8	48.4	93.2	10
	48.8	39.9	40.9	26.8	51.6	78.7 †	74.9	67.1	97.8	15
	62.3	50.8	50.5	37.0	57.6	88.0	84.6 †	76.6	*100.0	20
	73.3	58.8	59.2	47.1	75.6 †	93.4	89.8	83.9	95.9	25
	81.0	65.3	68.8	57.3	71.9	97.2	93.8	90.5	92.7	30
	86.8	70.8	73.0	67.4	76.3	99.0	96.9	95.1	88.6	35
	90.8	76.1	78.3	75.6	83.0	94.4	94.3	98.6	79.9	40
	93.0	81.4	82.1	82.9 †	86.6	96.2	97.7	*100.0 †	70.3	45
	95.0	85.4	86.1	87.8	93.6	99.0	*100.0	98.8	63.9	50
	94.5	88.4	88.8 †	91.6	96.5	95.3	94.6	98.0	57.3	55
	97.0	90.8	91.6	94.7	94.7	97.2	97.0	97.2	51.0	60
	96.5	92.8	93.8	96.3	98.2	98.2	94.6	96.2	45.7	65
	96.5	95.1	95 6	97.8	*100.0	*100.0	93.0	94.8	40.9	70
	95.0	97.1	97.2	98.7	99.2	99.0	91.4	92.6	31.4	75
	97.0	98.1 †	98.0	*100.0	98.2	98.2	89.0	89.8	21.7	80
	98.0	99.1	100.0	98.7	96.8	97.2	86.9	86.5	16.0	85
	98.6 †	99.6	*100.0	96.8	94.1	95.8	84.6	84.0	10.4	90
	99.1	*100.0	99.3	94.5	91.5	94.4	81.9	80.9	5.1	95
In *water*	100.0	99.5	98.3	91.8	88.3	92.5	79.0	77.0	0.0	100

92. There

92. There the *porcupine quill* ſhews no *retrogradation*; however, conſiſtent with its tribe, it had ſome in other experiments. Its laſt ſteps have the unſteadineſs of the *ſtationary* ſtate, and thereby are ſubject to anomalies. From the ſame cauſe, none of the other *threads* have exactly the ſame ſteps in any two experiments, though on the whole their *march* remains eſſentially the ſame. The *march* here given of the *hair* hygrometer comparatively with mine, is the mean reſult of three experiments, with three different ſets of inſtruments; one of the *hair* hygrometers that I have employed was ſent to me by Mr. PAUL, of Geneva, and its point of *extreme moiſture* was determined in a *fog*. The ſmall and changeable *retrogradation* of the *thread* of *whalebone* and of *hair* might have been overlooked, were it not for other *threads* in which the *retrogradation* begins before that period where the ſtate of *moiſture* is difficult to aſcertain; but from theſe *threads*, that phænomenon is placed in a clear light, which is reflected on the others. I have marked with an * the greateſt *elongation* of each of them, and with a + a point near which their *elongation* begins, and to which they return at laſt. Theſe ſigns will guide the eye in the above TABLE, which ſhews clearly, that no *thread* can be truſted to for the HYGROMETER.

TABLE

TABLE *of the correſpondent* marches *of* SLIPS, *or of fibrous vegetable and animal ſubſtances taken acroſs the fibres, and of ſuch as have no ſenſible fibres.*

	Gooſe quill.	Porcupine quill.	Slip of wh.bone.	Box.	Deal.	Ivory. breadthwiſe.	Ivory lengthwiſe.	Tortoiſe-ſhell.	Horn breadthwiſe.	Horn lengthwiſe.
Ext. dryneſs	0.0	0.0	0	0.0	0,0	0.0	0.0	0.0	0.0	0 0
	4.8	4.8	5	4.5	5.4	6.1	8.3	11.0	9.5	13 8
	9.7	8.8	10	9.5	11.2	12.7	16.6	21.5	18.5	26.8
	14.4	12.0	15	14.5	16.5	18.7	24.6	31.5	27.5	38.8
	19.2	17.0	20	20.0	21.9	24.9	31.5	38.5	37.0	48.8
	23.9	23.4	25	25.7	27.2	30.4	37.6	45.4	46.5	58.0
	28.5	29.4	30	31.5	32.7	35.4	43,6	51.9	54.5	64.6
	33.3	36.0	35	38,0	38.3	41.9	49.7	58.3	62.1	71.0
	38.3	41.4	40	45.5	43.7	47.4	56.3	63.8	68.7	75.5
	42.9	45.4	45	51.4	49.2	53.5	62.4	69.0	72.3	78 5
	47.4	49.8	50	56.5	54.6	58.5	67.4	72.8	77.0	82.2
	52.4	54.8	55	61.2	59.9	63.5	71.6	76.4	79.7	86.2
	56.9	59.7	60	65.7	64.9	68.0	76.1	79.4	84.3	89.6
	61.9	64.4	65	69.7	69.7	72.1	79.1	82.4	86.4	92.4
	67.2	68.5	70	73.7	74.5	76.1	82.9	84.9	88.4	93.4
	72.2	73.5	75	77.7	79.0	80.1	86.7	88.2	90.2	94.4
	77.8	78.9	80	81.5	83.5	84.5	90.4	91.2	92.0	95.4
	82.8	83.9	85	85.9	87.5	87.8	92.4	93.8	94.0	96.6
	88.2	88.9	90	90.5	92.0	92.0	94.5	96.2	96.1	97.8
	94.0	94.4	95	95.3	96.0	96.0	97.5	98.6	98.1	99.0
In *water*	100.0	100.0	100	100.0	100.0	100.0	100.0	100,0	100,0	100.0

93. This

93. This laſt TABLE is the moſt important, as it contains a claſs of *hygroſcopes* which poſſeſs in common the following firſt requiſites for an *hygrometer*; 1ſt, that they may indicate, without any illuſion, both *extreme dryneſs* and *extreme moiſture*; 2dly, that they move conſtantly in the ſame direction as *moiſture* itſelf; 3dly, that they move always when *moiſture* changes. It ſhould ſeem as if the *march* of the *ſlip* of *horn* taken *lengthwiſe*, from its very decreaſing progreſſion, came very near that of the thin *porcupine quill*; but, as I have ſaid, among the ſteps of the latter there are accidental *retrogradations*, and it ſometimes has a final one; and I have never obſerved that diſpoſition in the former, which, in its laſt ſmall ſteps, follows conſtantly the motions of every other *ſlip*.

94. The agreement of all the *ſlips* in this laſt reſpect is a very eſſential circumſtance in hygrometry, as it aſſures us, that we cannot miſtake the caſes when *moiſture* is *extreme* in the *atmoſphere*; a very important point for diſcovering the nature of many *meteorological* phænomena. No *ſlip* will create deception in that reſpect; while, on the contrary, every *thread* may deceive in dubious caſes, and even create great error, if, unknown to the obſerver, it happened to be in the beginning of its *elongation*. There was, however, a queſtion to be decided in that reſpect, namely, whether or not a great *moiſture* in the *medium* was a cauſe of alteration in the *march* of any *hygroſcope*, by producing in its ſubſtance a ſudden irregular *lengthening*. That accidental queſtion is anſwered in the negative by all the *hygroſcopes* of both claſſes: for, in reſpect of the *threads*, inſtead of *lengthening* ſuddenly in that period of *moiſture*, they have then a *retrograde* motion, either continuing or only beginning; and as for the *ſlips*, they, by *lengthening* in the ſame period, only follow their former *laws*: the *ſlips*

which, comparatively to that of *whalebone*, have at firſt ſmall *ſteps*, and which conſequently move in an increaſing progreſſion, continue only to follow that progreſſion; and thoſe which at firſt have greater *ſteps*, and conſequently a decreaſing march, have then ſmall *ſteps* conformable to their individual *law*; therefore, none of thoſe *hygroſcopes* of both claſſes have any *ſudden ſtart*, produced by any degree of *moiſture* in the *medium*, or by the application of concrete *water*; each of them follows, from one end to the other of its ſcale, it own *progreſſion*; and in reſpect of *ſlips*, moiſture is never *extreme* in the ambient *medium*, as long as, in their reſpective *progreſſions*, they have not attained their *greateſt length*.

95. Our common *hygrometer* muſt then be made of one of the *ſlips*; but with that great diſſimilarity obſerved in their *marches*, which of them ſhall we chooſe as indicating the real *march* of *moiſture*? None as yet from that conſideration, which I do not even think a primary one. It is true, that if we truſt to the increaſes of *weight* in thoſe ſubſtances, as being a means of aſcertaining the real progreſs of *moiſture* in the ambient *medium*, the mean rate of ſix experiments of that kind related in this Paper, give the preference to the *ſlip* of *whalebone*; but this I do not yet conſider as deciſive, farther than in what relates to the comparative *marches* of *ſlips* and *threads*; however, as my reaſons of doubt on a more abſolute concluſion cannot be expreſſed in a tranſitory manner, I muſt lay them aſide for the preſent.

96. But, as I have ſaid before, this is not what ought to determine our choice on the *ſubſtance* of a common *hygrometer*, ſince the obſervations themſelves are diſtinct from the conſequences to be drawn from them. Let us ſuppoſe the caſe (which I do not give up) that, with time and reſearches, ſome

proceſs be found by which *known* quantities of *moiſture* may be ſucceſſively produced in the *medium* itſelf. The uſe of that proceſs for hygrometry will be, as M. DE SAUSSURE has begun to do it, to obſerve, on ſome *hygrometer*, the ſucceſſive effects of thoſe known quantities of *moiſture*, from which may be formed a *table* of the correſpondence between the *equal degrees* of the ſcale of the choſen inſtrument, and the real quantities of *moiſture* in the *medium*; and that *table* will ſerve to correct as well *paſt* as *future* obſervations made with that inſtrument. Therefore it matters not what that *hygrometer* ſhall be, provided it is convenient in other reſpects. Let us then examine which of the *ſlips* poſſeſſes the moſt eſſential properties of an *hygrometer*, ſuch as ſhould be in common uſe for comparative obſervations, and to which conſequently future diſcoveries in reſpect of the real proportions between the quantities of *moiſture* itſelf would be applied.

97. *Steadineſs* is ſurely a firſt requiſite for ſuch an inſtrument; and in that reſpect no *ſlip* comes in competition with that of *whalebone*. That property was the firſt motive of my choice; and as an inſtance of it I ſhall only mention, that I have juſt now plunged into *water* an inſtrument of that ſort, of above ten years ſtanding, which is come to its point of *extreme moiſture* as if it had been fixed yeſterday; for, without regard to the diſtance of obſervations, there may be between them a difference of ſome *tenths* of a *degree*. Some other *ſlips* may be brought to a certain degree of *ſteadineſs* by ſtudying what is the degree of *ſtretch* which they may bear; but that attention is not neceſſary for the *ſlip* of *whalebone*: if, for inſtance, when its point of *extreme moiſture* has been fixed while it was *ſtretched* to a certain degree, that *ſtretch* is

5 much

much increaſed, it will acquire ſome abſolute *length*; but it will be *ſteady* again for a new point taken then in *water*.

98. Another property of the *ſlip* of *whalebone*, which at firſt ſhould ſeem contradictory to the former, is its great *expanſibility*, in which alſo it ſurpaſſes all the ſubſtances which I have tried. Such a *ſlip* lengthens above *one-eighth* of itſelf from *extreme dryneſs* to *extreme moiſture*, which produces many advantages in the conſtruction and obſervation of that inſtrument. In reſpect to obſervation, when it is expoſed to the wind, the difference between the chords of the arches of its bends and its real *length* is ſo ſmall, comparatively with its *hygroſcopic* variations, that the indetermination of its *index* will remain confined in a ſpace of one or two *degrees*, when it becomes impoſſible to obſerve *hygrometers* whoſe ſubſtance has but little *expanſion*. Laſtly, of all the ſubſtances which I have reduced to *ſlips*, none is ſo eaſily made thin and narrow as *whalebone*. I have found means for producing eaſily ſuch *ſlips* of it as, with a length of eight *inches*, weigh only about $\frac{1}{10}$th of a *grain*, and are thereby as *quick* as is convenient in other reſpects. All thoſe diſtinctive properties of the *ſlip* of *whalebone* ſeem to point out an *hygroſcopic* ſubſtance fit for our *common hygrometer*.

Deſcription of the whalebone hygrometer.

99. I have now only to deſcribe the conſtruction of that inſtrument as I have fixed it after a long experience. The *fig.* 1. (Tab. IX.) ſhews its form for common uſe. Some of thoſe inſtruments are of the ſame ſize as the *figure*, and they may eaſily be made ſmaller, but commonly they are half as large again in every dimenſion. Their frame will ſufficiently be known from the *figure*,

therefore I ſhall confine myſelf to the deſcription of ſome particulars. The *ſlip* of *whalebone* is repreſented by *a*, *b*; and at its end *a* is ſeen a ſort of *pincers*, made only of a flattened bent wire, tapering in the part that holds the *ſlip*, and preſſed by a ſliding ring. The end *b* is fixed to a moveable bar *c*, which is moved by a ſcrew for adjuſting at firſt the *index*. The end *a* of the *ſlip* is hooked to a thin braſs wire; to the other end of which is alſo hooked a very thin ſilver gilt *lamina*, that has at that end *pincers* ſimilar to thoſe of the *ſlip*, and which is fixed by the other end to the *axis* by a pin in a proper hole. The *ſpring d*, by which the *ſlip* is ſtretched, is made of ſilver gilt wire; it acts on the *ſlip* as a *weight* of about 12 grains, and with this advantage over a *weight* (beſides the avoiding ſome other inconveniencies of this) that, in proportion as the *ſlip* is weakened, in its lengthening by the penetration of moiſture, the *ſpring*, by unbending at the ſame time, loſes a part of its power. The *axis* has very ſmall *pivots*, the *ſhoulders* of which are prevented from coming againſt the frame, by their ends being confined, though freely, between the flat bearing of the heads of two *ſcrews*, the front one of which is ſeen near *f*. The ſection of that *axis*, of the ſize that belongs to a *ſlip* of about 8 inches, is repreſented in *fig.* 2.; the *ſlip* acts on the diameter *a*, *a*, and the *ſpring* on the ſmaller diameter *b*, *b*.

100. Another conſtruction of the ſame inſtrument is repreſented, of half its dimenſions, by *fig.* 3. (Tab. X.) The eſſential parts of that *hygrometer* are the ſame as in the other; therefore I ſhall only mention the differences adapted to the uſe of fixing it out of a window. The *figure* repreſents the manner in which it is fixed, with its *dial* turned half-way towards the obſerver, by a proper head of the hooks which hold it at the top and the bottom.

bottom. The ſteady part of the inſtrument is repreſented by *fig.* 4.; it is partly an half tube, cut longitudinally through the axis, and having ſidewiſe two vertical rows of large holes. A whole tube goes over that half one; for which purpoſe the piece *a*, *a*, is taken off by unſcrewing the female ſcrew *b* which belongs to the part *d*; this laſt is a ſmall open tube, cut outſide in ſcrew. The large external tube has alſo two vertical rows of holes, at ſuch a diſtance from one another, that, when one of them correſponds to one of the rows of the half tube, the other is in front. The two oppoſite poſitions that tube can be brought to are in order that the inſtrument may be placed either ſide of a window, and each of thoſe poſitions is determined by one end of a cut at the bottom of the tube in *a*, *fig.* 3. which then holds againſt the ſteady pin *c*, *fig.* 4. The rows of holes of the tube are to be turned towards the room, to prevent the rain from falling on the *ſlip*; and the *dial* being incloſed in a box with a glaſs in front, no rain can get into the inſtrument. It muſt alſo be fixed in a place not much expoſed to the ſun, or be ſcreened from it without preventing the circulation of the air. The communication of the *ſlip* with the external air through the rows of holes and the open bottom *d*, *fig.* 4. is ſufficient for that claſs of obſervations. By the manner of its being hooked, it may be eaſily taken off for carrying elſewhere; and, if a quick obſervation is wanted, the tube alſo may eaſily be taken off.

I have the honour of preſenting one of thoſe inſtruments to the Royal Society; and, as it is very deſirable that ſome hygrometer be added to the other meteorological inſtruments uſually obſerved, I wiſh this may deſerve a place in their Obſervatory for that purpoſe.

Fig. 1.



Fig. 2.



J. B. Hans del.

Basire Sc.

Fig. 3.



Fig. 4.

J. B. Hans del.

Basire sc.

PRESENTS

MADE TO THE

ROYAL SOCIETY

From November 1790 to June 1791;

WITH

The NAMES of the DONORS.

1790	Presents.	Donors.
Nov. 4.	The Nautical Almanacs for the Years 1793, 1794, 1795, and 1796. 4 Vols. London, 1790. 8°	The Commissioners of Longitude.
	A Collection of Ordinances and Regulations for the Government of the Royal Household. London, 1790. 4°	The Society of Antiquaries.
	Transactions of the Society for the Encouragement of Arts, Manufactures, and Commerce. Vol. VIII. London, 1790. 8°	The Society for the Encouragement of Arts, Manufactures, and Commerce.
	Report of the Committee of the Highland Society of Scotland, on the Subject of Shetland Wool. Edinburgh, 1790. 4°	The Highland Society of Scotland.
	Traité Analytique des Mouvemens apparens des Corps céleſtes, par M. Dionis du Séjour. Tome II. Paris, 1789. 4°	M. Dionis du Sejour, F. R. S.
	N. J. Jacquin Collectanea ad Botanicam, Chemiam et Hiſtoriam Naturalem. Vol. III. Vindobonæ, 1789. 4°	Profeſſor de Jacquin, F. R. S.
	Cautions to the Heads of Families, by A. Fothergill. Bath, 1790. 8°	Anthony Fothergill, M. D. F. R. S.
	Machine hydraulique d'une nouvelle invention, par T. Wex. Nantes, 1787. 8°	M. Wex.

1790	Presents.	Donors.
Nov. 11.	Memoirs of the Literary and Philosophical Society of Manchester. Vol. III. Warrington, 1790. 8°	The Literary and Philosophical Society of Manchester.
	Medical Communications. Vol. II. London, 1790. 8°	The Society for promoting Medical Knowledge.
	A Treatise on the Digestion of Food, by G. Fordyce. London, 1790. 4°	George Fordyce, M. D. F. R. S.
	Linnæan Zoology, by G. Shaw; the Figures by J. Sowerby. N° I. London, 1790. 4°	George Shaw, M D.F.R.S. and Mr. James Sowerby.
	The Naturalist's Miscellany, by G. Shaw; the Figures by F. P. Nodder. Vol. I. London, 1790. 8°	George Shaw, M.D.F.R.S. and Mr. F. P. Nodder.
	Principes sur les Mésures en Longueur et en Capacité, sur les Poids et les Monnoies, par M. Bonne. Paris, 1790. 8°	M. Bonne.
	Prælectio Academica simpliciores et salubriores comprehendens de Febribus notiones, a Jos. Pinilla Vizcayno. Compluti, 1790. 4°	Professor Pinilla Vizcayno, of Alcala.
18.	Nova Acta Academiæ Scientiarum Imperialis Petropolitanæ. Tomus V. Petropoli, 1789. 4°	The Imperial Academy of Sciences of Petersburg.
	Zwo abhandlungen über die Nutritions kraft, welche von der Kayserl. Academie der Wissenschaften in St. Petersburg den Preis getheilt erhalten haben. St. Petersburg, 1789. 4°	
	Commentarii de Rebus in Scientia Naturali et Medicina gestis. Volumen XXXI. Lipsiæ, 1789. 8°	Mr. Hurlock, F. R. S.
	Observations on Poisons, by T. Houlston. Edinburgh, 1787. 8°	Mr. Houlston.
25.	Kongl. Vetenskaps Academiens Nya Handlingar, Tom. X. for 1789, 3d and 4th Quarter; and Tom. XI. for 1790, 1st and 2d Quarter. Stockholm. 8°	The Royal Academy of Sciences of Stockholm.
	Maniera di migliorare e dirigere i Palloni aerei, inventata e dimostrata dai Fratelli Gerli. Roma, 1790. 8°	Sign. Agostino Giosseppe e Carlo Giosseppe Gerli.
Dec. 9.	A Portrait of Dr. Bradley, late Astronomer Royal and F. R. S.	The Rev. Mr. Peach.
	A Portrait of the late Benjamin Franklin, LL.D. F. R. S.	Caleb Whitefoord, Esq. F. R. S.
	Meteorological Journal kept at Bengal, by the late Colonel T. D. Pearse, from June 28, 1787, to April 9, 1789. MS. fol.	Sir Joseph Banks, Bart. P. R. S.

1790	Presents.	Donors.
Dec. 9.	Oratio Anniversaria in Theatro Coll. Reg. Med. Lond. ex Harveii instituto habita a J. Ash. Londini, 1790. 4°	John Ash, M. D. F. R. S.
	Index Ornithologicus, studio Jo. Latham. Londini, 1790. 2 Vols. 4°	Mr. Latham, F. R. S.
	Extrait des Observations Astronomiques et Physiques faites à l'Observatoire Royal en l'Année 1789. 4°	M. de Cassini, F. R. S.
23.	Géographie des Grecs analysée, par M. Gosselin. Paris, 1790. 4°	M. Gosselin.
1791		
Jan. 13.	The London Medical Journal. Vol. XI. London, 1790. 8°	Samuel Foart Simmons, M. D. F. R. S.
20.	A Narrative of the Building, and a Description of the Construction, of the Edystone Lighthouse with Stone, by J. Smeaton. London, 1791. fol.	John Smeaton, Esq. F. R. S.
	Journal of a Voyage to New South Wales, by John White, Esq. London, 1790. 4°	Thomas Wilson Esq.
27.	Magnetic Atlas, or Variation Chart, by J. Churchman.	Mr. John Churchman.
	An Explanation of the Magnetic Atlas, or Variation Chart, by J. Churchman. Philadelphia, 1790. 8°	———
Feb. 3.	Observations générales sur la Réspiration et sur la chaleur animale, par M. Séguin. 4°	M. Seguin.
	Relacion de las Epidemias, que han afligido á la Ciudad de Cartagena, por D. Mart. Rodon y Bell. Cartagena, 1787. 4°	Don Martin Rodon y Bell.
	Discurso que para dar Principio a los primeros Exercicios Publicos de la Botanica dixo el D. D. Mart. Rodon. Cartagena, 1785. 4°	———
	The Antiquaries Museum, by J. Schnebbelie. N° I. 4°	Mr. Schnebbelie.
17.	Libellus de Natura, Causa, Curationeque Scorbuti. Auctore N. Hulme. Londini, 1768. 8°	Nathaniel Hulme, M. D.
	A Treatise on the Puerperal Fever, by N. Hulme. London, 1772. 8°	———
	Oratio de re Medica cognoscenda et promovenda, Auctore N. Hulme. Londini, 1777. 4°	———

1791	Presents.	Donors.
Feb. 17.	A safe and easy Remedy, proposed for the Relief of Stone and Gravel, &c. by N. Hulme. London, 1788. 4°	Nathaniel Hulme, M. D.
Mar. 10.	Letters on the Elements of Botany, by J. J. Rousseau, translated into English, with 24 additional Letters, by T. Martyn. London, 1791. 8°	The Rev. Thomas Martyn, B. D. F. R. S.
	Thirty-eight Plates, with Explanations, adapted to the Letters on the Elements of Botany, by T. Martyn. London, 1788. 8°	
	17. A Treatise of the Plague, by P. Russell. London, 1791. 4°	Patrick Russell, M. D. F. R. S.
	24. Observationes Siderum habitæ Pisis ab Anno 78 ad Annum 82 vertentis Sæculi, a Jos. Slop de Cadenberg. Pisis, 1789. fol.	Professor Slop, of Pisa.
	31. A Treatise on the Angina Pectoris, by W. Butter. London, 1791. 8°	William Butter, M. D.
April	4. Icones selectæ Plantarum, quas in Japonia collegit et delineavit Eng. Kæmpfer. Londini, 1791. fol.	Sir Joseph Banks, Bart. P. R. S.
	An historical Developement of the present political Constitution of the Germanic Empire, by J. S. Pütter. Translated from the German, with Notes, by J. Dornford. London, 1790. 3 Vols. 8°	Josiah Dornford, LL.D.
	A Discourse on the Principle of Vitality, by B. Waterhouse. Boston, 1790. 4°	Prof. Waterhouse, of Cambridge in America.
	14. Recherches sur cette Question: la Chaleur naturelle de l'Homme peut-elle être considérée comme un Terme fixe? par M. Gauffen. Montpellier, 1787. 8°	M. Gauffen, of Montpellier.
	Dissertation sur le Thermomètre de Reaumur, par M. Gauffen. Beziers, 1789. 8°	
May	5. Pharmacopœia Collegii Regalis Medicorum Londinensis. Londini. 8°	The College of Physicians.
	An Account of the Life and Writings of Thomas Day, Esq. London, 1791. 8°	James Keir, Esq. F. R. S.
	Eléments de l'Art de la Teinture, par M. Berthollet. Tomes II. Paris, 1791. 8°	M. Berthollet, F. R. S.
	A short Account of a Passage from China late in the Season. London, 1788. 4°	Mr. George Robertson.
	Charts of the China Navigation, by G. Robertson. London, 1788. fol.	

1791	Presents.	Donors.
May 5.	Chart of the China Sea, including the Philippina, Mollucca, and Banda Islands, by G. Robertson, 1791.	Mr. George Robertson
	Memoir of a Chart of the China Sea, by G. Robertson, 1791. 4°	
12.	Transactions of the Royal Irish Academy, 1789. Dublin. 4°	The Royal Irish Academy.
	Astronomisches Jahrbuch für das Jahr 1793, von J. E. Bode. Berlin, 1790. 8°	Mr. J. E. Bode, F. R. S.
26.	Illustrations of British History, Biography, and Manners, exhibited in a Series of original Papers, by E. Lodge. 3 Vols. London, 1791. 4°	Edmund Lodge, Esq.
	Lettre de M. Van Marum, contenant la Déscription d'une Machine électrique, construite d'une manière nouvelle. 4°	Martinus Van Marum, M. D. of Harlem.
	Observations générales sur les Sensations, par M. Séguin. 8°	M. Seguin.
June 2.	Saggio di Litologia Vesuviana, dal Cav. G. Gioeni. Napoli, 1790. 8°	Cavaliere Giuseppe Gioeni.
	Bestimmung der Höhen der bekanntern Berge des Canton Bern, von J. G. Tralles. Bern, 1790. 8°	Professor Tralles, of Bern.
9.	A Print of John Smeaton, Esq. F. R. S. engraved by W. Bromley.	Alexander Aubert, Esq. F. R. S.
	Observations on the Diseases, Defects, and Injuries in all Kinds of Fruit and Forest Trees; with an Account of a Cure invented by W. Forsyth. London, 1791. 8°	Mr. William Forsyth.
	Essai d'une Classification des Couleurs. MS. 4°	M. T. F. Opitz, of Tschaslau in Bohemia.
23.	The Philosophy of Ancient Greece investigated in its Origin and Progress, by W. Anderson. Edinburgh, 1791. 4°	The Rev. Walter Anderson, D. D.
July 7.	Icones stirpium rariorum Descriptionibus illustratæ, Auctore R. A. Salisbury. Londini, 1791. fol.	Richard Anthony Salisbury, Esq. F. R. S.
14.	Oriental Repertory, published by A. Dalrymple. N° I. London, 1791. 4°	Alexander Dalrymple, Esq. F. R. S.
	Medical Facts and Observations. Vol. I. London, 1791. 8°	Samuel Foart Simmons, M. D. F. R. S.
	A Whalebone Hygrometer.	J. A. De Luc, Esq. F. R. S.

AN INDEX TO THE EIGHTY-FIRST VOLUME OF THE PHILOSOPHICAL TRANSACTIONS.

A.

ABSTRACT of a regiſter of the barometer, thermometer, and rain, at Lyndon in Rutlandſhire, for the year 1789, p. 89.

Account of ſome appearances attending the converſion of caſt into malleable iron, p. 173.

Acid, nitrous, though conſiſting of the ſame elements with pure water, contains a greater proportion of dephlogiſticated air, p. 219. Phoſphoric acid, an ingredient in James's powder, p. 345.

Air, extreme moiſture exiſts in, p. 19. The maximum of evaporation in a maſs of incloſed air, far from being identical with the maximum of moiſture, p. 20. Obſervation on the production of heavy inflammable air, p. 176. Fixed air contains a much larger portion of oxygène than of charcoal, p. 177. Obſervations on the decompoſition of fixed air, p. 182. Fixed air produced by charcoal, ibid. Vital air more ſtrongly attracted by charcoal than by phoſphorus, p. 183. Experiments relating to the decompoſition of dephlogiſticated and inflammable airs, p. 213. Inflammable air procured from iron by means of ſteam, p. 216. Nitrous acid, though conſiſting of the ſame elements with pure water, contains a greater proportion of dephlogiſticated air, p. 219. In a ſtagnant air every evaporating ſubſtance has an atmoſphere of extreme moiſture, p. 403.

Alcohol,

Alcohol, ſugar has no ſenſible affinity with, p. 11. Alcohol, having entered ſugar, by means of its capillary pores, evaporates, and leaves the ſugar ſenſibly as it was before, ibid.

Aleppo, diſtance between it and Buſſorah, p. 137.

Ambergris, on the production of, p. 43. Ambergris found in a ſpermaceti whale, ibid. Moſt likely to be found in ſickly whales, p. 47.

Angles, deſcription of a ſimple micrometer for meaſuring ſmall ones with the teleſcope, p. 283.

Antimonialis, pulvis, experiments on, p. 349.

Antimony and antimonial calces, various experiments on, p. 351. Antimonial calx, an ingredient in James's powder, p. 345. Sulphur of antimony no way neceſſary to the formation of James's powder, p. 360.

Arch. Conſiderations on the convenience of meaſuring an arch of the meridian, and of the parallel of longitude, having the Obſervatory at Geneva for their common interſection, p. 106.

Atmoſpheric electricity, meteorological journal principally relating to, p. 185.

B.

Barker, Thomas, abſtract of a regiſter, for the year 1789, of the barometer, thermometer, and rain, at Lyndon in Rutlandſhire, p. 89.

Barometer, abſtract of a regiſter of one, kept at Lyndon in Rutlandſhire, for the year 1789, p. 89.

Baſaltes and granite, obſervations on the affinity between, p. 48. Baſaltes and granite gradually approach and change into one another, p. 50. Feldſpath, ſhoerl, and mica, ſubſtances common both to baſaltes and granite, and tend to eſtabliſh the near relation between theſe two kinds of rock, p. 54. Baſaltes and granite ſo involved in one another, that they are ſuppoſed to have undergone the ſame operations of nature at the ſame time, p. 58.

Beddoes, Dr. on the affinity between baſaltes and granite, p. 48. On ſome appearances attending the converſion of caſt into malleable iron, p. 173.

Buſſorah, its diſtance from Aleppo, p. 137.

C.

Cacholong, Hydrabad Tabaſheer reſembles, p. 369.

Calces, antimonial, experiments on, p. 351.

Calculi, human, experiments on, p. 223.

Camels, on the rate of travelling, as performed by; and its application, as a ſcale, to the purpoſes of geography, p. 129. A camel's load 500 or 600 pounds, p. 136.

Capella, a ſtar of the firſt magnitude, culminates between the zenith of Geneva and St. Jean Maurienne, p. 115.

Capillary pores, water enters ſugar by means of, p. 11. Alcohol enters ſugar by the ſame means, ibid. The penetration of animal ſubſtances produced by the faculty of capillary pores, p. 12.

Caſes in which horny excreſcences have appeared on the human body, p. 96.

Cavallo, Mr. Deſcription of a ſimple micrometer for meaſuring ſmall angles with the teleſcope, p. 283.

Cauſeway, Giant's, many of its pillars conſiſt of fine-grained, dark-coloured whinſtone, p. 51.

Chalk, a great deal of chalky ground in the ſouthern parts of England, p. 281. Vein of chalk runs from the ſea in Devonſhire, along the ſouthern counties of England to Dover, ibid. Another vein parts off from that about Reading in Berkſhire, and continues its courſe by Dunſtable, Baldock, and Gogmagog Hills, to the ſea in Norfolk, ibid. Conſiderable quantities of chalk in the wolds of Yorkſhire, ibid. Bed of chalk diſcovered near Ridlington in Rutland, ibid. Chalk about Ridlington harder than that of the ſouth of England, ibid.

Chermes Lacca, account of, p. 228. Frequently adheres to the branches of the Mimoſa cinerea, ibid. Vaſt quantities of exceedingly ſmall animals obſerved upon the lac and branches, ibid. Deſcription of the male lac inſect in its perfect ſtate, p. 231. Deſcription of the female lac inſect in its perfect ſtate, ibid. Mr. Hellot's proceſs for extracting the colouring matter from dry lac, p. 233.

Conſiderations on the convenience of meaſuring an arch of the meridian, and of the parallel of longitude, having the Obſervatory of Geneva for their common interſection, p. 106.

D.

Dalby, Iſaac, on the longitudes of Dunkirk and Paris from Greenwich, deduced from triangular meaſurement, ſuppoſing the earth to be an ellipſoid, p. 236.

Dephlogiſticated air. See *Air*.

Deſcription of a ſimple micrometer for meaſuring ſmall angles with the teleſcope, p. 283. Deſcription of the whalebone hygrometer, p. 419.

Devonſhire, vein of chalk runs from the ſea coaſt of, along the ſouthern counties of England to Dover, p. 281. A vein of chalk runs from the ſea coaſt of Devonſhire to the ſea coaſt of Norfolk, ibid.

Diſtance between Aleppo and Buſſorah, p. 137.

Dover, vein of chalk, which runs from Devonſhire, terminates at, p. 281.

Dryneſs, abſolute, can only be produced by fire, p. 1. Obſervations on abſolute dryneſs, p. 2. Dryneſs produced by pot-aſh, p. 4. Point of dryneſs capable of fixation by lime, p. 5.

Dunkirk, its longitude from Greenwich deduced from triangular meaſurement, ſuppoſing the earth to be an ellipſoid, p. 236.

E.

Electricity, meteorological journal, principally relating to atmoſpheric, kept at Knightsbridge, p. 185. Deſcription of an inſtrument for collecting atmoſpheric electricity, ibid. Remarks on ſome phænomena exhibited by a rod on Aug. 31, 1789, p. 211.

Evaporation, the cauſe of hygroſcopic bodies loſing part of their moiſture, p. 2. Experiments on the maximum of evaporation, and its correſpondence with the maximum of moiſture in a medium, p. 17. The maximum of evaporation in a maſs of incloſed air, far from being identical with the maximum of moiſture, p. 20. In a ſtagnant air every evaporating ſubſtance has an atmoſphere of extreme moiſture, p. 403.

Excreſcences, obſervations on certain ones of the human body, p. 95.

Experiments on hygrometry, p. 2—42. Experiments on abſolute dryneſs, p. 2. On extreme moiſture, p. 10. On the maximum of evaporation, and its correſpondence with the maximum of moiſture in a medium, p. 17. On two diſtinct claſſes of hygroſcopes, p. 20. On the ſcale of the hygrometer between the two fixed points, p. 24. On the comparative changes of weight and dimenſions of ſome hygroſcopic ſubſtances, p. 27. On the recoil of hygroſcopic threads, p. 39. On the converſion of caſt into malleable iron, p. 173. Concluſions drawn from experiments on iron, p. 175. Experiments relating to the decompoſition of dephlogiſticated and inflammable airs, p. 213. Inflammable air procured by means of ſteam, p. 216. Experiments on human calculi, p. 223. Experiments and obſervations to inveſtigate the compoſition of James's powder, p. 317. Experiments on the ſenſible properties of James's powder, ibid. On its ſpecific gravity, p. 318. Experiments on the effects of fire on this powder, ibid. Experiment ſhewing that antimony, mixed with earthy matter, is uſed in the compoſition of James's powder, p. 320. Experiments with different menſtrua applied to James's powder, ibid. Experiment ſhewing the proportion in which James's powder diſſolves in water, p. 324. Experiment with marine acid applied to James's powder, which had not been expoſed to the action of nitrous acid, or any other menſtruum, p. 344. Experiments on James's powder with fixed alkalies, p. 345. Synthetic experiments on James's powder, ibid. Experiments on the Pulvis antimonialis, p. 349. Various experiments on antimonial calces, p. 351. Chemical experiments on Tabaſheer, p. 368. Experiments on Tabaſheer, treated with water, p. 370. With vegetable colours, p. 371. At the fire, ibid. With acids, p. 373. With liquid alkalies, p. 375. With dry alkalies, p. 379. With other fluxes, p. 380. Table of experiments on comparative changes in the weight and the length of the ſame ſubſtance by increaſing moiſture, p. 411.

F.

Feldſpath, a ſubſtance common both to baſaltes and granite, p. 54.

Fever powders, receipts for making, p. 346, 347.

Fire, considered as the cause of heat, the only agent by which absolute dryness can be immediately produced, p. 1. Fire capable of producing any required degree of compactness, as lavas abundantly shew, p. 55. The effects of fire on James's powder, p. 318. Fire powerful in the change of the colour of bodies, p. 362. Fire the cause of heat, p. 402.

Fixed air. See *Air*.

Flints, rows of, in a chalk-pit near Ridlington in Rutland, p. 281. Flints about Ridlington not so black as those of the southern counties, but veined, ibid.

Fyfield, in Hampshire, abstract of a register of rain kept at, for the year 1789, p. 89.

G.

Geneva. Considerations on the convenience of measuring an arch of the meridian, and of the parallel of longitude, having the Observatory of Geneva for their common intersection, p. 106. Capella, a star of the first magnitude, culminates between the zeniths of Geneva and St. Jean Maurienne, p. 115.

Geography. On the rate of travelling, as performed by camels; and its application as a scale, to the purposes of geography, p. 129.

Giant's Causeway, many of its pillars consist of fine-grained, dark-coloured whinstone, p. 51.

Granite and basaltes, observations on the affinity between, p. 48. Almost all granites melt into a black glass, p. 60. Granitic lavas are granite rocks fused, p. 63. Malvern Hills, composed of granite, p. 65.

Greenwich, the longitude of Dunkirk from, deduced from triangular measurement, supposing the earth to be an ellipsoid, p. 236. Longitude of Paris from Greenwich, ibid.

H.

Heat, at its maximum in a body, when it is incandescent, p. 3. Heat caused by fire, p. 402.

Herschel, Dr. on nebulous stars, p. 71.

Home, Everard, on cerain horny excrescences of the human body, p. 95.

Human body, observations on certain horny excrescences of, p. 95. The case of Mrs. Lonsdale, p. 96. The case of Mrs. Allen, p. 98.

Hydrabad Tabasheer, experiments on, p. 369. Resembles Cacholong, ibid.

Hygrometer, fundamental propositions for the construction of an, p. 1. Absolute dryness can only be produced by fire, ibid. Extreme moisture can only be produced by means of water, ibid. 16. Observations on absolute dryness, p. 2. An hygroscopic body cannot lose any part of its moisture, but by evaporation, ibid. Dryness capable of fixation by lime, p. 5. An hygroscopic body, which is brought to incandescence, cannot contain any evaporable water, p. 9. The point of extreme moisture of an hygrometer known by the excess of water, p. 17. Observations on two distinct classes

classes of hygroscopes, p. 20. Observations on the scale of the hygrometer between the two fixed points, p. 24. Experiments on the comparative changes of weight and dimensions of some hygroscopic substances, p. 27. Experiments on the recoil of hygroscopic threads, p. 39. Hygrometers variously affected by scattered atmospheres of extreme moisture, p. 401. Steadiness the first requisite for an hygrometer, p. 418. Description of the whalebone hygrometer, p. 419.

Hygrometry, observations on, p. 1. 389.

I.

James's powders, experiments and observations to investigate the composition of, p. 317. Its nature hitherto unknown to the public, ibid. Sensible properties of the powder, ibid. Smooth to the feel, with some rather rough particles, ibid. Gritty in the mouth, p. 318. Perfectly inodorous, ibid. On the specific gravity of James's powder, ibid. Much heavier than any of the common earths and stones in a pulverized state, ibid. The effects of fire on this powder, ibid. Antimony mixed with earthy matter, found by experiment in James's powder, p. 320. Experiments with different menstrua applied to James's powder, ibid. The proportion in which James's powder dissolves in water shewn, p. 324. Conclusions drawn from experiments on James's powder, p. 325. Experiment with marine acid applied to James's powder, which had not been exposed to the action of nitrous acid, or any other menstruum, p. 344. Phosporic acid, lime, and antimonial calx, ingredients in James's powder, p. 345. Experiments on James's powder with fixed alkalies, ibid. Sulphur of antimony no ways necessary to the formation of James's powder, p. 360. The ingredients made use of in the formation of James's powder enumerated, p. 367.

Infinite series, observations on, p. 146. New method of investigating the sums of infinite series, p. 295.

Inflammable air, observation on the production of heavy, p. 176. Experiments relating to the decomposition of inflammable air, p. 213. Inflammable air procured from iron by means of steam, p. 216.

Insect. Description of the male lac insect in its perfect state, p. 231. Description of the female lac insect in its perfect state, ibid.

Journal, meteorological. See *Meteorological Journal*.

Iron, Account of some appearances attending the conversion of cast into malleable, p. 173. Conclusions drawn from experiments on iron, p. 175. Grey iron contains a large portion of plumbago, p. 177.

Ivory, the temperature of water has a sensible effect on the expansion of, p. 15.

L.

Lac. See *Chermes Laccæ*.

Lambeth, South, abstract of a register of rain kept at, for the year 1789, p. 89.

Lane, Timothy, on human calculi, p. 223.

Lavas, abundantly ſhew, that fire is capable of producing any required degree of compactneſs, p. 55. Granitic lavas are granite rocks fuſed, p. 63.

Life. On the method of determining, from the real probabilities of life, the values of contingent reverſions in which three lives are involved in the ſurvivorſhip, p. 246.

Lime, an ingredient in James's powder, p. 345.

Liſt of Preſents, p. 423.

Longitude, conſiderations on the convenience of meaſuring an arch of the parallel of, p. 106.

Longitudes of Dunkirk and Paris from Greenwich, deduced from triangular meaſurement, ſuppoſing the earth to be an ellipſoid, p. 236.

Luc, J. A. de, on hygrometry, p. 1. 389.

M.

Macie, Mr. on Tabaſheer, p. 368.

Malvern Hills, compoſed of granite, p. 65.

Meteorological Journal, relating to atmoſpheric electricity, p. 185. See *Electricity*.

——— kept at the Apartments of the Royal Society, for January 1790, App. p. 2. February, p. 4. March, p. 6. April, p. 8. May, p. 10. June, p. 12. July, p. 14. Auguſt, p. 16. September, p. 18. October, p. 20. November, p. 22. December, p. 24.

Mica, a ſubſtance common both to baſaltes and granites, p. 54.

Micrometer, deſcription of a ſimple one for meaſuring ſmall angles with the teleſcope, p. 283. Teleſcopical micrometers, divided into two claſſes, ibid.

Milky way, conſiſts entirely of ſtars, p. 71.

Moiſture, in hygroſcopic bodies, can only be produced by means of water, p. 1. Hygroſcopic bodies cannot loſe part of their moiſture, but by evaporation, p. 2. Quick-lime poſſeſſes a great capacity for moiſture, p. 4. Obſervations on extreme moiſture, p. 10. Moiſture, a quantity of inviſible water, either evaporable, or evaporated, p. 14. The maximum of moiſture known by the exceſs of water, p. 17. Extreme moiſture exiſts in the air, p. 19. The idea of two ſorts of extreme moiſture without foundation, p. 402. In a ſtagnant air every evaporating ſubſtance has an atmoſphere of extreme moiſture, p. 403.

Morgan, Mr. on the method of determining, from the real probabilities of life, the values of contingent reverſions in which three lives are involved in the ſurvivorſhip, p. 246.

N.

Nebulous ſtars, obſervations on, p. 71. Diffuſed nebuloſities obſerved about the conſtellation of Orion, p. 85.

P.

P.

Paris, its longitude from Greenwich deduced from triangular meaſurement, ſuppoſing the earth to be an ellipſoid, p. 236.

Pearſon, Dr. on the compoſition of James's powder, p. 317.

Phoſphorus, not obtainable by paſſing marine acid through a compound of bones and charcoal, when red-hot, p. 184. The attraction of phoſphorus and lime for vital air exceeds the attraction of charcoal by a greater force than that ariſing from the attraction of marine acid for lime, ibid.

Pictet, Mark Auguſtus, on the convenience of meaſuring an arch of the meridian, and of the parallel of longitude, having the Obſervatory of Geneva for their common interſection, p. 106.

Pores, capillary, water enters ſugar by means of, p. 11. Alcohol enters ſugar by the ſame means, ibid. The penetration of animal ſubſtances produced by the faculty of capillary pores, p. 12.

Pot-aſh, with ſome other alkaline ſubſtances produce dryneſs, p. 4.

Powder. See *James's Powder*.

Preſents, liſt of, p. 423.

Prieſtley, Dr. on the decompoſition of dephlogiſticated and inflammable airs, p. 213.

Problems reſpecting ſurvivorſhips, p. 248, 253, 256, 258, 261, 263, 265, 267, 269, 272. Problems relating to angles, p. 290, 291.

Propoſitions, fundamental, for the conſtruction of an hygrometer, p. 1.

——— relating to infinite ſeries, p. 299, 300, 302, 303, 308, 309, 310, 311, 312, 314.

Pulvis antimonialis, experiments on, p. 349.

Q.

Quicklime, poſſeſſes a great capacity for moiſture, p. 4.

R.

Rain, abſtract of a regiſter of, kept at Lyndon in Rutlandſhire, at South Lambeth in Surrey, and at Selbourn and Fyfield in Hampſhire, for the year 1789, p. 89. A leſs quantity of rain fell in the year 1788, than in any year ſince 1750, p. 90.

Read, Mr. on atmoſpheric electricity, p. 185.

Receipts for making fever powders, p. 346, 347.

Rennell, Mr. on the rate of travelling, as performed by camels; and its application, as a ſcale, to the purpoſes of geography, p. 129.

Reſults from a ſeries of experiments on hygrometry, p. 41.

1. Fire, as cauſe of heat, the only means of obtaining extreme dryneſs, ibid.
2. Water, in its liquid ſtate, the only means of determining the point of extreme moiſture on the hygrometer, ibid.
3. No fibrous or vaſcular ſubſtance, taken lengthwiſe, proper for the hygrometer, ibid.

4. Light thrown on the march of a chosen hygrometer, by comparing it with the correspondent changes in weight of many hygroscopic substances, ibid.

Reversions. See *Survivorships.*

Ridlington, in Rutland, bed of chalk discovered near, p. 281. Flints found about Ridlington not so black as those of the more southern counties of England, p. 281.

Roxburgh, Dr. on the Chermes Lacca, p. 228.

S.

Selbourn, in Hampshire, abstract of a register of rain kept at, for the year 1789, p. 89.

Series, infinite, observations on, p. 146. See *Infinite Series.*

Shoerl, a substance common both to basaltes and granite, p. 54.

Solutions of problems relating to survivorships, p. 248, 253, 256, 258, 262, 263, 265, 267, 269, 272. Solutions of problems respecting angles, p. 290, 291.

Solutions of propositions relating to infinite series, p. 299, 300, 302, 303, 309, 310, 311, 312, 314.

Squid, a fish, supposed to be the food of the spermaceti whale, p. 44, 46

Stars, nebulous observations on, p. 71. Milky-way consists intirely of stars, ibid.

Sugar, has an affinity with water, and no sensible one with alcohol, p. 11. Water, entering sugar, by the faculty of its capillary pores dissolves it, ibid. Alcohol, entering sugar, by the same means, evaporates, and leaves the sugar sensibly as it was before, ibid.

Sulphur of antimony no ways necessary to the formation of James's powder, p. 360.

Survivorships. On the method of determining, from the real probabilities of life, the values of contingent reversions in which three lives are involved in the survivorship, p. 246.

T.

Tabasheer, chemical experiments on, p. 368. Hydrabad Tabasheer resembles fragments of Cacholong, p. 369. Of a disagreeable earthy taste, resembling that of magnesia, ibid. Experiments on Tabasheer treated with water, p. 370. With vegetable colours, 371. At the fire, ibid. With acids, p. 373. With liquid alkalies, p. 375. With dry alkalies, p. 379. With other fluxes, p. 380.

Tables.

A table of hygroscopic and thermoscopic results, p. 33.

A table of hygroscopic results, on whalebone, quill, and deal, p. 35.

Meteorological journal, principally relating to atmospheric electricity, kept at Knightsbridge, from May 9, 1789, to May 8, 1790, p. 189--210.

Table containing a monthly account of electrical sparks, and of positive and negative electricity, as indicated by a pith-ball electrometer, and sometimes by only flaxen threads without balls to them, p. 211.

Table

Table ſhewing the diſtance anſwering to any angle from one minute to one degree, which is ſubtended by an extenſion of a foot, p. 293.

Table ſhewing the diſtance anſwering to any angle from one minute to a degree, which is ſubtended by an extenſion of ſix feet, p. 294.

Tables of ſums of reciprocals, uſeful in the inveſtigation of the ſummation of infinite ſeries, p. 296, 297, 298, 299.

Table of the comparative changes in the weight and in the length of the ſame ſubſtances, by the ſame increaſes of moiſture, correſpondent to the march of the ſlip of whalebone from 5 to 100 of its degrees, p. 409.

Table of experiments on the comparative changes in the weight and the length of the ſame ſubſtance by increaſe of moiſture, p. 411.

Table of the correſpondent marches, by the ſame increaſes of moiſture, of different threads, or vegetable and animal fibrous ſubſtances taken lengthwiſe, p. 413.

Table of the correſpondent marches of ſlips, or of fibrous vegetable and animal ſubſtances taken acroſs the fibres, and of ſuch as have no ſenſible fibres, p. 415.

Meteorological journal, kept at the Apartments of the Royal Society, for January App. 1790, p. 2. February, p. 4. March, p. 6. April, p. 8. May, p. 10. June, p. 12. July, p. 14. Auguſt, p. 16. September, p. 18. October, p. 20. November, p. 22. December, p. 24.

Table containing the greateſt, leaſt, and mean heights of the thermometer and barometer; and alſo the mean quantity of rain in inches which fell, for every month in the year 1790, p. 26.

Teleſcope, deſcription of a ſimple micrometer for meaſuring ſmall angles with, p. 283.

Teleſcopical micrometers. See *Micrometer*.

Tennant, Smithſon, on the decompoſition of fixed air, p. 182.

Thermometer, abſtract of a regiſter of one, kept at Lyndon in Rutlandſhire, for the year 1789, p. 89.

Travelling, on the rate of, as performed by camels, and its application, as a ſcale, to the purpoſes of geography, p. 129. Camels will not permit themſelves to be overloaded, ibid. Their hourly rate, in travelling, aſcertained, p. 137.

V.

Vince, Rev. Mr. on the inveſtigation of infinite ſeries, p. 295.

W.

Waring, Dr. on infinite ſeries, p. 146.

Water, the only ſure immediate means of producing extreme moiſture in hygroſcopic bodies, p. 1, 16. The temperature of water has a ſenſible effect on the expanſion of ivory, p. 15. The maximum of moiſture known by the exceſs of water, p. 17.

Glaſs contains concrete water very imperfectly, p. 401. Water the cauſe of moiſture, p. 402.

Whale, ſpermaceti, ambergris found in, p. 43. Spermaceti whales, when ſtruck, generally void their excrement; but if they do not, ambergris is found in them, p. 47. Ambergris moſt likely to be found in ſickly whales, ibid.

Whalebone hygrometer, deſcription of, p. 419.

Wolds of Yorkſhire, conſiderable quantities of chalk found in, p. 281.

Y.

Yorkſhire, wolds of, conſiderable quantities of chalk found in, p. 281.

FROM THE PRESS OF JOHN NICHOLS.

ERRATA.

Page. Line.

144. 18. *insert* it *after* if

276. 2. *insert* comma *after* lives, *and dele after* are

311. 3. from bottom, *between* &c. *and* $-\frac{1}{2}$ *insert* -1

1, and 2. from bottom, *for* $\frac{8}{3}$ *read* $\frac{5}{3}$

312. 5. *for* $\frac{8}{3}$ *read* $\frac{5}{3}$